

EUR Research Information Portal

Differences in prescribing errors between electronic prescribing and traditional prescribing among medical students

Published in:

British Journal of Clinical Pharmacology

Publication status and date:

E-pub ahead of print: 23/03/2024

DOI (link to publisher):

[10.1111/bcp.16053](https://doi.org/10.1111/bcp.16053)

Document Version

Publisher's PDF, also known as Version of record

Document License/Available under:

CC BY-NC-ND

Citation for the published version (APA):

El Abdouni, S., Kalfsvel, L. S., Rietdijk, W. J. R., Van der Kuy, H., & van Rosse, F. (2024). Differences in prescribing errors between electronic prescribing and traditional prescribing among medical students: A randomized pilot study. *British Journal of Clinical Pharmacology*. Advance online publication. <https://doi.org/10.1111/bcp.16053>

[Link to publication on the EUR Research Information Portal](#)

Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.

In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: openaccess.library@eur.nl. Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

ORIGINAL ARTICLE

Differences in prescribing errors between electronic prescribing and traditional prescribing among medical students: A randomized pilot study

Samir El Abdouni  | Laura S. Kalfsvel  | Wim J. R. Rietdijk |
Hugo Van der Kuy  | Floor van Rosse

Department of Hospital Pharmacy, Erasmus MC, University Medical Center, Rotterdam, The Netherlands

Correspondence

Samir El Abdouni, Department of Hospital Pharmacy, Erasmus MC, University Medical Center, Dr. Molewaterplein 40, 3015GD Rotterdam, The Netherlands.
Email: s.elabdouni@erasmusmc.nl

Aims: This randomized controlled pilot study aimed to assess the differences in the frequency, type and severity of prescribing errors made by medical students when assessed in an electronic (e-)prescribing system compared to a traditional prescribing method (e.g., typing out a prescription).

Methods: Fourth year medical students in the period of 1 November to 31 July 2023, were asked to participate in this single-centre prospective, randomized, controlled intervention study. Participants performed a prescribing assessment in either an e-prescribing system (intervention group) or in a more traditional prescribing platform (control group). The prescriptions were checked for errors, graded and categorized. Differences in prescribing errors, error categories and severity were analysed.

Results: Out of 334 students, 84 participated in the study. Nearly all participants (98.8%) made 1 or more prescribing errors, primarily involving inadequate information errors. In the intervention group, more participants made prescribing errors involving the prescribed amount (71.4 vs. 19.0%; $P < .01$), but fewer involving administrative errors (2.4 vs. 19.0%; $P = .03$). Prescribing-method-specific errors were identified in 4.8 and 40.5% of the intervention and control group, respectively, with significant differences in overlapping errors as well.

Conclusion: This pilot study shows the importance of training e-prescribing competencies in medical curricula, in addition to traditional prescribing methods. It identifies prescribing-method-specific prescribing errors and emphasizes the need for further research to define e-prescribing competencies. Additionally, the need for an accessible real-life-like e-prescribing environment tailored to educators and students is essential for effective learning and incorporation of e-prescribing into medical curricula.

KEYWORDS

e-prescribing, medical education, prescribing, prescribing errors

F. van Rosse

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *British Journal of Clinical Pharmacology* published by John Wiley & Sons Ltd on behalf of British Pharmacological Society.

1 | INTRODUCTION

Prescribing errors are a known cause of negative patient health outcomes, including decreased health-related quality of life, drug-related hospital admissions and death.¹⁻⁷ For this reason, it is important to reduce the number of prescribing errors. Junior doctors are more likely to make prescribing errors^{8,9} and for a large part feel to have been inadequately prepared by their medical curriculum.¹⁰⁻¹⁴ This incentivizes improvement of the prescribing curriculum.

One way of achieving this, is through practicing and assessing in electronic (e-)learning environments. Earlier studies have shown the benefit of practicing and assessing in e-learning environments.¹⁵⁻¹⁷ For example, Kalfsvell *et al.* have shown that students who practice longer and more frequently in the web-based learning environment Pscribe are more likely to pass their clinical pharmacotherapy (CPT) assessments at the first attempt.¹⁵

When addressing the characteristics of safe and rational prescribing, the concept of competencies frequently arises. While some categorize knowledge, skill and attitude as individual *prescribing competencies*, and others prescribing itself as a competency, it is evident that all these elements are fundamental in achieving actual prescribing competence.^{10,11,18} In this context, we adopt the former definition with knowledge, skill and attitude being individual prescribing competencies necessary for achieving prescribing competence (Figure 1).

Despite substantial research identifying essential prescribing competencies and medical students' competency levels,^{10,11,15,18-21} e-prescribing and its distinct competencies have evaded scholarly attention. In their study, Bakkum *et al.* provided an overview of the digital education resources used in European CPT education in 2019.¹⁶ Despite its widespread use, only a few pertained to the training or assessment of e-prescribing, through the use of (copies) of the electronic prescribing system (EPS)²¹ or a prescribing simulator. This oversight is concerning, given that—as in many other countries—e-prescribing has been the standard of practice in the Netherlands for close to a decade.^{22,23}

We hypothesize that distinct competencies may be required for e-prescribing (e.g., navigating the EPS and patient and drug selection) and *traditional* prescribing (e.g., adhering to the correct format), resulting in prescribing method-specific errors. The latter being supported by studies performed on e-prescribing errors in (clinical) practice, which have shown the existence of various types of e-prescribing specific errors.²⁴⁻²⁷

In light of this, our study aims to assess the differences in the frequency, type and severity of prescribing errors made by medical students in an EPS compared to a more traditional prescribing method. Our objective is addressing the gap regarding the distinctions between e-prescribing and traditional prescribing and offer recommendations for improving the prescribing education curriculum through harmonizing it with the contemporary practices.

What is already known about this subject

- Little is known about e-prescribing competencies.
- Prescribing education may not sufficiently prepare medical students for the future responsibilities.

What this study adds

- On top of differences in frequency in overlapping prescribing error types, medical students make distinct prescribing method-specific errors.
- Both e-prescribing and traditional prescribing competencies should be addressed in medical curricula.
- A more integrated approach to prescribing education in medical curricula, encompassing both prescribing method-specific and non-prescribing method-specific competencies.

2 | METHODS

2.1 | Setting

In the Netherlands, medical education is organized through a 6-year curriculum consisting of a 3-year bachelor and 3-year master's phase. During the bachelor in the Erasmus MC University Medical Center, (clinical) pharmacotherapy education is focused mostly on knowledge, whereas in the master's phase students it is focused on applying this knowledge. Specifically, prescribing education is taught in this phase. This education is interlaced in the curriculum and is primarily organized by the hospital pharmacy, and classes are generally taught by pharmacists and clinical pharmacologists (in-training). On top of the classes, students have access to various (non)mandatory educational materials, including e-learning modules and podcasts on pharmacology and pharmacotherapy.

In the first year of the master's phase, students are taught how to write a paper prescription, how to use the World Health Organization (WHO) 6-step model for rational prescribing²⁸ and perform a formative skill-based prescribing assessment. The latter consists of 6 knowledge-based questions and 2 cases for which they prescribe a predefined drug. This formative assessment is performed in Pscribe,²⁹ an online e-learning platform (see Section 2.4). Here, students are presented with a case description, after which they prescribe according to the WHO 6-step model (see Figure S1). In the second year, students are required to perform the Dutch National Pharmacotherapy assessment,³⁰ a knowledge-based summative assessment. This is also the year in which students receive a 1-h practical class in which they




FIGURE 1 Prescribing competencies.

practice e-prescribing in a real-life-like setting, using the sandbox of the institution's EPS. In the final year, students perform a summative skill-based prescribing assessment in Pscribe, in a fashion similar to the formative assessment. Here, the students prescribe drugs for 4 junior-doctor level cases. In 1 of the cases, the students are instructed to use the WHO 6-step model to choose and prescribe the drug. In the remaining 3 cases, the drugs are predefined, placing emphasis on prescribing the correct form, dosage and dispensing quantity. Except for the 1-h e-prescribing practical, all education materials and assessments related to prescribing are located within the e-learning module Pscribe. The Medical Ethics Committee of the Erasmus MC determined that the Medical Research Involving Human Subjects Act (WMO) was not applicable to this study and approved the study (MEC-2022-0681). All participants provided written informed consent prior to the assessment. Upon registering in Pscribe, students consented to their data's use for research purposes. Thus, no additional consent was required for comparisons between participants and nonparticipants.

2.2 | Population and study procedure

In this prospective, randomized, controlled pilot intervention study, we included medical students in the first year of their master's phase from 1 November 2022 to 31 July 2023. During a mandatory class on medication safety, students were introduced to the study. At this stage, the students had already received some prescribing education and had completed the formative prescribing assessment 2 weeks earlier. Those willing to participate were invited to perform an additional prescribing assessment, either in an EPS (intervention group) or in Pscribe (control group), based on 1:1 randomization, respectively. This assessment took place during the week following the introduction and was conducted in a group setting in a designated classroom as the

examination area. Randomization took place at the time of the study by order of arrival to the computer lab. Every other student was assigned to either the intervention group (assessment in the EPS) or the control group (assessment in Pscribe).

2.3 | The prescribing assessment

For the prescribing assessment used in this study, the participants were asked to prescribe a drug for 2 distinct patients. The 2 cases were identical for both the intervention and control groups. The first case involved a 7-year-old child diagnosed with allergic rhinitis, requiring a prescription of cetirizine. The second case involved a woman seeking to switch from an intrauterine device to an oral contraceptive, prompting the prescription of ethinylloestradiol/levonorgestrel. A team of 3 teachers carefully designed these cases based on their experience, aiming for a balance between the complexity and the potential for prescribing errors. As with the formative and summative assessments, students were allowed to use references but not do discuss. An invigilator was present throughout each assessment.

Considering that the students had not had their EPS practical at the time of intervention, they received a short class on e-prescribing prior to it. This class consisted of a 5-min presentation on the basics of using an EPS (e.g., navigating the EPS and prescribing a drug). Additionally, we asked the participants in the intervention group to record the starting and ending time of each assessment case, as this is automatically tracked for the control group in Pscribe.

After each assessment, the prescriptions were extracted from the EPS and Pscribe and checked for prescribing errors twice by a teacher (S.E.). The prescriptions were graded using a rubric (see Table S1), with a maximum achievable score of 9 points. The rubric was decided upon through a series of focus group sessions with pharmacotherapy

teachers, ultimately leading to consensus. The errors were subsequently categorized by type (see Table S2) and classified by severity (Table S3), in a fashion similar to our previous studies.^{21,31} Additionally, we distinguished between prescribing errors that were exclusively made by a particular group and those that overlapped between the groups (*overlapping prescribing errors*). Within the first category, we further classified errors as either specific to a prescribing method or not.

A second teacher (L.K.) double-checked a 10% sample of the prescriptions. All uncertainties or discrepancies, stemming from the double check or otherwise, were discussed by S.E., L.K. and F.R. until consensus was achieved. The baseline characteristics for all participants were extracted from Pscribe, as each participant had an active account on the platform. All data were entered into a Castor database and all participants received personal feedback on their prescriptions.

2.4 | Prescribing platforms

To perform the assessment, the intervention group used the HiX sandbox (Chipsoft, Amsterdam, The Netherlands). The version of HiX used in our institution is a clinical electronic health record, integrated with an EPS. The sandbox EPS is a copy of the institution's EPS and contains both anonymized and test patients. In this environment, which is primarily used for testing updates, no action reaches actual patients. Within this copy of the EPS, the participants were able to prescribe as they would in actual practice.

The control group performed the prescribing assessment in Pscribe, an online e-learning platform that is currently used for the prescribing education and assessment. Here, participants type out a prescription in a template. We classify this as a traditional prescribing method because, although it is generated digitally, it still involves the manual transcribing of prescriptions, resembling the traditional act of writing. Furthermore, it lacks features associated with e-prescribing, such as EPS navigation, patient selection and structured data entry. Examples are provided in Figures S2–S4.

2.5 | Statistical analysis

As this was a pilot study, we did not perform a formal sample size calculation. We aimed to include a total of 180 participants, drawn from 3 cohorts of students. If this sample size was not reached, we would incorporate 1 more cohort. We analysed the data in 3 steps. Firstly, the baseline characteristics of the participants and the assessment metrics are described. Continuous variables are presented as means and standard deviation or as medians and interquartile ranges depending on the normality of a variable. Differences between 2 groups were examined using an independent sample *t*-test or Mann–Whitney *U* test for normally or non-normally distributed variables. Normality was assessed by the Shapiro–Wilk test. Secondly, the primary endpoint was the difference in prevalence of prescribing error types

between the intervention and control groups, presented as numbers and percentages. When there were at least 5 observations per cell, between-group differences were compared using the Chi-square test. If not, the Fisher's exact test was used. Multiple-group comparison was done using the Fisher's exact test. Thirdly, the secondary endpoints were the prevalence of prescribing method-specific errors, differences in the prevalence of prescribing error subtypes and the difference in prevalence of prescribing error severities, which were presented and compared similar to the primary endpoint. A *P*-value <.05 was considered significant. No imputation was used in case of missing values. All the collected data were analysed using Python (Python Software Foundation. Python Language Reference, version 3.10.4). The data was visualized using the Plotly library (Plotly Technologies Inc., version 5.7.0, Montréal, Canada).

3 | RESULTS

Out of the 334 students that were asked to participate in this study from November 2022 to July 2023, 84 (25.1%) were included. As we did not reach the aimed sample size by the third cohort, we included 1 more. No statistically significant differences were found in the grades for the formative test between the participants and nonparticipants ($P = .14$; see Table S4). An overview of the inclusion and analysed prescriptions is shown in Figure 2. Sixty-two (73.8%) of the included participants were female and 46 (54.8%) participants had a passing grade for the formative assessment (see Table S5). No significant differences were found in sex or formative assessment grade between the intervention and control groups.

3.1 | Assessment metrics

An overview of assessment related results is shown in Table 1. No statistically significant differences were found between the intervention and control groups in mean time spent on the assessment, the median number of points achieved and the number of participants with at least 1 prescribing error. The intervention group, however, had significantly more participants with at least 1 prescribing error in both cases (95.2 vs. 59.5%). When excluding errors related to omission of the bodyweight of a child, which should be explicitly documented on a prescription and an error made by every participant in the intervention, no such difference is found.

3.2 | Prescribing error types

As shown in Figure 3, most prescribing errors were made in the category Inadequate information (e.g., wrong or missing usage instructions), with 100.0 and 88.1% ($P = .06$) of the participants making at least 1 prescribing error in this category for the intervention and control group, respectively. Significantly more participants in the intervention group made at least 1 prescribing error involving the

FIGURE 2 Included participants and analysed prescriptions.^a Two participants were excluded due to a technical problem with extracting the prescriptions.^b One participant did not make a prescription for the second case of the assessment.




TABLE 1 Assessment metrics and prescribing errors.

	Intervention (n = 42)	Control (n = 42)	P
Mean duration (min) per participant in:			
Both cases	10.3 ± 2.9	11.5 ± 3.2 ^a	.10
Mean difference between Case 1 and Case 2 (min) ^b	2.6 ± 4.5 ^a	2.9 ± 4.1 ^a	.82
Median number of points ^c per participant in:			
Both cases	15.0 (13.3–16.0)	14.0 (11.0–17.0)	.87
Case 1	7.5 (6.3–8.0)	7.0 (5.0–9.0) ^a	.79
Case 2	8.0 (7.0–8.0)	8.0 (7.0–8.0)	.82
Number of participants with an error in:			
At least 1 case	42 (100.0)	39 (92.9)	.24
Both cases	40 (95.2)	25 (59.5)	<.01
Number of participants with an error (excluding missing weight errors) in:			
At least 1 case	41 (97.6)	37 (88.1)	.20
Both cases	22 (52.4)	22 (52.4)	>.99
Mean number of errors per participant in:			
Both cases	2.9 ± 0.7	2.9 ± 1.8	.95
Case 1	1.6 ± 0.6	2.0 ± 1.1	.07
Case 2	1.4 ± 0.5	1.6 ± 0.7	.07

Note: Values are expressed as a mean ± standard deviation, median (interquartile range) or n (%).

^a1 missing value.

^bThe difference was calculated by subtracting the time spent on Case 1 by the time spent on Case 2.

^cThe participants could receive a maximum of 9 points per case.




FIGURE 3 Forest plot of relative risk for prescribing errors between intervention and control.

prescribed amount (71.4 vs. 19.0%; $P < .01$) and less participants made at least 1 prescribing error involving administrative errors (2.4 vs. 19.0%; $P = .03$).

3.3 | Prescribing method-specific prescribing error types

In Table 2 we present all the prescribing errors made, stratified by subcategory. We identified 4 prescribing error subcategories made exclusively by the intervention group and 5 made exclusively by the control group. Among these errors, 2 were specific to the intervention group (EPS) and 3 in the control group (non-EPS). Overall, 4.8% of the participants made at least 1 EPS-specific prescribing error and 40.5% at least 1 non-EPS-specific prescribing error. The nonmethod-specific prescribing distinct errors pertained to errors that theoretically could have been made using either method.

3.4 | Overlapping prescribing error types

Among the overlapping prescribing errors (see Table 2), we found significant differences between the intervention and control group in errors related to the omission of the current bodyweight (100 vs. 57.1%, $P < .01$), errors related to a too high dispensing amount for the necessary treatment (21.4 vs. 2.4%, $P = .02$), errors related to a too high dispensing amount for newly started chronic drugs (23.8 vs. 4.8%, $P = .03$).

3.5 | Prescribing error severity

As shown in Table 3, significantly more participants in the intervention group made at least 1 B-class (100.0 vs. 57.1%; $P < .01$) and at least 1 D-class error (83.3 vs. 54.8%; $P = .01$) compared to the control group. Furthermore, when looking at the most severe prescribing error per participant, we find a significant difference in D-class errors as well.

Upon closer investigation, we found that, except for a single error in the control group, all B-class errors involved omitting the child's

bodyweight from the prescription. Most D-class errors involved missing usage instructions for both the intervention (88.6%) and control (91.3%) groups. Specifically, the omission of any instructions pertaining when to start an oral contraceptive and additional contraceptive measures. Furthermore, except for 1 participant in each group, none of the D-class errors were prescribing method-specific.

A detailed overview of the individual prescribing error (sub)categories corresponding with the severities is shown in Table S6.

4 | DISCUSSION

In our pilot study, we aimed to assess the differences in the frequency, type and severity of prescribing errors made by medical students when assessed in an EPS compared to a traditional prescribing method (e.g., typing out a prescription). We found that nearly all participants, both in the intervention and the control group, made 1 or more prescribing error (96.4%), with the majority of them making errors related to inadequate information in both the intervention (100.0%) and control (88.1%) groups, in line with the findings from earlier studies.²¹

Although no significant differences were found in the number of errors, we found that in the intervention group fewer participants made administrative errors (2.4 vs. 19.0%), but more participants made errors related to the prescribed amount (71.4 vs. 19.0%) when compared to the control group. The importance of integrating e-prescribing into the medical curriculum is apparent, as we identified EPS-specific prescribing errors. The required prescribing competencies for avoiding these errors are—as of now—not being taught sufficiently. For instance, the error involving the prescription of an insufficient amount due to an erroneous end-time highlights the need to teach that fields within an EPS are interconnected and can influence the final prescription. Entering data in a wrong field, such as specifying the dosage in the notes section, can lead to potentially harmful situations. Especially considering that—in the Dutch context—most EPSs also include a clinical decision support-system that relies on accurate data entry. Neglecting this can compromise patient safety, particularly when prescribers expect the system to intervene when necessary. In this study, participants in the intervention group would have received an alert when prescribing too high a dose.

TABLE 2 Prescribing errors.

Error category	Subcategory	Intervention (n = 42)	Control (n = 42)	P	Distinct?	Specific to
Administrative error	Value entered in wrong field (EPS) ^a	1 (2.4)	0 (0)	n/a ^f	Y	EPS
	Missing patient information (excl. bodyweight) ^b	0 (0)	8 (19)	n/a ^f	Y	Non-EPS
Inadequate information	No/incomplete dosage ^c	0 (0)	1 (2.4)	>.99	Y	-
	Wrong usage instructions ^d	0 (0)	2 (4.8)	.49	Y	-
	No concentration or dosage stated ^e	0 (0)	12 (28.6)	n/a ^f	Y	Non-EPS
	Wrong dose/concentration (combination) ^f	0 (0)	1 (2.4)	n/a ^f	Y	Non-EPS
	Confusing information ^g	3 (7.1)	10 (23.8)	.07	N	-
	Missing usage instructions ^h	37 (88.1)	31 (73.8)	.16	N	-
	No amount to supply stated ⁱ	1 (2.4)	1 (2.4)	>.99	N	-
Wrong drug dose	No weight of child ^j	42 (100)	24 (57.1)	<.01	N	-
	Dose too high ^k	1 (2.4)	0 (0)	>.99	Y	-
Wrong prescribed amount	Dose too low ^l	7 (16.7)	7 (16.7)	>.99	N	-
	Too little due to erroneous end datetime ^m	1 (2.4)	0 (0)	n/a ^f	Y	EPS
	Insufficient prescribed to finish treatment ⁿ	1 (2.4)	0 (0)	>.99	Y	-
	Insufficient prescribed which makes the prescription patient-unfriendly ^o	10 (23.8)	5 (11.9)	.25	N	-
	Too much prescribed for necessary treatment ^p	9 (21.4)	1 (2.4)	.02	N	-
	Too much prescribed for newly started chronic drugs ^q	10 (23.8)	2 (4.8)	.03	N	-

Abbreviation: EPS, electronic prescribing system.

^aFor example, dose entered in the notes-field as opposed to the dose-field.

^bOmission of relevant patient information (e.g., name, date of birth or address).

^cFor example, cetirizine 1 mg/mL 2 times a day as opposed to cetirizine 1 mg/mL oral liquid 5 mg 2 times a day.

^dFor example, the instruction take with a glass of water in the case of a cetirizine oral liquid.

^eFor example, Cetirizine oral liquid as opposed to Cetirizine 1 mg/mL oral liquid.

^fFor example, ethinylloestradiol/levonorgestrel 10/150 microg. tablet as opposed to ethinylloestradiol/levonorgestrel 30/150 microg. tablet.

^gFor example, the presence of information related to another therapeutic indication or side-effects.

^hFor example, the absence of instructions pertaining to when to start taking an oral contraceptive and/or until when to take additional contraceptive measures.

ⁱOmission of the amount to dispense (e.g., 200 mL or 63 tablets).

^jOmission of the child's current weight, which should be explicitly stated in the prescription.

^kFor example, 10 mg 2 times a day as opposed to 5 mg 2 times a day.

^lFor example, 5 mg 2 times a day as opposed to 10 mg 2 times a day.

^mFor example, by entering a therapy end date that is too soon, the electronic prescribing system calculates the wrong amount to dispense.

ⁿFor example, prescribing an amount only sufficient for 1–2 days.

^oFor example, prescribing 63 tablets of a contraceptive as opposed to 90 tablets for a daily dosage of one tablet, when explicitly not prescribing a stopping week.

^pFor example, 90 tablets of an oral contraceptive, as opposed to 63 tablets, for 3 months when a stop week was prescribed.

^qFor example, prescribing 900 mL of cetirizine oral liquid where 200 mL would have sufficed.

^rNot applicable to prescribing method-specific errors.

TABLE 3 Prescribing error severities.

Severity class	Number of participants per error class ^a			Number of participants by most severe error per participant ^b		
	Intervention (n = 42)	Control (n = 42)	P-value	Intervention (n = 42)	Control (n = 42)	P-value
B	42 (100.0)	24 (57.1)	<.01	1 (2.4)	2 (4.9)	>.99
C	31 (73.8)	30 (71.4)	>.99	6 (14.3)	13 (31.0)	0.12
D	35 (83.3)	23 (54.8)	.01	35 (83.3)	23 (54.8)	.01

Note: Values are presented as n (%). The percentages represent the proportion of the entire group. B: Error occurred but would not have reached the patient. C: Error would have reached the patient but would not have had the potential to cause harm. D: Error would have reached the patient and would have had required additional monitoring to confirm that it resulted in no harm and/or would have required intervention to preclude harm.

^aThe percentages exceed 100%, as some participants made multiple prescribing errors of various severities.

^bThe total percentage may not add up to 100% as not all participants made a prescribing error.

Interestingly, only a single student made this error in the intervention group, but none in the control group.

Furthermore, we found significant differences in the overlapping prescribing errors between the 2 groups, particularly in errors related to the prescribed amount and those related to inadequate information. Concerning errors related to the prescribed amounts, we found that errors related to overprescribing (e.g., 900 mL of cetirizine oral liquid instead of the required 200 mL) were significantly more common in the intervention group (21.4 vs. 2.4%). This results from omitting the specific quantity or duration, which—in our EPS—defaults to a 3-month supply. Errors related to underprescribing for the necessary treatment (only enough cetirizine oral liquid to last 2 days) exclusively occurred in the intervention group, although this was a single occurrence. We found the opposite for inadequate information errors, where there was less variety in the subcategories between the 2 groups. Nonetheless, we found a significant difference in the number of participants that omitted the bodyweight on a prescription for a child between the intervention (100%) and the control group (57.1%). It is remarkable that not 1 participant in the intervention group entered—or at least asked a question about—the bodyweight in the prescription. It is possible that due to the novelty of prescribing in an EPS, this aspect was simply forgotten. In the control group, errors related to confusing information were more common (23.8 vs. 7.1%), albeit not statistically significant. This difference may be attributed to the limitations of the EPS, which imposes constraints on the amount of information that can be entered into the notes section. This may hinder students from copying and pasting irrelevant information, potentially leading to confusion. This phenomenon is something we do not only observe in the control group, but also recognize from assessments performed by medical students as part of the current curriculum. Additionally, we found that participants in the intervention group (83.3%) made significantly more severe errors than those in the control group (54.8%). The fact that the vast majority of these errors were not prescribing-method specific further emphasizes the need for integration of e-prescribing into the medical curriculum.

Nonetheless, our study also revealed the existence of non-EPS-specific prescribing errors, such as omitting essential patient information or the drug concentration or strength, which does not occur in an EPS where drug details are part of the description of the selectable drugs. These findings underscore the importance of teaching medical students how to write traditional paper prescriptions, making them aware of the differences between the prescribing methods. Despite e-prescribing being the current practice, situations may arise where physicians must resort to traditional prescribing methods, such as during technical failures or emergencies.

In *A Competency Framework for all Prescribers*, developed by the Royal Pharmaceutical Society, prescribing is presented as a singular competency, supported by several statements.¹⁵ Two of these statements refer to e-prescribing and the use of EPSs, namely, that a prescriber “electronically generates and/or writes legible, unambiguous and complete prescriptions which meet legal requirements” and “effectively uses the systems necessary to prescribe drugs”. No other explicit references to e-prescribing related competencies were found. Nonetheless, to effectively incorporate e-prescribing competencies into our medical curricula, the distinct competencies need to be identified, properly defined and incorporated into prescribing frameworks. This requires further research and could be achieved extracting competencies from prescribing errors, as shown in Figure 4.

To our knowledge, there are currently no dedicated real-life-like e-prescribing environments available for medical students to hone their e-prescribing competencies. Utilizing the actual EPS of an institution is challenging, primarily due to patient safety concerns and the necessity for direct physician supervision. In practice, we see that students are allowed to prescribe in the institution's EPS during their clerkships, with the requirement for subsequent co-signing by a physician. However, not all systems have this functionality and the learning opportunity is limited to the clerkships, and, while our study utilized a sandboxed version of our institution's EPS, similar to what students experience during their e-prescribing practical, logistical challenges were evident. Prior to the intervention, the students had to be




FIGURE 4 (Electronic-)prescribing competencies.

authorized individually in the EPS by the IT department. During the intervention, issues included participants not having access to the EPS, limited export options and frequent system refreshing. The latter, leading to the loss of the prescriptions of 2 participants. These administrative burdens hinder the educational experience for both students and teachers. Furthermore, the institution's EPS is tailored to the clinical setting, making it less suitable for teaching general (e-)prescribing. As such, there is a pressing need for an accessible real-life-like educational prescribing platform to ensure efficient and effective learning process.

Our study had several limitations, including its small sample size and the single-centre setting, which limit the generalizability of our findings and reduce the statistical power of the study. The low response rate (25%) may have introduced selection bias, as it is likely that the more motivated students participated in the study. Although there was no statistically significant difference in the distribution of the grades on the formative assessment between participants and nonparticipants, twice as many participants received a good grade. Moreover, we only included 46.7% of our intended inclusions. One contributing factor may have been the scheduling constraints, as the proposed timeslots for assessments may not have aligned with the students' availability. The limited number of assessment cases offered us valuable insights, while a larger variety of cases might offer a more comprehensive evaluation. For future research, it may be beneficial to offer an online assessment option with the freedom for participants to complete it at their convenience, which may increase participation rates. Regarding the checking, although all discrepancies were discussed until consensus was achieved, we did not calculate the interrater reliability. Additionally, we relied on the sandboxed version of our institution's EPS, which may not fully replicate the diversity of EPSs used in real clinical practice. Finally, it is important to acknowledge that EPSs and practices can vary significantly between countries, which should be considered when interpreting our results outside of the Dutch healthcare setting.

The results of our study emphasize the need for re-evaluating prescribing education within our medical curricula. It is essential to incorporate prescribing competencies tailored to both specific prescribing methods as well as those applicable. To achieve this, a follow-up to this pilot study is required, involving further research to identify these competencies across a larger participant population with a wider array of prescriptions. In parallel, the development of an accessible educational prescribing platform, is imperative. Meanwhile, educators should try to keep in mind the intended prescribing competencies when designing prescribing assessments.

5 | CONCLUSIONS

This study shows the importance of training e-prescribing competencies in medical curricula, in addition to traditional prescribing methods. It identifies prescribing-method-specific prescribing errors and emphasizes the need for further research to define e-prescribing competencies. Additionally, the need for an accessible real-life-like

e-prescribing environment tailored to educators and students is essential for effective learning and incorporation of e-prescribing into medical curricula.

AUTHOR CONTRIBUTIONS

Samir El Abdouni, Laura S. Kalfsvel, Wim J.R. Rietdijk and Floor van Rosse designed the study. Data classification was done by Samir El Abdouni, Laura S. Kalfsvel and Floor van Rosse. Samir El Abdouni and Laura S. Kalfsvel processed the data. Samir El Abdouni performed the analysis and drafted the manuscript. All authors discussed the results and commented on the manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Samir El Abdouni  <https://orcid.org/0000-0002-3873-2202>

Laura S. Kalfsvel  <https://orcid.org/0000-0002-1545-7623>

Hugo Van der Kuy  <https://orcid.org/0000-0002-7128-8801>

REFERENCES

- Pont L, Alhawassi T, Bajorek B, Krass I. A systematic review of the prevalence and risk factors for adverse drug reactions in the elderly in the acute care setting. *Clin Interv Aging*. 2014;9:2079-2086. doi:10.2147/cia.s71178
- Beijer HJM, De Blaey CJ. Hospitalisations caused by adverse drug reactions (ADR): a meta-analysis of observational studies. *Pharm World Sci*. 2002;24(2):46-54. doi:10.1023/a:1015570104121
- Tangiisuran B, Graham Davies J, Wright JE, Rajkumar C. Adverse drug reactions in a population of hospitalized very elderly patients. *Drugs Aging*. 2012;29(8):669-679. doi:10.1007/bf03262282
- Oscanoa TJ, Lizaraso F, Carvajal A. Hospital admissions due to adverse drug reactions in the elderly. A meta-analysis. *Eur J Clin Pharmacol*. 2017;73(6):759-770. doi:10.1007/s00228-017-2225-3
- Lazarou J, Pomeranz BH, Corey PN. Incidence of adverse drug reactions in hospitalized patients. *JAMA*. 1998;279(15):1200. doi:10.1001/jama.279.15.1200
- Klopotowska JE, Wierenga PC, Stuijt CCM, et al. Adverse drug events in older hospitalized patients: results and reliability of a comprehensive and structured identification strategy. *PLoS ONE*. 2013;8(8):e71045. doi:10.1371/journal.pone.0071045
- Leendertse AJ, Egberts AC, Stoker LJ, van den Bemt PM, Group HS. Frequency of and risk factors for preventable medication-related hospital admissions in the Netherlands. *Arch Intern Med*. 2008;168(17):1890-1896.
- Ashcroft DM, Lewis PJ, Tully MP, et al. Prevalence, nature, severity and risk factors for prescribing errors in hospital inpatients: prospective study in 20 UK hospitals. *Drug Saf*. 2015;38(9):833-843. doi:10.1007/s40264-015-0320-x
- Ryan C, Ross S, Davey P, et al. Prevalence and causes of prescribing errors: the PRescribing Outcomes for Trainee Doctors Engaged in Clinical Training (PROTECT) study. *PLoS ONE*. 2014;9(1):e79802. doi:10.1371/journal.pone.0079802 PONE-D-13-18651 [pii]
- Brinkman DJ, Tichelaar J, Schutte T, et al. Essential competencies in prescribing: a first European cross-sectional study among 895 final-

- year medical students. *Clin Pharmacol Ther.* 2017;101(2):281-289. doi:10.1002/cpt.521
11. Brinkman DJ, Tichelaar J, Graaf S, Otten RHJ, Richir MC, Van Agtmael MA. Do final-year medical students have sufficient prescribing competencies? A systematic literature review. *Br J Clin Pharmacol.* 2018;84(4):615-635. doi:10.1111/bcp.13491
 12. Geoghegan SE, Clarke E, Byrne D, et al. Preparedness of newly qualified doctors in Ireland for prescribing in clinical practice. *Br J Clin Pharmacol.* 2017;83(8):1826-1834. doi:10.1111/bcp.13273
 13. Han WH, Maxwell SR. Are medical students adequately trained to prescribe at the point of graduation? Views of first year foundation doctors. *Scott Med J.* 2006;51(4):27-32. doi:10.1258/RMSMJ.51.4.27
 14. Heaton A, Webb DJ, Maxwell SR. Undergraduate preparation for prescribing: the views of 2413 UK medical students and recent graduates. *Br J Clin Pharmacol.* 2008;66(1):128-13434. doi:10.1111/j.1365-2125.2008.03197.x
 15. Kalfsvell L, Versmissen J, van Doorn A, van den Broek W, van der Kuy H, van Rosse F. Better performance of medical students on pharmacotherapy knowledge and skills tests is associated with practising with e-learning program P-scribe. *Br J Clin Pharmacol.* 2022;88(3):1334-1346. doi:10.1111/bcp.15077
 16. Bakkum MJ, Tichelaar J, Papaioannidou P, et al. Harmonizing and improving European education in prescribing: an overview of digital educational resources used in clinical pharmacology and therapeutics. *Br J Clin Pharmacol.* 2021;87(3):1001-1011. doi:10.1111/bcp.14453
 17. Gordon M, Chandratilake M, Baker P. Improved junior paediatric prescribing skills after a short e-learning intervention: a randomised controlled trial. *Arch Dis Child.* 2011;96(12):1191-11944. doi:10.1136/archdischild-2011-300577
 18. Royal Pharmaceutical Society. *A Competency Framework for all Prescribers.* Royal Pharmaceutical Society. Accessed 01/09/2023. <https://www.rpharms.com/Portals/0/RPS%20document%20library/Open%20access/Prescribing%20Competency%20Framework/RPS%20English%20Competency%20Framework%203.pdf?ver=mctnrKo4YaJDh2nA8N5G3A%3d%3d>
 19. Liu J, Wong S, Foster G, Holbrook A. Prescribing competency of medical students: National Survey of Medical Education Leaders. *J Popul Ther Clin Pharmacol.* 2018;25(1):e18-e24. doi:10.22374/1710-6222.25.1.2
 20. Mucklow J, Bollington L, Maxwell S. Assessing prescribing competence. *Br J Clin Pharmacol.* 2012;74(4):632-6399. doi:10.1111/j.1365-2125.2011.04151.x
 21. Kalfsvell L, Hoek K, Bethlehem C, et al. How would final-year medical students perform if their skill-based prescription assessment was real life? *Br J Clin Pharmacol.* 2022;88(12):5202-5217. doi:10.1111/bcp.15427
 22. IGJ. *Elektronisch voorschrijven (EVS).* Inspectie Gezondheidszorg en Jeugd. Accessed 01/09/2023. <https://www.igj.nl/onderwerpen/medicatieveiligheid/toezicht-voorschrijven-geneesmiddelen/elektronisch-voorschrijven>
 23. KNMG. In: GGD Nederland GN, KNMP, KNOV, LHV, NHG, NPCF, NVAVG, NVZ, NVZA, OMS, V&VN, Verenso, VGN, VHN, ed. *Richtlijn Elektronisch voorschrijven*; 2013.
 24. Odukoya OK, Stone JA, Chui MA. E-prescribing errors in community pharmacies: exploring consequences and contributing factors. *Int J Med Inform.* 2014;83(6):427-437. doi:10.1016/j.ijmedinf.2014.02.004
 25. Molitor R, Friedman S. Electronic prescription errors in an ambulatory pharmacy. *J Manag Care Pharm.* 2011;17(9):714-7155. doi:10.18553/jmcp.2011.17.9.714
 26. Palchuk MB, Fang EA, Cygielnik JM, et al. An unintended consequence of electronic prescriptions: prevalence and impact of internal discrepancies. *J Am Med Inform Assoc.* 2010;17(4):472-476. doi:10.1136/jamia.2010.003335
 27. Brown CL, Mulcaster HL, Triffitt KL, et al. A systematic review of the types and causes of prescribing errors generated from using computerized provider order entry systems in primary and secondary care. *J Am Med Inform Assoc.* 2017;24(2):432-440. doi:10.1093/jamia/ocw119
 28. de Vries TPGM, Richir MC, Tichelaar J. *Guide to Good Prescribing.* World Health Organization (WHO); 1994.
 29. Pscribe. Accessed 12/06/2023. 2023. <https://www.pscribe.nl/en-GB/Entrance/Home/Index>
 30. Kramers C, Janssen BJ, Knol W, et al. A licence to prescribe. *Br J Clin Pharmacol.* 2017;83(8):1860-1861. doi:10.1111/bcp.13257
 31. Kalfsvell L, Wilkes S, van der Kuy H, et al. Do junior doctors make more prescribing errors than experienced doctors when prescribing electronically using a computerised physician order entry system combined with a clinical decision support system? A cross-sectional study. *Eur J Hosp Pharm.* 2023. doi:10.1136/ejpharm-2023-003859

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: El Abdouni S, Kalfsvell LS, Rietdijk WJR, Van der Kuy H, van Rosse F. Differences in prescribing errors between electronic prescribing and traditional prescribing among medical students: A randomized pilot study. *Br J Clin Pharmacol.* 2024;1-10. doi:10.1111/bcp.16053