



Global access to technologies to support safe and effective inguinal hernia surgery: prospective, international cohort study

National Institute for Health and Care Research (NIHR) Global Health Research Unit on Global Surgery

Correspondence to: Maria Picciochi, National Institute for Health and Care Research (NIHR) Global Health Research Unit on Global Surgery, Institute of Applied Health Research, University of Birmingham, Heritage Building, Mindelsohn Way, Birmingham B15 2TH, UK (e-mail: m.picciochi@bham.ac.uk)

Members of the National Institute for Health and Care Research (NIHR) Global Health Research Unit on Global Surgery are co-authors of this study and are listed under the heading Collaborators.

This paper won the BJS European Hernia Society Competition prize and was presented at the European Hernia Society Conference, Prague, 2024 (<https://ehs2024.org>).

Introduction

Technological advancement is important to improve healthcare quality and safety, especially in surgery¹. For patients with an inguinal hernia, mesh and minimally invasive surgery are the two main technologies that have improved healthcare quality and safety^{2,3}. The use of mesh is proven to reduce recurrence^{4,5}. This avoids the need for further repairs, which are technically more challenging and have a higher risk for patients⁶. The use of minimally invasive surgery has proven advantages in bilateral hernias and in female patients^{2,3} and is recommended in unilateral repair where appropriate expertise is available^{2,3}.

Access to these technologies and the expertise required are not widely or equitably distributed at a global level. As it is the case for other technologies, countries in the Global South have more limited access¹. At the same time, in this part of the globe, there is a higher prevalence and a higher burden of disease associated with inguinal hernias⁷. Several barriers to implementation in the Global South have been identified previously, including costs, distribution, and training^{8,9}. To overcome these, studies reporting the use of mesh based on mosquito net mesh and evaluating training programmes have been conducted^{10,11}. With these efforts and with global investment in new technologies and the expansion of existing technologies, it was expected that there would be an increase in their use in low–middle-income countries. Data assessing this variability have not been collected in a standardized way and are usually reported from single-country or single-region studies^{5,12}. Therefore, identification of areas where improvement is most needed will be key to better inform policymakers.

The overarching aim of this study was to evaluate access to technologies that are relevant to the treatment of inguinal hernia patients to identify the areas where improvement is needed. Therefore, the primary aim of this study was to evaluate the use of mesh and predictors of mesh use in elective inguinal hernia repairs and the secondary aims of this study were to evaluate the use of minimally invasive surgery and predictors of minimally invasive surgery use and to evaluate the

safety associated with the use of mesh and the use of minimally invasive surgery.

Methods

Study design

This was a pre-planned analysis of an international, multicentre, prospective cohort study of patients undergoing inguinal hernia surgery. Routine and anonymized data were collected and no changes in patient care were made. The study protocol is publicly available (globalsurgeryunit.org/clinical-trials-holding-page/hippo) and was registered in ClinicalTrials.gov (NCT05748886). Approvals were obtained by local principal investigators in each hospital taking part, according to local and national regulations. This study is reported in line with STROBE guidelines¹³.

Inclusion and exclusion criteria

Any hospital performing inguinal hernia repair was eligible to take part. Each participating hospital identified consecutive patients undergoing primary inguinal hernia repair as the main procedure during a 4-week inclusion window between 30 January and 21 May 2023. Adult patients, defined as older than 16 years, undergoing elective primary inguinal repair were included. Patients operated on via midline incision or converted to midline incision were excluded, considering the complexity inherent to this approach.

Outcome definitions

The use of mesh in open surgery was defined as the primary outcome and was compared across the different income groups, as defined by the World Bank. The use of minimally invasive surgery and complications at 30 days were secondary outcomes. Minimally invasive surgery included both laparoendoscopic and robotic approaches and was defined as per intention-to-treat, therefore converted surgeries to open were included in this group. Postoperative complications were defined according to

Received: January 31, 2024. **Revised:** April 10, 2024. **Accepted:** June 09, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of BJS Foundation Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

the Clavien–Dindo classification and these data were collected at 30 days after surgery¹⁴. To comprehensively evaluate postoperative complications, surgical-site infection rates and reoperation rates (mapped to surgical approach and use of mesh) were also collected at 30 days after surgery.

Data management

Data were collected and stored online using a secure server running the Research Electronic Data Capture (REDCap) web application¹⁵. The service was managed by the Global Surgery REDCap system hosted at the University of Birmingham, Birmingham, UK. Its security was governed by the policies of the University of Birmingham. Each collaborator involved in data collection was identified and registered by the hospital lead and received personal login details. This allowed secure data entry and storage in REDCap.

Data validation

The data collection methodology was validated previously, in terms of case ascertainment and data accuracy^{16,17}. The hospital lead had access to the data entered by their team. They were responsible for data accuracy and data completeness collected and uploaded from their site. The data were checked centrally and when there were missing data or invalid data, the hospital lead was contacted to complete and correct the data entered. After this, participating hospitals with data completeness less than 95% were excluded.

Sample size

There was no formal sample size calculation for the analysis proposed and all eligible patients were included. To ensure global generalizability of the results and to justify the resources put into the study, a minimum number of 300 centres contributing patient-level data from 70 countries was estimated, based on previous cohort studies (that is GlobalSurg and COVIDSurg studies)^{16,17}. Assuming an average of 30 patients per centre, a minimum sample size of 10 000 patients was predicted. Assuming that the prevalence of mesh use ranges between 70% and 95%, sample size considerations for building a prediction model showed that approximately 2500 subjects would be required to build a model with 7 predictor variables, a prevalence of 95%, and a C-statistic of 0.7 (see [Table S1](#) for full details)¹⁸. Sample sizes were estimated using the `pmsampsiz` command in Stata, version 18.0 (StataCorp).

Statistical analysis

Data were mapped to country income groups, defined according to the World Bank (low-income countries, lower-middle-income countries, upper-middle income countries, and high-income countries), as their importance in relation to healthcare access, safety, and quality has been widely recognized⁶.

Continuous non-normally distributed hospital-, patient-, and intraoperative-related variables are presented as median (interquartile range (i.q.r.)) values, whereas categorical variables are presented as frequencies and percentages. The use of mesh and minimally invasive surgery are presented as frequencies and rates across income groups. Postoperative complications, surgical-site infection, and reoperation are presented as frequencies and rates across surgical approach and mesh use groups.

Multilevel logistic regression models were used to test factors that could be associated with higher mesh use in open surgery and the use of minimally invasive surgery. Plausible hospital

and clinical factors agreed by the Study Management Group were considered and hospital was included as a random effect. For the above analyses, appropriate model fit diagnostics were checked to confirm that validity and model assumptions were maintained for the data. Categories were collapsed when category event rates were too low to be efficiently included in the model, as was the case for income groups. All statistical analyses were performed using R (R Foundation for Statistical Computing, Vienna, Austria; version 4.0.2). $P < 0.050$ was considered statistically significant.

Results

Included patients

Data were collected from 18 058 patients across 640 centres located in 83 countries. For this study, 14 768 adults undergoing elective primary inguinal hernia repair in 612 centres located in 81 countries were included, as shown in [Fig. 1](#) and [Fig. S1](#). Most of them were operated on in high-income countries (60.4%, 8916 of 14 768).

Included patients had a median age of 60 (i.q.r. 47.0–70.0) years ([Table 1](#)), with an absolute median difference of 10 years between patients operated on in high- and low-income countries. Most patients were male (91.7%, 13 539 of 14 768). Regarding their perioperative risk, most were ASA grade I–II (85.9%, 12 691 of 14 768) and without co-morbidities, which was observed across all income groups. The majority of patients presented with symptomatic hernias (86.2%, 12 729 of 14 768) that were unilateral and with an extension limited to the inguinal region (79.4%, 11 733 of 14 768). Intraoperatively, greater than 90.0% of the operations were classified as clean (14 419 of 14 768). Hernia defect size was variable in all income groups, with defects of 1.5–3 cm being the most reported (40.8%, 6031 of 14 768). Hospitals where these patients were operated on were mostly tertiary-level centres (62.4%, 9126 of 14 768) and their funding was mainly provided by the public sector (86.6%, 12 152 of 14 768) ([Table S2](#)).

Surgical variation and intraoperative outcomes

In all income groups, patients were more commonly operated on by a senior surgeon (71.0%, 10 487 of 14 768) ([Table 2](#)). Most had previous experience of greater than 200 inguinal hernia repairs (53.1%, 7833 of 14 768). There was heterogeneity in the surgical technique chosen to repair inguinal hernia, as shown in [Fig. 2](#). The Lichtenstein technique was used for greater than half of patients in all groups (61.7%, 9117 of 14 768). Of the techniques using a minimally invasive approach, transabdominal preperitoneal repair was twice as commonly used as totally extraperitoneal repair (7.8%, 1155 of 14 768). The use of both decreased from high- to low-income countries. The use of soft tissue repair was more commonly used in low-income countries (28.0%, 178 of 635). More details regarding surgical technique variation are available in [Table S3](#).

Overall, 94.8% of the patients had mesh placed to repair the inguinal hernia (13 995 of 14 768) ([Table 2](#)). When the approach was open, mesh was used in 93.2% of the repairs (13 995 of 14 768) ([Fig. 3](#)). There was a reduction in mesh use from high-income countries (98.9%) to low-income countries (72.1%). In the group of patients where mesh was used, the most frequent type of mesh was permanent synthetic (90.2%, 12 620 of 13 995) and the most common suture used to fix the mesh was non-absorbable (52.5%, 7343 of 13 995).

Less than a quarter of patients underwent minimally invasive surgery (24.8%, 3661 of 14 768) ([Fig. 3](#)).




Fig. 1 Flow chart of included patients

HIC, high-income country; UMIC, upper-middle-income country; LMIC, lower-middle-income country; LIC, low-income country.

Table 1 Preoperative and intraoperative characteristics of adults undergoing elective inguinal hernia repair

	HIC (n = 8916)	UMIC (n = 2575)	LMIC (n = 2642)	LIC (n = 635)	Total (n = 14 768)
Age (years)					
Median (interquartile range)	63.0 (52.0–73.0)	57.0 (45.0–67.0)	52.0 (37.0–63.0)	53.0 (36.0–63.5)	60.0 (47.0–70.0)
Sex					
Male	8094 (90.8)	2356 (91.5)	2500 (94.6)	589 (92.8)	13 539 (91.7)
Female	821 (9.2)	219 (8.5)	142 (5.4)	46 (7.2)	1228 (8.3)
Missing, n	1	0	0	0	1
ASA grade					
I–II	7211 (80.9)	2337 (90.8)	2536 (96.0)	607 (95.6)	12 691 (85.9)
III–V	1645 (18.4)	234 (9.1)	105 (4.0)	21 (3.3)	2005 (13.6)
Not recorded	60 (0.7)	4 (0.2)	1 (0.0)	7 (1.1)	72 (0.5)
Missing, n	0	0	0	0	0
Co-morbidities					
None	6613 (74.2)	1961 (76.2)	2195 (83.1)	526 (82.8)	11 295 (76.5)
One	1713 (19.2)	491 (19.1)	367 (13.9)	100 (15.7)	2671 (18.1)
Two	456 (5.1)	93 (3.6)	72 (2.7)	9 (1.4)	630 (4.3)
Three or more	130 (1.5)	26 (1.0)	5 (0.2)	0	161 (1.1)
Missing, n	4	3	3	0	10
Symptoms					
Asymptomatic	1106 (12.4)	293 (11.4)	580 (22.0)	60 (9.4)	2039 (13.8)
Symptomatic	7810 (87.6)	2282 (88.6)	2062 (78.0)	575 (90.6)	12 729 (86.2)
Missing, n	0	0	0	0	0
Hernia size					
Limited to inguinal region	7576 (85.0)	1977 (76.8)	1698 (64.3)	482 (75.9)	11 733 (79.4)
Limited to scrotum	1263 (14.2)	569 (22.1)	903 (34.2)	147 (23.1)	2882 (19.5)
Extend to mid-thigh or beyond	77 (0.9)	29 (1.1)	41 (1.6)	6 (0.9)	153 (1.0)
Missing, n	0	0	0	0	0
Hernia site					
Bilateral	1362 (15.3)	347 (13.5)	386 (14.6)	59 (9.3)	2154 (14.6)
Unilateral	7554 (84.7)	2228 (86.5)	2256 (85.4)	576 (90.7)	12 614 (85.4)
Missing, n	0	0	0	0	0
Hernia defect size (cm)					
<1.5	1866 (20.9)	456 (17.7)	425 (16.1)	138 (21.7)	2885 (19.5)
1.5–3	3654 (41.0)	1018 (39.5)	1063 (40.2)	296 (46.6)	6031 (40.8)
>3	1804 (20.2)	820 (31.8)	920 (34.8)	175 (27.6)	3719 (25.2)
Not known	1590 (17.8)	281 (10.9)	234 (8.9)	26 (4.1)	2131 (14.4)
Missing, n	2	0	0	0	2
Contamination					
Clean	8858 (99.3)	2383 (92.5)	2546 (96.4)	632 (99.5)	14 419 (97.6)
Clean-contaminated	53 (0.6)	190	93	3 (0.5)	339 (2.3)
Contaminated	4 (0.0)	2 (0.1)	2 (0.1)	0 (0.0)	8 (0.1)
Dirty	1 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	2 (0.0)
Missing, n	0	0	0	0	0

Values are n (%) unless otherwise indicated. HIC, high-income country; UMIC, upper-middle-income country; LMIC, lower-middle-income country; LIC, low-income country.

Table 2 Surgical variation and outcomes across income groups

	HIC (n = 8916)	UMIC (n = 2575)	LMIC (n = 2642)	LIC (n = 635)	Total (n = 14 768)
Primary operator					
Senior surgeon	6568 (73.7)	1868 (72.5)	1653 (62.6)	398 (62.7)	10 487 (71.0)
Trainee surgeon	2338 (26.2)	665 (25.8)	951 (36.0)	232 (36.5)	4186 (28.3)
Non-surgeon	10 (0.1)	42 (1.6)	38 (1.4)	5 (0.8)	95 (0.6)
Missing, n	0	0	0	0	0
Previous experience (number of repaired hernias)					
0–50	1472 (16.5)	523 (20.3)	534 (20.2)	223 (35.1)	2752 (18.7)
51–200	2400 (27.0)	591 (23.0)	1004 (38.0)	172 (27.1)	4167 (28.2)
>200	5030 (56.5)	1461 (56.7)	1102 (41.7)	240 (37.8)	7833 (53.1)
Missing, n	14	0	2	0	16
Surgical approach					
Open	6303 (70.7)	1865 (72.4)	2311 (87.5)	628 (98.9)	11 107 (75.2)
Laparoendoscopic	2397 (26.9)	699 (27.1)	316 (12.0)	6 (0.9)	3418 (23.1)
Robotic	144 (1.6)	4 (0.2)	4 (0.2)	0 (0.0)	152 (1.0)
Converted	72 (0.8)	7 (0.3)	11 (0.4)	1 (0.2)	91 (0.6)
Missing, n	0	0	0	0	0
Use of mesh					
Yes	8842 (99.2)	2465 (95.7)	2231 (84.4)	457 (72.0)	13 995 (94.8)
No	74 (0.8)	110 (4.3)	411 (15.6)	178 (28.0)	773 (5.2)
Missing, n	0	0	0	0	0
Type of mesh used*					
Permanent synthetic	7964 (90.1)	2184 (88.6)	2055 (92.2)	417 (91.2)	12 620 (90.2)
Absorbable synthetic	608 (6.9)	193 (7.8)	155 (7.0)	39 (8.5)	995 (7.1)
Biological	7 (0.1)	9 (0.4)	1 (0.0)	1 (0.2)	18 (0.1)
Composite	261 (3.0)	79 (3.2)	18 (0.8)	0 (0.0)	358 (2.6)
Missing, n	2	0	2	0	4
Suture used to fix the mesh*					
Absorbable	2112 (23.9)	542 (22.0)	373 (16.7)	49 (10.7)	3076 (22.0)
Non-absorbable	3870 (43.8)	1449 (58.8)	1617 (72.5)	407 (89.1)	7343 (52.5)
Glue	641 (7.3)	6 (0.2)	3 (0.1)	0 (0.0)	650 (4.6)
Tackers	872 (9.9)	305 (12.4)	186 (8.3)	0 (0.0)	1363 (9.7)
Not fixed	1345 (15.2)	163 (6.6)	50 (2.2)	1 (0.2)	1559 (11.1)
Missing, n	2	0	2	0	4

Values are n (%) unless otherwise indicated. *Only evaluated in patients undergoing inguinal hernia repair with mesh (n=13 995). HIC, high-income country; UMIC, upper-middle-income country; LMIC, lower-middle-income country; LIC, low-income country.




Fig. 2 Variation in surgical technique across income groups

TAPP, transabdominal preperitoneal repair; TEP, totally extraperitoneal repair; HIC, high-income country; UMIC, upper-middle-income country; LMIC, lower-middle-income country; LIC, low-income country.




Fig. 3 Use of technologies across income groups

*Of the patients undergoing minimally invasive surgery, 99.6% (3648 of 3661) had mesh repair. There were no missing data for surgical approach and mesh use. MIS, minimally invasive surgery; HIC, high-income country; UMIC, upper-middle-income country; LMIC, lower-middle-income country; LIC, low-income country.




Fig. 4 Predictors of mesh use in open surgery

Only patients undergoing open surgery were included in the model ($n=11\,107$). HIC, high-income country; LMIC, low-middle income countries, including upper-middle-, lower-middle-, and low-income countries.

Laparoendoscopic surgery accounted for most of the minimally invasive surgery across all income groups, as shown in Table 2. In general, there was a higher proportion of

patients operated on by senior surgeons in laparoendoscopic surgery (87.6%, 3074 of 3508) with a higher previous experience, as shown in Table S4.




Fig. 5 Predictors of use of minimally invasive surgery

MIS, minimally invasive surgery; HIC, high-income country; LMIC, low-middle income countries, including upper-middle-, lower-middle-, and low-income countries.

Table 3 Variation in complications, surgical-site infection, and reoperation at 30 days with regard to surgical approach and use of mesh

	Complications at 30 days	Surgical-site infection at 30 days	Reoperation at 30 days	Total
Surgical approach*				
Open surgery	1518 (13.7)	397 (3.6)	45 (0.4)	11 107
Minimally invasive surgery	415 (11.3)	48 (1.3)	30 (0.8)	3661
Missing, n	0	0	0	0
Use of mesh				
Yes	1828 (13.1)	408 (2.9)	67 (0.5)	13 995
No	105 (13.6)	37 (4.8)	8 (1.0)	773
Missing, n	0	0	0	0

Values are n (%) unless otherwise indicated. *Surgical approach classified as intention-to-treat. Minimally invasive surgery includes laparoendoscopic, robotic, and converted surgeries.

Predictors of access to technologies

Patients undergoing open surgery where mesh was not used were younger (median age of 49.0 years *versus* 61.0 years), had fewer co-morbidities, and had larger hernias, as shown in [Table S5](#). However, in the adjusted analysis, being operated on in a low-middle-income country was associated with lower mesh use (adjusted OR 0.02 (95% c.i. 0.01 to 0.06); $P < 0.001$) ([Fig. 4](#)). Being female was the only other factor that was associated with lower use of mesh in open surgery (adjusted OR 0.53 (95% c.i. 0.35 to 0.81); $P = 0.004$). Of the other factors tested, none had a significant association with use of mesh.

Patients undergoing hernia repair in low-middle-income countries was associated with lower odds of minimally invasive surgery use (adjusted OR 0.11 (95% c.i. 0.07 to 0.18); $P < 0.001$) ([Fig. 5](#)). Having an inguinal hernia limited to the scrotum or that

extended to the mid-thigh or beyond was associated with lower odds of minimally invasive surgery. Of the hospital factors tested, being operated on in a private hospital (adjusted OR 9.20 (95% c.i. 4.84 to 17.51); $P < 0.001$) and in a tertiary-level hospital (adjusted OR 3.05 (95% c.i. 1.43 to 6.53); $P = 0.004$) were both associated with higher odds of use of minimally invasive surgery. Of the patient factors tested, having a bilateral hernia repair was associated with higher use of minimally invasive surgery (adjusted OR 7.87 (95% c.i. 6.82 to 9.09); $P < 0.001$).

Postoperative complications

The postoperative complication rate was 13.7% and most of the postoperative complications were minor, i.e. Clavien–Dindo grade I–II (93.5%, 1808 of 1933) ([Table S6](#)). Patients undergoing minimally invasive surgery had a postoperative complication rate

of 11.3% (415 of 3661) (Table 3). Patients where mesh was used had a postoperative complication rate of 13.1% (1828 of 13 995). Overall, at 30 days, the surgical-site infection rate was 3.0% (445 of 14 768) and the reoperation rate was 0.5% (75 of 14 768).

Discussion

The principal finding of this study is the lack of access to mesh observed in low- and middle-income countries. This was shown in open repair, as well as in all hernia repairs included, regardless of the approach. With regard to open surgery, having the hernia repair in low–middle-income countries was the most important factor found to be associated with lower use of mesh. Lower use of mesh not only has a direct impact on patients, who will have a higher risk of hernia recurrence⁴, but also demonstrates that access to mesh technology is limited.

This study also shows low use of minimally invasive surgery across all income groups. Overall, less than a quarter of patients were operated on using minimally invasive surgery and, when used, laparoendoscopic-based techniques were preferred. This was even lower in low–middle-income countries. However, having the repair in a private or tertiary-level hospital and having a bilateral hernia were all associated with higher use of minimally invasive surgery.

The data from this study are relevant for developing plans to expand the use of mesh and minimally invasive surgery.

There is a global need to increase access to and training programmes for mesh inguinal hernia repair in low–middle-income countries¹⁹. Mesh is a simple device that has been recommended by international guidelines for the treatment of inguinal hernias² and is recognized as standard practice by several hernia societies globally^{20–22}. Using mesh reduces recurrence rates, avoids further operations, and has been shown to be cost-effective^{4,23}. Therefore, upscaling mesh use in inguinal hernia patients should be a first priority in providing access to more advanced technologies²⁰. Supply chains, training surgical teams, and reducing costs of mesh for patients are all factors that have been identified previously as barriers to access to mesh technology and that could be targeted¹.

Expansion of minimally invasive surgery in well-resourced settings will require expansion of dedicated training programmes and a focus on patients who will benefit most. In settings where expertise is available, minimally invasive surgery is the recommended approach for inguinal hernia repair, according to international guidelines². However, the low use of minimally invasive surgery in this study, even in high-income countries, leads to concerns regarding inherent training challenges and slower learning curves^{24,25}. There is also the potential lack of agreement in the wider general surgical community regarding the clinical benefit outside of selected groups of patients, such as patients with bilateral hernias, patients with recurrent hernias, and female patients²⁶.

There are limitations associated with this study. Representation by low–middle-income countries in this setting was low. Countries with better access to technologies might not have been captured. Also, in the high-income group, there was a lack of representation by centres from countries that are reported to have higher rates of minimally invasive surgery (for example Sweden and Denmark)²⁷ and this might have resulted in a low estimation of minimally invasive surgery use amongst this group. Complication data were only collected at 30 days after surgery, which limits the evaluation of recurrence, which is an important longer-term outcome of hernia repair. However,

there is already good evidence showing higher recurrence rates when mesh is not placed, even in low-income settings^{5,11}.

Future research is still needed. Full understanding of the payment mechanisms available in different countries will help to identify economic barriers to access to mesh technology. National evaluation of payment options and avoidance of out-of-pocket expenses might improve access to mesh and other technologies, by protecting patients from catastrophic expenditure.

This study provides relevant information to policymakers on potential targets to improve access to simple technologies, such as mesh. To achieve medium- to long-term improvement, it will be essential to train surgical teams on site. Partnerships between high-income countries and low–middle-income countries could be useful to co-develop a recognized global training package. Involving hernia societies and national surgical colleges based in low–middle-income countries could expand the training programme, while monitoring its quality and safety. Expanding mesh use could be a first step, before expanding the use of more advanced technologies, for which training is more demanding, supply chains are more complex, and the costs are higher.

Collaborators

National Institute for Health and Care Research (NIHR) Global Health Research Unit on Global Surgery

Full authorship listed in the supplementary material. Writing group (study management group): M Picciochi (University of Birmingham, Birmingham, UK), AO Ademuyiwa (University of Lagos, Lagos, Nigeria), A Adisa (Obafemi Awolowo University, Ile-Ife, Nigeria), AE Agbeko (Komfo Anokye Teaching Hospital, Kumasi, Ghana), JA Calvache (Universidad del Cauca, Popayan, Colombia and Erasmus MC Rotterdam, The Netherlands), D Chaudhry (University Hospitals of North Midlands NHS Trust, Stoke-on-Trent, UK), R Crawford (Chris Hani Baragwanath Hospital, Soweto, South Africa), AC Dawson (The University of Newcastle, Central Coast Clinical School, Newcastle, Australia), M Elhadi (University of Tripoli, Tripoli, Libya), A Ghaffar (Aga Khan University, Karachi, Pakistan), D Ghosh (Christian Medical College, Ludhiana, India), J Glasbey (University of Birmingham, Birmingham, UK), P Haque (Christian Medical College, Ludhiana, India), E Harrison (University of Edinburgh, Edinburgh, UK), A Isik (Istanbul Medeniyet University, Istanbul, Turkey), I Jakaityte (University of Exeter Medical School, Exeter, UK), SK Kamarajah (University of Birmingham, Birmingham, UK), O Kouli (The Walton Centre, Liverpool, UK), I Lawani (University of Abomey Calavi, Cotonou, Benin), S Lawani (University of Abomey Calavi, Cotonou, Benin), V Ledda (University of Birmingham, Birmingham, UK), E Li (University of Birmingham, Birmingham, UK), J Martin (University of Western Ontario, Ontario, Canada), A Minaya Bravo (Hospital Universitario HM Rivas, Madrid, Spain), D Morton (University of Birmingham, Birmingham, UK), D Nepogodiev (University of Birmingham, Birmingham, UK), F Ntirenganya (University of Rwanda, Kigali, Rwanda), O Omar (University of Birmingham, Birmingham, UK), SZY Ooi (Morrison Hospital, Swansea, UK), R Oppong (University of Birmingham, Birmingham, UK), F Pata (University of Calabria, Rende, Italy), A Ramos-De la Medina (Hospital Español de Veracruz, Veracruz, Mexico), M Sampaio-Alves (Faculdade de Medicina da Universidade do Porto, Porto, Portugal), JFF Simoes (Hospital Garcia de Orta, Almada, Portugal), M Steinruecke (University of Cambridge School of Clinical Medicine, Cambridge,

UK), S Tabiri (Tamale Teaching Hospital, Tamale, Ghana), A Bhangu (University of Birmingham, Birmingham, UK).

Funding

This study was supported by a National Institute for Health and Care Research (NIHR) Global Health Research Unit on Global Surgery grant (NIHR133364) and a project research grant from the Portuguese Hernia and Abdominal Wall Society (Sociedade Portuguesa de Hernia e Parede Abdominal). The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author and analysis group had full access to all the data in the study and the corresponding author and the writing committee had final responsibility for the decision to submit for publication. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, or the UK Department of Health and Social Care.

Author contributions

Writing group (study management group) co-authors were involved in conceptualisation, methodology, investigation, resources, data curation, writing-original draft, review and editing, visualisation, supervision, project administration and funding acquisition. Data handling and management co-authors were involved in validation, investigation, resources and project administration. Dissemination Committee co-authors were involved in conceptualisation, investigation, resources, data curation, and project administration. Hospital-lead co-authors were involved in validation, investigation, resources, data curation and project administration. Collaborator co-authors were involved in investigation, resources, data curation and project administration.

Disclosure

M. Picciochi reports a project research grant from the Portuguese Hernia and Abdominal Wall Society (Sociedade Portuguesa de Hernia e Parede Abdominal). A. Bhangu reports a National Institute for Health Research (NIHR) Global Health Research Unit grant. The authors declare no other conflict of interest.

Supplementary material

Supplementary material is available at BJS online.

Data availability

Anonymized data are available upon request from the writing group and successful completion of a data sharing agreement through an Application Programming Interface linked to the REDCap data server hosted at the University of Birmingham, Birmingham, UK.

References

- Howitt P, Darzi A, Yang GZ, Ashrafian H, Atun R, Barlow J et al. Technologies for global health. *Lancet* 2012;**380**:507–535
- HerniaSurge Group. International guidelines for groin hernia management. *Hernia* 2018;**22**:1–165
- Stabilini C, van Veenendaal N, Aasvang E, Agresta F, Aufenacker T, Berrevoet F et al. Update of the international HerniaSurge guidelines for groin hernia management. *BJS Open* 2023;**7**:zrad080
- Lockhart K, Dunn D, Teo S, Ng JY, Dhillon M, Teo E et al. Mesh versus non-mesh for inguinal and femoral hernia repair. *Cochrane Database Syst Rev* 2018; **(9)**CD011517
- Tabiri S, Owusu F, Atindaana Abantanga F, Moten A, Nepogodiev D, Omar O et al. Mesh versus suture repair of primary inguinal hernia in Ghana. *BJS Open* 2019;**3**:629–633
- Guillaumes S, Juvany M. Inguinal hernia repairs performed for recurrence in Spain: population-based study of 16 years and 1,302,788 patients. *Hernia* 2022;**26**:1023–1032
- Ma Qiuyue, Jing Wenzhan, Liu Xiaoli, Liu Jue, Liu Min, Chen Jie. The global, regional, and national burden and its trends of inguinal, femoral, and abdominal hernia from 1990 to 2019: findings from the 2019 Global Burden of Disease Study - a cross-sectional study. *Int J Surg* 2023. DOI:10.1097/JS9.0000000000000217
- Chao TE, Mandigo M, Opoku-Anane J, Maine R. Systematic review of laparoscopic surgery in low- and middle-income countries: benefits, challenges, and strategies. *Surg Endosc* 2016;**30**:1–10
- WHO. *Medical Devices: Managing the Mismatch: An Outcome of the Priority Medical Devices Project*. 2010. https://iris.who.int/bitstream/handle/10665/44407/9789241564045_eng.pdf?sequence=1 (accessed 29 January 2024)
- Ashley T, Ashley HF, Wladis A, Nordin P, Ohene-Yeboah M, Rukas R et al. Standardised competency-based training of medical doctors and associate clinicians in inguinal repair with mesh in Sierra Leone. *World J Surg* 2023;**47**:2330–2337
- Patterson T, Currie P, Patterson S, Patterson P, Meek C, McMaster R. A systematic review and meta-analysis of the post-operative adverse effects associated with mosquito net mesh in comparison to commercial hernia mesh for inguinal hernia repair in low income countries. *Hernia* 2017;**21**:397–405
- Wang YT, Meheš MM, Naseem HR, Ibrahim M, Butt MA, Ahmed N et al. Assessing the impact of short-term surgical education on practice: a retrospective study of the introduction of mesh for inguinal hernia repair in sub-Saharan Africa. *Hernia* 2014;**18**:549–556
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Epidemiology* 2007;**18**:800–804
- Clavien PA, Strasberg SM. Severity grading of surgical complications. *Ann Surg* 2009;**250**:197–198
- Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform* 2019;**95**:103208
- GlobalSurg Collaborative, National Institute for Health Research Global Health Research Unit on Global Surgery. Global variation in postoperative mortality and complications after cancer surgery: a multicentre, prospective cohort study in 82 countries. *Lancet* 2021;**397**:387–397
- COVIDSurg Collaborative, GlobalSurg Collaborative. Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. *Anaesthesia* 2021;**76**:748–758
- Riley RD, Ensor J, Snell KI, Harrell FE, Martin GP, Reitsma JB et al. Calculating the sample size required for developing a clinical prediction model. *BMJ* 2020;**368**:m441

19. Ryan I, Shah KV, Barrero CE, Uamunovandu T, Ilbawi A, Swanson J. Task shifting and task sharing to strengthen the surgical workforce in Sub-Saharan Africa: a systematic review of the existing literature. *World J Surg* 2023;**47**:3070–3080
20. Sani R. Endorsement of the HerniaSurge guidelines by the Afro Middle East Hernia Society. *Hernia* 2018;**22**:175
21. Sharma A. Endorsement of the HerniaSurge guidelines by the Asia Pacific Hernia Society. *Hernia* 2018;**22**:173
22. Campanelli G. Endorsement of the herniasurge guidelines by the European Hernia Society. *Hernia* 2018;**22**:169
23. Techapongsatom S, Tansawet A, Pattanaprateep O, Attia J, McKay GJ, Thakkestian A. Cost-effectiveness analysis of mesh fixation techniques for laparoscopic and open inguinal hernia surgeries. *BMC Health Serv Res* 2022;**22**:1125
24. Sivakumar J, Chen Q, Hii MW, Cullinan M, Choi J, Steven M et al. Learning curve of laparoscopic inguinal hernia repair: systematic review, meta-analysis, and meta-regression. *Surg Endosc* 2023;**37**:2453–2475
25. Merola G, Cavallaro G, Iorio O, Frascio M, Pontecorvi E, Corcione F et al. Learning curve in open inguinal hernia repair: a quality improvement multicentre study about Lichtenstein technique. *Hernia* 2020;**24**:651–659
26. McCormack K, Scott NW, Go PM, Ross S, Grant AM; EU Hernia Trialists Collaboration. Laparoscopic techniques versus open techniques for inguinal hernia repair. *Cochrane Database Syst Rev* 2003; **(1)**CD001785
27. Organization for Economic Cooperation and Development (OECD). *Surgical operations and procedures statistics*. 2023. https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Surgical_operations_and_procedures_statistics#Number_of_surgical_operations_and_procedures (accessed 12 December 2023).