

# EUR Research Information Portal

## Family perspectives and experiences on implementing telehealth in pediatric palliative care

**Published in:**

European Journal of Pediatrics

**Publication status and date:**

Published: 08/04/2025

**DOI (link to publisher):**

[10.1007/s00431-025-06124-6](https://doi.org/10.1007/s00431-025-06124-6)

**Document Version**

Publisher's PDF, also known as Version of record

**Document License/Available under:**

CC BY

**Citation for the published version (APA):**

Peñarrubia-San-Florencio, L., Godoy Junior, C., Ricart, S., Vilarrubí, S. N., Ruiz-Herguido, C., Esteve, A. V., Meca-Santamaria, J., Blanco-Blanco, J., & Valls, L. L. (2025). Family perspectives and experiences on implementing telehealth in pediatric palliative care: a qualitative approach. *European Journal of Pediatrics*, 184(5), Article 287. <https://doi.org/10.1007/s00431-025-06124-6>

[Link to publication on the EUR Research Information Portal](#)

**Terms and Conditions of Use**

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.

In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

**Take-down policy**

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: [openaccess.library@eur.nl](mailto:openaccess.library@eur.nl). Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.



# Family perspectives and experiences on implementing telehealth in pediatric palliative care: a qualitative approach

Lucia Peñarrubia-San-Florencio<sup>1,2,3</sup> · Carlos Godoy Junior<sup>4</sup> · Silvia Ricart<sup>1,2</sup> · Sergi Navarro Vilarrubí<sup>1,2</sup> · Cristina Ruiz-Herguido<sup>2,5</sup> · Arnau Valls Esteve<sup>2,6</sup> · Julia Meca-Santamaria<sup>5,6</sup> · Joan Blanco-Blanco<sup>3,7,8</sup> · Laura Lahuerta Valls<sup>3,9</sup>

Received: 9 August 2024 / Revised: 24 March 2025 / Accepted: 2 April 2025  
© The Author(s) 2025

## Abstract

Telehealth is seen as a promising avenue to improve accessibility and quality of pediatric palliative care (PPC) yet engaging users with digital tools (DT) over the long term poses a challenge. Aligning telehealth initiatives with the actual needs and expectations of families is crucial for sustainable care models. To explore family perspectives and experiences with telehealth in PPC, focusing on their needs, perspectives, concerns, and hopes to refine digital care models. This research employed a qualitative methodology, gathering data through a combination of semi-structured interviews and focus group discussions with primary caregivers (PCGs) of PPC patients. PCGs of children utilizing telehealth services in PPC contexts. Interviews with seven individuals and a focus group of eight revealed three main themes: the experience of engaging with a PPC service that utilizes telehealth, the transformative potential of telehealth within PPC, and the barriers and facilitators influencing its adoption. A key finding was the collective desire among PCGs for telehealth solutions that support a patient and family-centered, holistic care model that integrates various health services without compromising the humanized essence of care. **Conclusions:** Families are generally positive about integration of telehealth into PPC, emphasizing the need for telehealth models that maintain the core values of humanized care. Success depends on engaging end-users throughout development, implementing user-friendly technology that fits into family life, and focusing on customization for patients.

## What is known:

- Telehealth has emerged as an innovative response to meet the increasing demand for highly specialized care in PPC.
- Digital health solutions often face increasing abandonment rates over time.

## What is new:

- Families caring for children with palliative needs are receptive to telehealth, identifying essential features for its long-term success.
- They value the preservation of personal attention in care, emphasizing the importance of maintaining human connection alongside digital innovations.

**Keywords** Pediatrics · Palliative medicine · Patient-centered care · Telemedicine · Caregivers

Communicated by Piet Leroy

✉ Joan Blanco-Blanco  
joan.blanco@udl.cat

<sup>1</sup> Pediatric Palliative Care Service, Sant Joan de Déu Hospital, Esplugues de Llobregat, Barcelona, Spain

<sup>2</sup> Institut de Recerca Sant Joan de Déu (IRSJD), Esplugues de Llobregat, Barcelona, Spain

<sup>3</sup> Faculty of Nursing and Physiotherapy and GESEC, Universitat de Lleida, Lleida, Spain

<sup>4</sup> Erasmus School of Health Policy and Management, Erasmus University Rotterdam, Rotterdam, Netherlands

<sup>5</sup> Sant Joan de Déu Foundation, Esplugues de Llobregat, Barcelona, Spain

<sup>6</sup> Innovation Department and Innovation, Health Technologies Group Sant Joan de Déu Hospital, Esplugues de Llobregat, Barcelona, Spain

<sup>7</sup> CIBER, Frailty and Healthy Aging (CIBERFES), Madrid, Spain

<sup>8</sup> Institute of Biomedical Research, GRECS, Lleida, Spain

<sup>9</sup> Granollers General Hospital, Granollers, Barcelona, Spain

## Abbreviations

COREQ	Consolidated criteria for reporting qualitative research
DT	Digital tools
PC	Palliative care
PCGs	Primary caregivers
PPC	Pediatric palliative care
WHO	World Health Organization

## Introduction

Pediatric palliative care (PPC), as defined by the World Health Organization (WHO), is a medical specialty that delivers comprehensive care encompassing physical, psychological, social, and spiritual aspects to pediatric patients affected by life-threatening and life-limiting conditions and their families, with the goal of enhancing their quality of life [1]. Initiated at diagnosis, this continuous care transcends the patient's therapeutic journey and can be delivered across settings, from hospitals to homes, ensuring personalized care that prioritizes the child's comfort in their most familiar surroundings [2].

As the prevalence of chronic and complex conditions increases, the demand for PPC intensifies, presenting a considerable challenge for healthcare providers to offer high-quality, tailored care to patients and their families [3, 4]. It is estimated that each year, over 21 million children worldwide have special needs that require pediatric palliative care PPC [5]. The scarcity of specialized services equipped to deliver all-encompassing care to children with life-threatening and life-limiting conditions, especially those in home settings who depend on medical technologies, adds layers of complexity to care provision [6]. Research underscores the necessity of a holistic and multifaceted approach to home care for advanced-stage diseases, one that upholds rigorous safety standards irrespective of the setting chosen by the patient's family [7, 8].

Information and communication technologies are now key in advancing healthcare quality, modernization, and sustainability [9]. Recognizing their value, WHO supports the incorporation of these technologies into homecare systems, citing the efficacy of hybrid models that integrate traditional and remote care methods. These models have not only expanded healthcare capabilities but have also yielded remarkable efficiencies [10, 11]. Although telehealth initiatives have predominantly catered to adults with chronic conditions that require consistent monitoring (chronic obstructive pulmonary disease, diabetes, heart diseases, or obesity) [12], they are now increasingly being adopted in palliative care (PC) settings. This shift towards including video consultations has expanded PC's capacity, a development

significantly hastened by the exigencies of the COVID-19 pandemic [13].

Despite showing promising outcomes in various medical domains, the widespread adoption and effectiveness of telehealth programs within PPC still present uncertainties [14]. Moreover, trends indicate a potential decline in user engagement and motivation over time concerning the continuous use of digital health services [15]. In response to these challenges, there is a growing emphasis on designing digital health tools that are centered around the user's needs and preferences. This study, conducted within the framework of the European AICCELERATE project [16], an initiative aimed at promoting the development of artificial intelligence and digital tools (DT) in healthcare, employs a qualitative methodology to delve into the perspectives, needs, and experiences of patients and primary caregivers (PCGs) engaged with a hybrid telehealth care model. The goal is to circumvent the common pitfall of creating sophisticated technologies that fail to address the real-world needs of their target demographic, thereby enhancing the success rate of technology adoption post-implementation.

## Study objectives and purpose

### Primary objective

Explore families' perceptions of using telehealth for PPC, aiming to understand their needs, perceptions, concerns, and expectations. This contributes to enhancing telehealth care models in PPC settings.

### Secondary objectives

To investigate families' views on integrating telehealth solutions into PPC, identifying perceived benefits and challenges.

To analyze key features families deem essential for the long-term viability of tele-assistance in home settings, thereby promoting the effective development and implementation of user-centered telehealth interventions.

## Methodology

### Study design

This study employed a qualitative approach, incorporating semi-structured interviews, focus groups, and field notes from the researcher to address the research objectives.

## Data quality criteria

The methods and reports adhered to the Consolidated Criteria for Reporting Qualitative Research (COREQ) checklist [17].

## Ethical considerations

Adhering to the Helsinki Declaration and Human Medical Research Law, the study received approval from the local ethics committee of Hospital Sant Joan de Déu (PIC-123–19). Informed consent was obtained from all participants prior to their inclusion, providing them detailed information about the study's objectives and procedures. Participation was entirely voluntary, with no material incentives offered.

## Population and sample

Participants were primary PCGs of children with life-threatening and life-limiting conditions, receiving care under the telehealth hybrid model of the PPC unit at Hospital Sant Joan de Déu, Barcelona. Since 2021, this PPC unit has integrated telehealth into its services, including telemedicine and digital platforms for information and communication, such as a patient portal [18]. Eligibility for inclusion required individuals to have been PCGs of a child under the PPC unit's supervision for at least 6 months prior to joining the study. Exclusion criteria included PCGs of children in their final days of life and language barriers (limited proficiency in Spanish or Catalan).

We employed purposive sampling to ensure a diverse and heterogeneous participant base. Recruitment was led by the principal investigator (LPS, PhD candidate). Recruitment involved detailed study explanations over the phone, sending informed consent documents through the institutional portal, and collecting signed consent during a physical visit with the research team.

## Data collection

Data collection utilized a blend of in-depth individual interviews, focus groups, and entries from a researcher's field journal capturing observations and reflective insights [19]. Both the semi-structured interviews and the focus groups were conducted remotely, with participants joining from their homes via smartphones, tablets, or personal computers. The data collection period spanned from January to April 2022. The semi-structured interviews and focus groups were led by the principal investigator (LPS) and a member of the associated research team (LLV, PhD). Both women researchers bring together over two decades of experience in health services research. The principal investigator, an expert PC

nurse, was familiar to the participants, although not serving as their direct care nurse.

An interview guide based on literature and the research group's experience was developed (Supplementary Material 1), covering topics on the conception and use of new technologies in PPC. Additionally, open-ended questions were included to explore how new information and communication technologies could enhance the care and well-being of the child and family (Supplementary Material 2). Concurrently, a member of the associated research team kept a field journal to record observations during the interviews and focus group (SRV, PhD).

Interviews were recorded in audio and video, then transcribed, and participants' names were anonymized by the principal investigator. After transcription, the audio and video recordings were deleted.

## Data analysis

The qualitative analysis adopted an inductive approach, enabling the identification of emerging patterns and the discovery of insightful concepts within the complexity of pediatric palliative experiences [20]. Coding and analysis were performed in pairs, parallel to data collection, including a team member (CGJ, PhD candidate) not directly engaged in conducting interviews or focus groups [21]. A coding framework was developed, modified, and refined using ATLAS.ti 22 software as new themes and subthemes emerged throughout the analysis (Supplementary Material 3). Two discussion meetings were held among the reviewers to aid in the interpretation of data and to validate the emerging themes identified.

The total number of interviews was determined by reaching data saturation. Following this, a focus group was conducted to specifically delve into the emerging categories from the semi-structured interviews and triangulate the collected information.

## Results

We conducted seven semi-structured individual interviews. To further validate and refine our findings, a focus group session was held with eight new participants, distinct from those in the initial interviews. This step facilitated the triangulation of data and the fine-tuning of previously identified categories. There were no declines to participate among those contacted. The average duration of the interviews was 31 min, with a range of 24 to 43 min, while the focus group session extended to 75 min.

All participants were in a private setting during their interview, except for one mother, who was accompanied by her breastfeeding infant, and a pair of caregiver

participants who joined the interview together. Table 1 outlines the sociodemographic details of both the semi-structured interview and focus group participants, including information about the patients under their care.

After the iterative analysis of the data, 12 codes were consolidated from the original 76 codes, which were grouped into 3 main categories after axial coding. Table 2 shows the main themes and subthemes derived from the analysis.

**Table 1** Demographic and clinical characteristics of caregivers and patients included in the study

Characteristics	Number (N)	Percentages (%)
<b>Caregivers' gender</b>		
Male	7	47
Female	8	53
<b>Caregivers' age</b>		
< 30	3	20
30–40	7	47
> 40	5	33
<b>Marital status</b>		
Married or cohabiting	14	93
Divorced or not cohabiting	1	7
<b>Patients' underlying illnesses</b>		
Oncological	3	20
Neurological	8	53
Other	4	27
<b>Patients' age</b>		
< 1	2	13
1–5	5	33
6–12	3	20
13–16	4	27
> 16	1	7
<b>Patients' gender</b>		
Male	9	60
Female	6	40
<b>Siblings</b>		
0	6	40
1	6	40
2	3	20
3 or +	0	0
<b>Medical devices at home</b>		
0	3	20
1	3	20
2	4	27
3 or +	5	33

## Insights into hybrid care: families' experiences with the support received from the current hybrid PPC

This category encapsulates the experiences, needs, and evaluations concerning the support patients and their families receive from the hybrid service provided by the PPC team. Central to the service provided by the hybrid telehealth PPC is the provision of humanized care, a fundamental need highlighted by all participants as a defining characteristic of the current care approach. Key elements of this humanized care include direct contact, personalized support tailored to the unique circumstances, and progression of the patient's condition (Q#1, Q#2); empowerment of PCGs through ongoing education offered by healthcare professionals; and clear, fluid communication with the clinical team (Q#3).

*Q#1*—“...hearing your voices is what relaxes us...”

*Q#2*—“...the peace of mind that you [PPC providers] can see it, if something [wrong] is happening...”

*Q#3*—“Seeing you [PPC providers] and speaking with a professional, I think, gives you [the PCGs] more confidence, more capability.”

A significant concern among participants was the restricted availability of the specialized PPC team, noting that access to these expert services is not always possible around the clock. This limitation poses considerable challenges, especially when PCGs encounter new, acute, or unexpected health crises. In such situations, the most commonly used alternative is to seek help from local emergency services, which may not be adequately equipped to address the specialized needs of PPC patients effectively (Q#4). This discontinuity in care exacerbates PCGs' uncertainty and distress. In the absence of immediate access to PPC specialists, some PCGs opt to travel to the hospital where the PPC team is based. However, this decision presents its own set of logistical challenges, including travel distances and the added strain of managing transportation (Q#5).

*Q#4*—“When you visit [the emergency service] for a respiratory issue, the immediate response in primary care emergencies is to administer salbutamol. However, in cases like my child's, salbutamol is inappropriate because the underlying condition differs. This reflects a systemic issue where individual patient needs are overlooked in favor of a one-size-fits-all approach.”

*Q#5*—“The hospital is always crowded, and for me, it's a hassle to get the child into the car, drive there, find parking, and it takes a long time to get to Barcelona depending on the time of day. If it's something urgent, then I have no choice but to go...”

**Table 2** List of main themes and sub-themes from the analysis

Themes	Sub-themes
C.1. Insights into hybrid care: families' experiences with the support received from the current hybrid PPC	C.1.1. Direct interaction with individuals and personalized attention C.1.2. Fluid communication and empowerment C.1.3. Restricted availability and discontinuity in care
C.2. Shaping the future: leveraging information and communication technologies to transform PPC	C.2.1. Enhancement of communication and information C.2.2. Coordination and integration among community health services (care network) C.2.3. Modernizing care approaches and enhancing family empowerment C.2.4. Comprehensive and holistic support
C.3. Advancing PPC through digital transformation: barriers and facilitators	C.3.1. Maintenance of human connections C.3.2. Security and privacy C.3.3 Initial instruction and ease of use C.3.4 Diligent use C.3.5 Preference for mobile devices

### Shaping the future: leveraging information and communication technologies to transform PPC

Participants engaged in envisioning scenarios and solutions for integrating digital technologies into the PPC service model. A significant number of participants showed enthusiasm for trying out new digital solutions, demonstrating an openness to adopting technological advances in PPC. They especially appreciated how these technologies could offer continuous insights into their children's health status, thereby minimizing the need for hospital visits and enhancing overall communication (Q#6, Q#7).

*Q#6*—"Having the ability to receive updates on my child's condition through virtual means or a simple phone call is incredibly helpful. It eliminates the need for in-person visits, which can be cumbersome.... for regular updates, this virtual approach is ideal and much appreciated."

*Q#7*—"Sometimes we don't really know how to interpret what the problem is. We can see trends, we can see things, but it's a bit hard [identifying and understanding the specific health issues or needs of patients]. I think it would be very good if there was a direct transfer of information [real-time remote monitoring]."

Moreover, participants viewed DT beneficial for better care coordination among all parties involved in a child's treatment (Q#8). Additionally, they highlighted the importance of providing parents with the necessary tools and information to manage the complexities of their child's care more effectively, thereby fostering greater empowerment and ensuring that care practices remain up to date with the most current advancements and resources (Q#9). Reflecting the holistic approach of PPC, they also recognized the value

of these technologies in delivering comprehensive support that spans beyond medical care, including emotional support (Q#10).

*Q#8*—"There needs to be very good communication between the home environment, the hospital environment, and the school environment, as children spend many hours at school. And the information, the ability to respond, all of this also needs to be present there."

*Q#9*—"Parent training should focus on helping us become more independent and informed about new developments of any new or existing support available. We're currently discovering resources we were previously unaware of, and this affects us all in various ways: logistics, family finances, and our child's well-being."

*Q#10*—"Emotional support is also crucial for parents."

### Advancing PPC through digital transformation: barriers and facilitators

This category identified central elements for integrating digital solutions into PPC, aiming for a balanced approach between technology and traditional care. At the core of this integration lies the principle of the maintenance of the humanized care, highlighted by participants' preference to keep personal interactions alive within digital care settings (Q#11). While some degree of automation is acceptable, participants emphasized that it should enhance rather than replace the human element and direct connection with the care team (Q#12). Additionally, PCGs expressed confidence in the care team's ability to handle digital information securely, placing responsibility for data protection on healthcare providers (Q#13).



*Q#11*—“Sometimes virtual assistance might not be sufficient. Sometimes the help of a professional at home is inevitable.”

*Q#12*—“I feel more secure talking to a person.....”

*Q#13*—“We know where the data goes; it will be sent directly to you. I don’t think it can get lost in the network; it should be secure enough, I believe....”

The need for simplicity and smooth integration of digital technologies into daily routines was another key theme (*Q#14*). Participants expressed feeling overwhelmed by the demands of caregiving at home, highlighting the necessity for DT to alleviate rather than exacerbate these challenges (*Q#15*).

*Q#14*—“Provided I receive proper training, and the devices aren’t overly complicated to use, I’m completely comfortable with it.”

*Q#15*—“We have so much work every day that if it’s going to involve more work on our part [the use of DT], then no.”

When it comes to digital instruments for collecting patient data that might require caregiver involvement, opinions varied: some preferred tools that demanded minimal, sporadic engagement (*Q#16*), whereas others were open to more frequent interaction if it tangibly aided their children (*Q#17*). Additionally, a preference emerged for using mobile devices like smartphones or tablets for these digital interactions (*Q#18*).

*Q#16*—“Sending the data whenever we notice something that warrants attention is crucial.”

*Q#17*—“It wouldn’t feel like a burden at all [completing online questionnaires], primarily because it concerns my son. I’m confident I can dedicate at least 10 min daily to provide all the necessary data about him.”

*Q#18*—“I have a computer at home that’s mostly collecting dust; my phone is what I primarily use now, given the computer’s obsolescence. Essentially, the smartphone is my go-to device.”

## Discussion

The main objective of this study was to delve into families’ perspectives on the use of telehealth and digital solutions in the field of PPC, to understand their needs, perceptions, concerns, and expectations. Our findings reveal an interest among families of children with complex illnesses and PPC needs towards the integration of these technologies into daily care. They foresee a future where technology is integrated with traditional care, forming a harmonious synergy. This integration is seen as a means to improve parental participation in home care settings, thereby strengthening home

care capabilities, increasing family autonomy, and streamlining the care process. Such a vision aligns with existing research highlighting the empowerment of families through the implementation of DT in PPC contexts [22].

Interestingly, our study participants displayed openness to adopting new digital innovations, without any discernible apprehension. This receptiveness to digital health solutions may be influenced by our sample’s demographic profile, mainly comprising individuals younger than 40. Previous studies have shown that this demographic, being more versed in digital technology, often possesses a high degree of digital fluency, essential for adapting to technological advancements [23]. Additionally, the experience gained from integrating various digital health solutions during the COVID-19 pandemic has likely fostered their adaptability and comfort with these technologies [24]. Notably, our findings also reveal a lack of significant concerns over privacy among participants. They seemed at ease with the idea of sharing sensitive data digitally. This contrasts with the prevalent privacy concerns observed in adult PC settings, where the introduction of telehealth tools often leads to anxieties over data privacy and management [25].

A paradox emerged in our study. While DT are valued for enhancing communication with PPC teams, there is a simultaneous worry that they may dilute the crucial human connections between healthcare professionals and families. PCGs recognize that, although digital solutions streamline care processes, they cannot substitute the essential elements of trust, empathy, and personal connection that are particularly crucial during the difficult times of illness. This concern aligns with similar research findings [26–28], suggesting that telehealth, despite its benefits, cannot replicate the comprehensive depth and quality of face-to-face care. Echoing the sentiments of prior studies, the need for DT to blend effortlessly into family life was emphasized [25]. Participants advocated for user-friendly technologies that are straightforward to use, set up, and manage, considering the significant emotional and physical stresses families face. The challenge of configuring and handling these devices adds an extra layer of burden.

The study has some limitations that should be taken into account for the correct interpretation of the results. The participant sample of the study was limited to a single PPC service in Barcelona, Spain. This team integrates an asynchronous communication pathway with patients and families through the patient portal, meaning that participants were not entirely unfamiliar with adopting or assessing new technologies. Given that healthcare practices and levels of digital literacy can vary significantly between regions, the applicability of our findings might be limited beyond this specific context [29]. Expanding the study to include caregivers from other centers, particularly those without prior exposure to digital tools, could provide valuable insights and

enhance the generalizability of the results. Future research could explore this broader inclusion to better assess digital adoption in more diverse care settings.

Furthermore, all interviews in our study were conducted online, either due to participants' preference or logistical reasons. Although existing research suggests that both face-to-face and virtual interviews generate similar amounts of data and thematic richness, face-to-face interactions tend to be more profound [30]. Despite this, remote interviews via video, phone, or online platforms remain reliable and credible methods [31]. Finally, the interviews and focus groups were carried out without the use of a physical prototype or actual DT that might be implemented in PPC. This lack of tangible examples may have restricted participants' grasp of certain practical applications, while encouraging speculation about various features and potentialities. Direct interaction with these tools could uncover unexpected benefits or obstacles, thereby altering users' viewpoints and expectations [32].

## Conclusion

This study highlights the critical importance of addressing the needs of PCGs and children in the realm of hybrid PPC. It brings attention to family concerns that technological advancements could potentially detract from the quality of care by replacing vital human interactions or by adding to their existing burdens. Such insights underscore the critical need for a thoughtful approach in the ideation, design, and development of future digital solutions, with an aim to augment rather than replace face-to-face care, prioritizing personal connections, customized care, and user-friendly technology.

Moving forward, it becomes imperative to also incorporate the viewpoints of clinical staff, another key user group, to ensure that the development of any new tool effectively responds to the comprehensive needs of the entire end-user community before its design is finalized.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00431-025-06124-6>.

**Acknowledgements** We thank the caregivers who have participated for their trust and knowledge contributed to this study.

**Authors' contributions** LPSF played a lead role in data curation, formal analysis and investigation. She was the main contributor to the original draft and led the review and editing of the writing. CGJ was supportive in data curation. He equally contributed to formal analysis, and methodology and was supportive in both the original draft and review and editing phases of writing. SR was supportive in conceptualization, project administration, methodology and supervision. She equally contributed to resource management, supported the review and editing of the written content. CRH; SNV; AVE i JMC offered equal contributions to project administration, investigation, and provided support in

writing review and editing. LLV shared equal responsibilities in project administration and resource management and was supportive in the writing review and editing process. Lastly, JBB equally contributed to conceptualization, project administration, methodology and supervision, and provided support in the original draft writing, formal analysis, and review and editing stages.

**Funding** Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature. This research is part of the AICCEL-ERATE project. This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement n° 101016902.

**Data availability** No datasets were generated or analysed during the current study.

## Declarations

**Ethics approval** The study was conducted according to the principles of the Declaration of Helsinki and in accordance with the Medical Research Involving Human Subjects Act (WMO) and was approved by the local ethics committee of Hospital Sant Joan de Déu (PIC- 123–19).

**Consent to participate** An informed consent was obtained from all interviewees before participation.

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Benini F, Spizzichino M, Trapanotto M, Ferrante A (2008) Pediatric palliative care. *Ital J Pediatr* 34(1):4. <https://doi.org/10.1186/1824-7288-34-4>
2. World Health Organization. (n.d.). Palliative care for children [Internet]. <https://www.who.int/europe/news-room/fact-sheets/item/palliative-care-for-children>. Accessed April 2, 2024.
3. Bjur KA, Wi C, Ryu E, Crow SS, King KS, Juhn YJ (2019) Epidemiology of children with multiple complex chronic conditions in a mixed urban-rural US community. *Hosp Pediatr* 9(4):281–290. <https://doi.org/10.1542/hpeds.2018-0091>
4. Fraser LK, Gibson-Smith D, Jarvis S, Norman P, Parslow RC (2021) Estimating the current and future prevalence of life-limiting conditions in children in England. *Palliat Med* 35(9):1641–1651. <https://doi.org/10.1177/0269216320975308>
5. Connor SR, Downing J, Marston J (2017) Estimating the global need for palliative care for children: a cross-sectional analysis. *J Pain Symptom Manage* 53(2):171–177. <https://doi.org/10.1016/j.jpainsymman.2016.08.020>
6. Together for Short Lives. (2018). The state of the UK children's hospice nursing workforce: a report on the demand and supply of



- nurses to children's hospices [Internet]. [https://www.togetherforshortlives.org.uk/app/uploads/2018/03/PolRes\\_Together\\_for\\_Short\\_Lives\\_-\\_The\\_state\\_of\\_children\\_s\\_palliative\\_care\\_nursing\\_in\\_2017.pdf](https://www.togetherforshortlives.org.uk/app/uploads/2018/03/PolRes_Together_for_Short_Lives_-_The_state_of_children_s_palliative_care_nursing_in_2017.pdf). Accessed February 1, 2024.
7. Winegard B, Miller EG, Slamon NB (2017) Use of telehealth in pediatric palliative care. *Telemedicine and e-Health* 23(11):938–940. <https://doi.org/10.1089/tmj.2016.0251>
  8. Normand, C., Martin-Moreno, J. M., Harris, M., Gorgojo, L., Clark, D., & Centeno, C. (2008). Palliative care in the European Union 2008 [Internet]. <https://policycommons.net/artifacts/1341562/palliative-care-in-the-european-union/>. Accessed June 2, 2024.
  9. World Health Organization. (2019, April 17). WHO releases first guideline on digital health interventions [Internet]. <https://www.who.int/news/item/17-04-2019-who-releases-first-guideline-on-digital-health-interventions>. Accessed February 10, 2024.
  10. World Health Organization. (2019). WHO guideline: recommendations on digital interventions for health system strengthening [Internet]. <https://www.who.int/publications/i/item/9789241550505>. Updated 2019. Accessed January 12, 2024.
  11. World Health Organization. (2016). Monitoring and evaluating digital health interventions: a practical guide to conducting research and assessment [Internet]. <https://iris.who.int/bitstream/handle/10665/252183/9789241511766-eng.pdf?sequence=1&isAllowed=y>. Accessed February 15, 2024.
  12. Paré G, Jaana M, Sicotte C (2007) Systematic review of home telemonitoring for chronic diseases: the evidence base. *J Am Med Inform Assoc* 14(3):269–277. <https://doi.org/10.1197/jamia.M2270>
  13. Vincent D, Peixoto C, Quinn KL et al (2022) Virtual home-based palliative care during COVID-19: a qualitative exploration of the patient, caregiver, and healthcare provider experience. *Palliat Med* 36(9):1374–1388. <https://doi.org/10.1177/02692163221116251>
  14. Holmen H, Riiser K, Winger A (2020) Home-based pediatric palliative care and electronic health: systematic mixed methods review. *J Med Internet Res* 22(2):e16248. <https://doi.org/10.2196/16248>
  15. Lie SS, Karlsen B, Oord ER, Graue M, Oftedal B (2017) Dropout from an eHealth intervention for adults with type 2 diabetes: a qualitative study. *J Med Internet Res* 19(5):e187. <https://doi.org/10.2196/jmir.7479>
  16. AICCELERATE. (2022). AICCELERATE smart hospital care pathway engine [Internet]. <https://aiccelerate.eu/>. Accessed July 25, 2024.
  17. Tong A, Sainsbury P, Craig J (2007) Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 19(6):349–357. <https://doi.org/10.1093/intqhc/mzm042>
  18. Hospital Sant Joan de Déu Barcelona. (n.d.). Atención paliativa y paciente crónico complejo [Internet]. <https://www.sjdhospitaibarcelona.org/es/servicios-asistenciales/atencion-paliativa-paciente-cronico-complejo>. Accessed January 12, 2024.
  19. Phillippi J, Lauderdale J (2018) A guide to field notes for qualitative research: context and conversation. *Qual Health Res* 28(3):381–388. <https://doi.org/10.1177/1049732317697102>
  20. Strauss, A., & Corbin, J. (1998). Basics of qualitative research: techniques and procedures for developing grounded theory (2nd ed.). Sage Publications, Inc.
  21. Palinkas LA (2014) Qualitative and mixed methods in mental health services and implementation research. *J Clin Child Adolesc Psychol* 43(6):851–861. <https://doi.org/10.1080/15374416.2014.910791>
  22. Harris N, Beringer A, Fletcher M (2016) Families' priorities in life-limiting illness: improving quality with online empowerment. *Arch Dis Child* 101(3):247–252. <https://doi.org/10.1136/archdischild-2015-308769>
  23. Le TV, Galperin H, Traube D (2023) The impact of digital competence on telehealth utilization. *Health Policy and Technology* 12(1):100724. <https://doi.org/10.1016/j.hlpt.2023.100724>
  24. Garfan S, Alamoodi AH, Zaidan BB et al (2021) Telehealth utilization during the Covid-19 pandemic: a systematic review. *Comput Biol Med* 138:104878. <https://doi.org/10.1016/j.combiomed.2021.104878>
  25. Steindal SA, Nes AAG, Godskesen TE et al (2023) Advantages and challenges of using telehealth for home-based palliative care: systematic mixed studies review. *J Med Internet Res* 25:e43684. <https://doi.org/10.2196/43684>
  26. Bradford N, Young J, Armfield NR et al (2012) A pilot study of the effectiveness of home teleconsultations in paediatric palliative care. *J Telemed Telecare* 18(8):438–442. <https://doi.org/10.1258/jtt.2012.gth103>
  27. Weaver MS, Robinson JE, Shostrom VK, Hinds PS (2020) Telehealth acceptability for children, family, and adult hospice nurses when integrating the pediatric palliative inpatient provider during sequential rural home hospice visits. *J Palliat Med* 23(5):641–649. <https://doi.org/10.1089/jpm.2019.0450>
  28. Hayes Bauer E, Ørsted Schultz AN, Brink A, Oechsle Jørgensen L, Bollig G (2021) Patients' and relatives' preferences for a palliative/oncology day ward and out-of-hours telemedicine-an interpretive description. *Healthcare (Basel, Switzerland)* 9(6):758. <https://doi.org/10.3390/healthcare9060758>
  29. Andreassen HK, Bujnowska-Fedak MM, Chronaki CE et al (2007) European citizens' use of e-health services: a study of seven countries. *BMC Public Health* 7:53. <https://doi.org/10.1186/1471-2458-7-53>
  30. Krouwel M, Jolly K, Greenfield S (2019) Comparing Skype (video calling) and in-person qualitative interview modes in a study of people with irritable bowel syndrome: an exploratory comparative analysis. *BMC Med Res Methodol* 19(1):219. <https://doi.org/10.1186/s12874-019-0867-9>
  31. Saarijärvi M, Bratt E (2021) When face-to-face interviews are not possible: tips and tricks for video, telephone, online chat, and email interviews in qualitative research. *Eur J Cardiovasc Nurs* 20(4):392–396. <https://doi.org/10.1093/eurjcn/zvab038>
  32. van Velsen L, Ludden G, Grünloh C (2022) The limitations of user- and human-centered design in an eHealth context and how to move beyond them. *J Med Internet Res* 24(10):e37341. <https://doi.org/10.2196/37341>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.