



ELSEVIER

Contents lists available at ScienceDirect

Geriatric Nursing

journal homepage: www.gnjournal.com

Involvement, worries and loneliness of family caregivers of people with dementia during the COVID-19 visitor ban in long-term care facilities

Marleen Prins, MSc^{a,b,*}, Bernadette Willemse, PhD^c, Claudia van der Velden, MSc^b,
Anne Margriet Pot, PhD^{a,d}, Henriëtte van der Roest, PhD^b

^a Erasmus School of Health Policy & Management, Health Care Governance (HCG), Burg. Oudlaan 50, 3000 DR Rotterdam, Zuid-Holland, The Netherlands

^b Netherlands Institute of Mental Health and Addiction, Department on Aging, Da Costakade 45, 3500 AS, Utrecht, Utrecht, The Netherlands

^c Care Farm Reigershoeve Foundation, Oosterweg 5B, 1968 KM, Heemskerk, Noord-Holland, The Netherlands

^d Optentia Research Focus Area, P O Box 1174, North-West University, Vanderbijlpark, 1900 South Africa



ARTICLE INFO

Article history:

Received 21 July 2021

Received in revised form 29 September 2021

Accepted 1 October 2021

Available online xxx

Keywords:

Dementia

Caregiving

Nursing homes

COVID-19

Loneliness

Worrying

Involvement

The Netherlands

ABSTRACT

To prevent COVID-19 from spreading in long-term care facilities (LTCFs), the Dutch government took restrictive measures, including a visitor-ban in LTCFs. This study examined the relationship between involvement of family caregivers (FCs) of people with dementia (PwD) living in LTCFs and FCs mental health during the visitor-ban, and whether this relationship was moderated by the frequency of alternative contact with PwD during the visitor-ban and FC resilience. This cross-sectional study collected data from 958 FCs. FCs who visited PwD more frequently before, were more worried during the visitor-ban than those with lower visiting frequency. FCs who visited the PwD daily before, but had minimal weekly contact during the visitor-ban, worried less. Resilient FCs who did social and task-related activities before, experienced less loneliness during the visitor-ban. It is advisable for healthcare professionals to reach out to these groups, to facilitate ongoing contact and help them overcome their loneliness.

© 2021 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Introduction

People with dementia living in long-term care facilities (LTCFs) are disproportionately at risk of becoming infected with COVID-19 because of their ageing immune system and comorbid diseases.^{1,2} Further, they have a decreased ability to comply with restrictive measures, e.g., social distancing and hand hygiene.³ The risk of COVID-19 outbreaks is especially high in LTCFs because residents often spend much of their time in communal spaces with other residents and have physical contact with care staff during the provision of care.^{4–6} A study on mortality associated with COVID-19 indicated that 46% of all people who died due to COVID-19 since the pandemic began until October 2020 lived in LTCFs.⁷

To protect people with dementia living in LTCFs from becoming infected with COVID-19, visitor bans have been implemented by governments worldwide.^{7,8} In the Netherlands, visitors were banned between mid-March and mid-May 2020, which meant that LTCFs closed their doors for visitors from outside as obligated by law.^{9,10} Similar measures taken within LTCFs prevented residents from going

outside and halted social and group activities.^{10–12} These restrictions, together with the lack of social interaction and physical closeness (e.g., holding hands, hugging) with family and friends are thought to have had an adverse impact on the well-being of both residents and their family and friends. Indeed, research has demonstrated the effects of the visitor ban on the well-being and autonomy of LTCF residents.^{10,11,13} Although other, “innovative” means of contact, like video calls, were used to facilitate interaction with family and friends, some residents nonetheless felt socially isolated,^{9,10,13} with potentially negative consequences for their health and well-being.^{14,15} On the other hand, there are indications that the visitor ban did not have a negative impact on the mood of a small proportion of residents because they experienced more peace and quietude in the LTCF.^{16,17}

Family caregivers, i.e., children, spouses, other family members, friends, neighbors, or legal guardians who have a personal relationship with the person with dementia and provide support, were presumably impacted by the visitor ban as well. This is because many family caregivers remain involved in the care and lives of people with dementia when they move to an LTCF.^{18,19} During the visitor ban, however, family caregivers were prohibited from such involvement. It remains unclear how this prohibition impacted family caregivers' mental health.

*Corresponding author.

E-mail address: prins@eshpm.eur.nl (M. Prins).

Before the pandemic, family caregivers contributed in different ways to the care of people with dementia. They provided task-related care (e.g., physical care tasks or helping with activities and meals), undertook pleasant personal or social activities, and gave psychosocial support.^{20–22} Also, family caregivers provided care staff with information about residents' life history, background, and needs. Further, they monitored the provision of care and acted as a link to the outside world for people with dementia.²³ Research has shown that the involvement of family caregivers in LTCFs contributes to improved quality of life of people with dementia as well as better quality of care.^{21,24}

For family caregivers, involvement in the care for people with dementia living in LTCFs is also important because it can have positive outcomes of benefit to their own mental health, including relationship continuity and emotional rewards like feeling appreciated and personal growth.^{25,26} Previous research on visitor bans as a consequence of the SARS outbreak revealed a range of negative effects on the physical and mental health of family caregivers.²⁷ In addition to missing the company of the person with dementia, family caregivers reported experiencing fear, worry, loss of control, frustration, and guilt.²⁷

Based on these previous research findings, we assumed that the visitor ban imposed during the COVID-19 pandemic had adverse effects on family caregivers' mental health.^{27,28} This is worrisome given the fact that their mental health is already under pressure. For example, research has shown that family caregivers experience ongoing feelings of guilt, burden and depression.^{29–31} There are individual differences in how family caregivers cope with caring for a person with dementia; these differences depend on various personal characteristics and circumstances. Resilience is one such characteristic, as it can serve as a protective barrier against developing certain psychological problems, including during the visitor ban.^{28,32} As such, resilience among family caregivers has been defined as a positive adaptation to adversity.³³

In the current study, the relationship between the visiting frequency and types of activities performed by family caregivers of people with dementia in LTCFs before the visitor ban and their experiences of worrying and loneliness during the ban was examined. Two possible moderators in this relationship, i.e., the contact

frequency during the visitor ban and the resilience of family caregivers, were studied as well.

Family caregivers might have experienced a loss of control and therefore increased worrying and loneliness during the visitor ban.²⁷ because they were unable to monitor or be involved in the provision of care.²³ Furthermore, family caregivers may have experienced more loneliness because they missed the company of the person with dementia²⁷ and there was less continuity in their relationship.^{25,26} The relationship between the visiting frequency before and worrying and loneliness during the visitor ban might have been moderated by how much contact family caregivers still had with the person with dementia during the visitor ban (for example, by telephone or video calls). Having more frequent contact might have helped family caregivers stay informed about how the person with dementia was doing and how they were being cared for. This could have in turn reduced worrying and loneliness. Resilience was a potential moderator as well, as it could have helped family caregivers better adapt to the changed situation, thereby mitigating worrying and loneliness. Fig. 1 depicts the hypothesized relationships between the investigated variables.

Hypothesis 1: More family involvement before the visitor ban leads to more worries of family caregivers during the visitor ban. This relationship is moderated by the frequency of contact during the visitor ban and the resilience of the family caregiver.

Hypothesis 2: More family involvement before the visitor ban leads to more experienced loneliness in family caregivers during the visitor ban. This relationship is moderated by the frequency of contact during the visitor ban and the resilience of the family caregiver.

Methods

Design

In this cross-sectional study, data were collected 6–10 weeks after the imposition of the national visitor ban, which lasted from April 30 until May 27, 2020, using a semi-open online survey administered to family caregivers of LTCF residents with dementia.¹³ This study was part of a larger study that aims to examine the impact of social isolation during the COVID-19 pandemic on socially vulnerable populations (www.coronatiijden.nl). Medical ethical approval for the study




Fig. 1. Models investigating the relationship between the involvement of family caregivers in LTCFs for people with dementia before the visitor ban and their worries (direct effect) and loneliness (direct effect), with moderators contact frequency during the visitor ban and the resilience of family caregivers (hypothesis 1 and 2).

was provided by the Medical Ethical Committee of Amsterdam's Institute for Social Science Research.

Sample and procedure

A total of 357 care organizations, registered nationally as nursing home care providers³⁴ under the Chronic Care Act (WLZ),³⁵ were invited by email to participate in the study. Information about the study and a link to the digital survey were provided. The care organizations were asked to distribute the information to the first contact persons of residents under their care, most of whom were family members, i.e. children, spouses or other family members, and a small number of friends, neighbors or a legal guardian. In this paper they are referred to as family caregivers to stress the personal relationship with the person with dementia. No confirmation of participation was sought from the care organizations. All data was gathered anonymously, a distinction between care organizations cannot be made. Informed consent from participants was obtained digitally prior to the start of the online survey.

This study used data from a sub-sample of participants. Only data provided by family caregivers who indicated dementia or memory problems as the main reason for the admission of their relative to the LTCF were included in the analyses ($N = 958$). The participants provided information about the type of LTCF in which the person with dementia resided in. These data indicated that 53.6% of people with dementia lived in a nursing home, 44.0% lived in a residential care home, and 2.4% lived in another type of LTCF (e.g., care farm). Of the LTCFs, 36.2% were considered to be of small size (<40 residents), 37.2% of medium (40–100 residents), and 20.7% of large size (>100 residents). The size of the LTCF was not indicated for the remaining 5.9%. Most LTCFs were located in regions where the COVID-19 infection rate during data collection was high (64.4%), as opposed to low (14.8%) and medium (20.8%).

Measures

Demographic information about the participants was gathered, i.e., age, gender, relationship to the person with dementia, age and gender of the person with dementia.

Independent variables

The frequency with which family caregivers visited the person with dementia before the visitor ban was measured with the following question: "How often did you visit your relative before the corona crisis in the LTCF?". Answer options were "every day" (high frequency), "at least once a week" (moderate frequency), "at least once a month" (low frequency), and "never, less than once a year and once or multiple times a year". Participants who did not visit or who visited only a few times per year were excluded from analyses, because they were considered to be incidental visitors.

Participants were asked to indicate, from a list of 11 activities, which activities they used to undertake with the person with dementia while visiting before the visitor ban. Participants had the option to add alternative activities to the list. As agreed upon by the researchers, the activities were categorized into social activities or task-related activities. Table 2 includes a list of these social and task-related activities.

Moderators

Contact frequency during the visitor ban was measured by asking participants "How often did you speak with your relative in the past four weeks?". Answer options were "daily", "multiple times a week", "weekly", "once in two weeks", "once" and, "never". These options were combined into three categories for the purpose of readability, namely high frequency (daily or multiple times a week), moderate

frequency (weekly), and low frequency (once in two weeks or once or never).

Two items of the translated version of the Brief Resilience Scale^{36,37} were used because they specifically measure resilience in stressful times: "I have a hard time making it through stressful events" (reverse-coded item) and "It does not take me long to recover from a stressful event". The mean "resilience in stressful times" score was calculated from the two items. The scores ranged from 1 to 5. Higher scores indicated a higher resilience level.

Dependent variables

The extent to which family caregivers worried about the person with dementia in the past four weeks was assessed with nine items formulated by the research consortium and based on research on the impact of the COVID-19 pandemic on people with severe mental illnesses.³⁸ The scores ranged from 0 (never) to 5 ((almost) always). A factor analysis was performed on these nine items. All but one item loaded on one factor; these eight items explained 48.5% of the variance. As the reliability of this eight-item scale, Cronbach's alpha = 0.841, was sufficient, it was used to assess the extent to which family caregivers worried about the person with dementia. The topics included in the scale were worries about the physical health of the person with dementia, being unable to visit the person with dementia in the LTCF, loneliness and unrest of the person with dementia, whether care for the person with dementia was of high quality, the day structure of the person with dementia, the possibility of the person with dementia passing away before family caregivers were able to visit them again, and the person with dementia no longer recognizing their family caregivers after the visitor ban had been lifted.

Loneliness was measured with the loneliness scale developed by de Jong Gierveld.³⁹ The scale consists of 11 items. Answer options are "no" (0), "more or less"¹ and "yes".¹ The total score, the sum of the dichotomized scores, ranged between 0 and 11. Higher scores indicated higher levels of loneliness.

Data analysis

Data were analyzed using SPSS version 27. Multiple regression analyses were performed to examine the relationship between the independent variables and the dependent variable. To create the interaction term, the independent and dependent variables were multiplied. The resilience score was centered to prevent multicollinearity.⁴⁰ Statistical significance was determined at $p < 0.05$. For each outcome measurement, i.e., worrying and loneliness, a separate model was investigated.

Results

Demographics

The mean age of the family caregivers was 60.30 (SD = 8.95, range between 16 and 89), and 71.7% were female. Three quarters (75.7%) of family caregivers indicated that the person with dementia was their father or mother. Spouses or partners represented 10.3% of the sample, while 14.0% had another type of relationship with the person with dementia (for example, other family members, friends, neighbors or legal guardians). The mean age of the person with dementia to whom family caregivers were related was 85.67 (SD=7.91, range between 46 and 105), and 73.5% were female. The demographics of the participants are presented in Table 1.

Independent variables

Of the family caregivers, 57.0% carried out social activities as well as task-related activities before the visitor ban, whereas 43.0% only engaged in social activities (Table 1). Table 2 shows the types of

Table 1

Family caregivers' and people with dementia demographics and scores on independent, dependent and moderator variables (N = 958).

Characteristics	Family caregivers
Mean age (SD)	60.30 (8.95)
Gender, n (%)	
Male	271 (28.3%)
Female	687 (71.7%)
Relationship to the resident, n (%)	
Spouse/partner	99 (10.3%)
Son/daughter	725 (75.7%)
Other (e.g. other family member, friend, neighbor, legal guardian)	134 (14.0%)
Visiting frequency before the visitor ban, n (%)	
Low (monthly, less than weekly)*	98 (10.2%)
Moderate (weekly, less than daily)	672 (70.1%)
High (daily)	188 (19.6%)
Types of activities performed while visiting before the visitor ban, n (%)	
Social activities	412 (43.0%)
Social and task related activities	546 (57.0%)
Contact frequency during the visitor ban, n (%)	
Low (less than weekly)	302 (31.5%)
Moderate (weekly)	221 (23.1%)
High (more than weekly)	435 (45.4%)
Mean resilience score (SD)	3.52 (0.76)
Mean worrying score (SD)	3.31 (0.79)
Mean loneliness score (SD)	6.84 (1.81)

* Respondents who paid no or few visits per year were excluded from the analyses.

activities undertaken by family caregivers before the visitor ban, their categorization into social versus task-related activities, and the number and percentage of family caregivers who undertook the aforementioned activities.

Moderators

The frequency with which the family caregivers contacted the person with dementia during the visitor ban was high for 45.4% of the caregivers, moderate for 23.1%, and low for 31.5% (Table 1). During the visitor ban, contact frequency increased for 17% of family caregivers and decreased for 25% as compared to visiting frequency before the ban. Family caregivers had alternative means of contact with the person with dementia via telephone (56.7%), video calls

Table 2

Types of activities family caregivers did before the visitor ban, their categorization into social versus task-related activities and the number and percentage of family caregivers that performed the activities (N=958)

Types of activities	Number of family caregivers that performed the activity (%)
<i>Social activities</i>	
Talking	875 (91.3)
Walking	570 (59.5)
Eating/drinking together	535 (55.8)
Going somewhere together (e.g. shopping/ outside/elsewhere)	437 (45.6)
Television/radio	254 (26.5)
Playing games together	197 (20.6)
Enjoyable activities (e.g. looking at pictures, reading or being read to, singing/music, etc.)	49 (5.1)
Religious activity	3 (0.3)
<i>Task-related activities</i>	
Sorting out the laundry	346 (36.1)
Personal care (e.g. hair, shaving, hands, etc.)	277 (28.9)
Preparing clothes	222 (23.2)
Helping care staff	121 (12.6)
Helping with meals	106 (11.1)
Other (e.g. doing administration, organizing things)	33 (3.4)

(53.2%), outside windows (29.5%), outside areas (23.7%), digital messages or social media (12.5%) or other ways (12.2%, for example, mail or hospital visits). Participants' mean resilience score was 3.52 (SD=0.76), ranging from 1 to 5 (Table 1).

Dependent variables

The mean score on the worrying scale was 3.31 (SD = 0.79), ranging from 1 to 5, as presented in Table 1. On the loneliness scale, participants scored a mean of 6.84 (SD = 1.81), ranging between 2 and 11 (Table 1).

Regression analyses

The final models that best fit the data are presented in Table 3. The final model for worrying examined the relationship between the involvement of family caregivers before and their worrying during the visitor ban, as moderated by contact frequency during the ban (Fig. 1). The results indicated that family caregivers who visited at least weekly before the visitor ban, worried more during the ban than those who visited less than weekly before the ban (direct effect). When family caregivers visited daily before and had weekly or more contact with the person with dementia during the visitor ban, they worried less as compared to family caregivers that visited less than weekly before the visitor ban and had contact less than weekly during the visitor ban (moderating effect). The final model, presented in Table 3, explained 10.3% of the variance (R²).

The final model for loneliness examined the relationship between the involvement of family caregivers before and their loneliness during the visitor ban, as moderated by the resilience of family caregivers (Fig. 1). For the dependent variable, loneliness, no significant direct effects were found. However, when family caregivers performed social as well as task-related activities before the visitor ban and were more resilient (moderating effect), they felt less lonely during the ban. The final model, presented in Table 3 explained 9.0% of the variance (R²).

Discussion/ conclusions

In this study, we examined the relationship between the visiting frequency of and types of activities undertaken by family caregivers of people with dementia before and their worrying and loneliness during the COVID-19 visitor ban in LTCFs. Two possible moderators in this relationship were studied: contact frequency during the visitor ban and the resilience of family caregivers.

In line with hypothesis 1, we found that family caregivers who visited more than weekly before the visitor ban worried more during the ban than those who visited monthly but less than weekly. This is in line with the findings from McCleary et al., as they found that the most stressful aspect of the visitor ban for family caregivers during the SARS outbreak was their inability to fulfill their responsibilities while they felt that their relatives' happiness and comfort depended on their presence. In addition, family caregivers in a study by McCleary et al. explained that their feelings during the visitor ban were similar to those they felt after the person with dementia moved into the LTCF, i.e., they experienced feelings of anger, worry, helplessness, grief, guilt, and failure.^{27,41} This is a possible explanation for the increased level of worrying we found in our study among family caregivers who visited the person with dementia more frequently before the visitor ban.

Further, this study revealed that family caregivers who visited the person with dementia more frequently before the visitor ban, worried less during the ban, when they had weekly or more contact. Family and professional caregivers were, however, compelled to find alternatives to physical visits during the visitor ban, e.g., telephone or video calls. When these methods allowed them to maintain frequent contact, they were likely better able to stay informed about the well-

Table 3

Results of moderated regression analyses with outcome variables 'worrying' and 'loneliness' and moderator variables 'contact frequency during the visitor ban' and 'resilience', corrected for confounders age and gender of the family caregiver and the relationship between the family caregiver and person with dementia ($N = 958$).

Variables	Final Worrying Model (β) ^{††}	Final Loneliness Model (β) ^{††}
<i>Visiting frequency before visitor ban</i>		
Low (reference) (monthly, less than weekly)		
Moderate (weekly, less than daily)	0.478***	-0.047
High (daily)	0.914***	-0.048
<i>Types of activities</i>		
Social activities only (reference)		
Task related and social activities	0.177	0.008
<i>Contact frequency during visitor ban</i>		
Low (reference) (less than weekly)		
Moderate (weekly)	0.222	
High (more than weekly)	0.251	
Resilience	nm [†]	-0.505
<i>Interaction effects (Visiting frequency before*contact during visitor ban)</i>		
Moderate before*moderate during	-0.448	nm [†]
Moderate before*high during	-0.409	nm [†]
High before*moderate during	-0.444*	nm [†]
High before*high during	-0.674**	nm [†]
<i>Interaction effects (Type of activities before*contact during visitor ban)</i>		
Task related and social activities*moderate during	0.006	nm [†]
Task related and social activities*high during	0.069	nm [†]
<i>Interaction effects (Visiting frequency before visitor ban*resilience)</i>		
Moderate before*resilience	nm [†]	-0.009
High before*resilience	nm [†]	-0.209
<i>Interaction effects (Type of activities*resilience)</i>		
Task related and social activities*resilience	nm [†]	-0.323*

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

[†] not in model

^{††} The confounders age and gender of the family caregiver and the relationship between the family caregiver and people with dementia have been taken into account in these analyses.

being of the person with dementia and monitor their care, at least to some degree, and therefore worry less. Although there is broad agreement that national visitor bans should be avoided in the future, in cases in which local visitor bans may be imposed, efforts should be taken to help family caregivers and people with dementia maintain ongoing contact.

Another outcome of our study was the moderating effect of resilience in the relationship between the types of activities in which family caregivers engaged before and their experience of loneliness during the visitor ban (hypothesis 2). When family caregivers undertook more diverse activities before the visitor ban, i.e., task-related and social activities, and they were more resilient, they experienced less loneliness during the ban. This is useful information given the harmful effects of loneliness on the health of family caregivers in general.⁴² Resilience seems to be a protective factor for a subgroup of family caregivers, i.e., those who undertake more diverse activities when visiting people with dementia in LTCFs. This also means that non-resilient family caregivers who generally only undertake social activities are more susceptible to loneliness. LTCFs should therefore consider the resilience of family caregivers of people with dementia and specifically this subgroup of family caregivers on occasions in which they have less contact with the person with dementia.

In our sample, one out of ten family caregivers were partners or spouses of the person with dementia. This might have impacted the results, as spouses may experience more loneliness than children or caregivers who have another type of relationship with the person with dementia. Spouses of people with dementia are often older themselves, and older people are known to be more at risk of experiencing loneliness or social isolation, especially when self-isolation measures are in effect, such as those imposed during the COVID-19 pandemic.¹⁵ It is possible that during the visitor ban, spouses who

used to visit their partner with dementia in the LTCF as part of their daily routine experienced greater loneliness, because they were prohibited from visitations. Therefore, in future analyses, a differentiation could be made between spousal caregivers and children of people with dementia to investigate whether they indeed experience disproportionately greater loneliness.

Although the models developed in this study explained a fair amount of variance, other factors, which we did not investigate, might also have had an effect on the worrying and loneliness experienced by family caregivers during the visitor ban. These factors include communication between family caregivers and care staff, as well as the provision of information. During the visitor ban, family caregivers depended on care staff to inform them about how the person with dementia was doing. Research has shown that informal communication between family caregivers and staff is essential for cultivating trust in staff and promoting family involvement.⁴³ To keep family caregivers engaged, care staff need to be proactive and creative.^{44,45} However, in this study, the extent to which this occurred among LTCF care staff members could have varied. COVID-19 outbreaks, for instance, could have influenced the degree to which care staff communicated with family caregivers. More specifically, such outbreaks could have made it more difficult for care staff to sufficiently inform family caregivers. The results indicate that regular contact between family caregivers and people with dementia is important and that such contact is often supported by care staff. Therefore, future research might explore the relationship between communication between family caregivers and care staff in times of crisis and the effects of this communication on the mental health of family caregivers.

The current study was unique in that we collected data during the COVID-19 visitor ban in the Netherlands. To our knowledge, very few studies collected data while the visitor ban was still active. Furthermore, we also collected retrospective data about the situation in LTCFs before the visitor ban, which made it possible to compare data about this situation before and during the visitor ban. Also, a substantial number of family caregivers who participated in the study lived in different regions in the Netherlands. Additionally, the explained variance of both investigated models (10.3% and 9.0%) was quite high considering the large scale of other factors that might contribute to worrying or loneliness, like dealing with uncertainty,⁴⁶ having diminished social contacts in times of crisis, e.g., the COVID-19 lockdown,⁴⁷ and unfamiliarity with digital tools for staying connected with the person with dementia.⁴⁸

In this study, we measured the visiting frequency before and contact frequency during the visitor ban differently. Concerning visiting frequency before the visitor ban, only physical visits were included, not contact in the form of, for example, telephone or video calls. This could mean that contact frequency before the visitor ban could have been higher if other types of contact were included. Regardless, during the visitor ban, family caregivers had less contact with the person with dementia. It is possible that if this construct had been measured differently, we would have found that contact frequency declined even more.

Further, we did not account for the way in which family caregivers had contact with the person with dementia during the visitor ban nor the duration of such interaction. For example, family caregivers might have depended on others to help them with video calls. This might also have been the case for people with dementia, who likely needed help from care staff, especially those with more advanced dementia. Time and tools must be made available to them as well. The duration of visits before and contact during the visitor ban also remains unclear. It was expected that experiences between family caregivers who visited the person with dementia before, on a daily basis and for hours, but who only spoke to them for five minutes at a time during the visitor ban via video calls might differ from family caregivers who

visited once a week for an hour before and called them for one hour per week during the visitor ban. Thus, more in-depth analyses are needed to investigate whether the type and duration of contact also influenced loneliness and worrying among family caregivers.

The extraordinary circumstances of the COVID-19 pandemic and the consequent LTCF visitor ban necessitated the development of a less traditional study design. This in turn imposed some limitations on the current study. To lower the threshold for participation as much as possible, LTCFs were permitted to participate fully anonymously and were not required to notify the research team about their participation. It seems that LTCFs in regions with high infection rates participated the most (64.4%), perhaps because they understood the urgency of contributing to research on the unique circumstances of the COVID-19 pandemic. Another consequence of the novel research design was that no response rates were available, and it is unclear how family caregivers were nested within LTCFs, i.e., multilevel analyses were not possible. This might have affected the research outcomes. Additionally, the policies and degrees of commitment of care staff to informing and involving family caregivers might also have differed between LTCFs. That said, as the visitor ban was imposed on the national level, the circumstances were likely the same in all participating LTCFs. In order to minimize the burden for participants during this stressful period, deliberate choices were made in the questionnaire to limit its length. In the case of the resilience measurements, this resulted in the inclusion of only two items of the Brief Resilience scale that deemed most important for our study purposes.

The results of this work demonstrate that a visitor ban in LTCFs is undesirable not just for people with dementia but also for care staff and family caregivers. Although there is widespread agreement that national visitor bans in LTCFs should be avoided in the future, temporary and local bans might sometimes still be necessary. If and when this happens, adequate attention should be given to the mental health of not only people with dementia and care staff but also family caregivers as well. The results of this study suggest that to reduce worrying among family caregivers, LTCFs should facilitate ongoing contact with people with dementia, specifically for highly involved family caregivers. Also, non-resilient family caregivers who generally only undertake social activities are more prone to loneliness. It is therefore advisable for healthcare and welfare professionals to reach out to this group to help them overcome their loneliness.

Acknowledgements

This study was funded by the The Netherlands Organization for Health Research and Development (ZonMw) (grant no. 10150062010007). The funding source had no involvement in determining the study design; in the collection, analysis and interpretation of data; in the writing of the articles; and in the decision to submit it for publication.

We would like to thank the consortium members Stephanie Steinmetz, Elske Stolte, Theo G. van Tilburg, and Danny H. de Vries for their contributions within the larger study on the impact of social isolation in times of the COVID-19 pandemic on socially vulnerable populations (www.coronatijden.nl).

References

- Brown EE, Kumar S, Rajji TK, Pollock BG, Mulsant BH. Anticipating and mitigating the impact of the COVID-19 pandemic on Alzheimer's disease and related dementias. *Am J Geriatr Psychiatry* [Internet]. 2020;28(7):712–721. <https://doi.org/10.1016/j.jagp.2020.04.010>. Available from.
- World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report – 51 March 11, 2020. 2020.
- Livingston G, Weidner W. COVID-19 and dementia: Difficult decisions about hospital admission and triage [Internet]. 2020. Available from: <https://www.alz.co.uk/news/adi-releases-position-paper-on-covid-19-and-dementia>.
- Yen MY, Schwartz J, King CC, Lee CM, Hsueh PR. Recommendations for protecting against and mitigating the COVID-19 pandemic in long-term care facilities. *J Microbiol Immunol Infect* [Internet]. 2020;53(3):447–453. <https://doi.org/10.1016/j.jmii.2020.04.003>. Available from.
- McMichael TM, Currie DW, Clark S, Pogosjans S, Kay M, Schwartz NG, et al. Epidemiology of COVID-19 in a long-term care facility in King County, Washington. *N Engl J Med*. 2020;382(21):2005–2011.
- Ayalon L, Zisberg A, Cohn-Schwartz E, Cohen-Mansfield J, Perel-Levin S, Bar-Asher Siegal E. Long-term care settings in the times of COVID-19: challenges and future directions. *Int Psychogeriatrics*. 2020;32(10):1239–1243.
- Comas-Herrera A, Zalakain J, Lemmon E, Henderson D, Litwin C, Hsu AT, et al. Mortality associated with COVID-19 in care homes: international evidence [Internet]. 2020. Available from: <https://lccovid.org/wp-content/uploads/2020/10/Mortality-associated-with-COVID-among-people-living-in-care-homes-14-October-2020-3.pdf>.
- Wang H, Li T, Barbarino P, Gauthier S, Brodaty H, Molinuevo JL, et al. Dementia care during COVID-19. *Lancet*. 2020;395(10231):1190–1191.
- Actiz, Alzheimer Nederland, LOC, NIP, Verenso, V&VN, et al. Handreiking bezoek en sociaal contact. Corona in verpleeghuizen [Guidance for visits and social contacts. Corona in LTCFs] [Internet]. Utrecht; 2020. Available from: <https://www.actiz.nl/nieuws/nieuwe-handreiking-aanpak-verpleeghuiszorg-bij-corona-gereed>.
- Verbeek H, Gerritsen DL, Backhaus R, de Boer BS, Koopmans RTCM, Hamers JPH. Allowing visitors back in the nursing home during the COVID-19 crisis: a Dutch national study into first experiences and impact on well-being. *J Am Med Dir Assoc* [Internet]. 2020;21(7):900–904. <https://doi.org/10.1016/j.jamda.2020.06.020>. Available from.
- Dichter MN, Sander M, Seismann-Petersen S, Köpke S. COVID-19: it is time to balance infection management and person-centered care to maintain mental health of people living in German nursing homes. *Int Psychogeriatrics*. 2020;32(10):1157–1160.
- Society British Geriatrics. COVID-19 : Managing the COVID-19 pandemic in care homes for older people Key recommendations [Internet]. *British Geriatrics Society*. 2020. Available from: <https://www.bgs.org.uk/resources/covid-19-managing-the-covid-19-pandemic-in-care-homes>.
- van der Roest HG, Prins M, van der Velden C, Steinmetz S, Stolte E, van Tilburg TG, et al. The impact of COVID-19 measures on well-being of older long-term care facility residents in the Netherlands. *J Am Med Dir Assoc* [Internet]. 2020;21(11):1569–1570. <https://doi.org/10.1016/j.jamda.2020.09.007>. Available from.
- Courtin E, Knapp M. Social isolation, loneliness and health in old age: a scoping review. *Health Soc Care Community*. 2017;25(3):799–812.
- Armitage R, Nellums LB. COVID-19 and the consequences of isolating the elderly. *Lancet Public Heal* [Internet]. 2020;5(5):e256. [https://doi.org/10.1016/S2468-2667\(20\)30061-X](https://doi.org/10.1016/S2468-2667(20)30061-X). Available from <http://dx.doi.org/>.
- van der Geest M. Onderzoek Amsterdam UMC: verpleeghuisbewoners kregen geen knauw door bezoekverbod. de Volkskrant [Internet]. 2020 Sep 26; Available from: <https://www.volkskrant.nl/nieuws-achtergrond/onderzoek-amsterdam-verpleeghuisbewoners-kregen-geen-knauw-door-bezoekverbod~b20001de?referer=https%3A%2F%2Fwww.google.com%2F>.
- Leontjevas R, Knippenberg IAH, Smalbrugge M, Plouvier AOA, Teunisse S, Bakker C, et al. Challenging behavior of nursing home residents during COVID-19 measures in the Netherlands. *Ageing Ment Heal* [Internet]. 2020;1–6. <https://doi.org/10.1080/13607863.2020.1857695>. Available from.
- Gaugler JE, Pearlin LI, Zarit SH. Family involvement following institutionalization: modeling nursing home visits over time. *Int J Aging Hum Dev* [Internet]. 2003;57(2):91–117. Available from: <http://journals.sagepub.com/doi/10.2190/8MNF-QMA3-A5TX-6QQ3>.
- Gladstone JW, Dupuis SL, Wexler E. Changes in family involvement following a relative's move to a long-term care facility. *Can J Aging / La Rev Can du Vieil* [Internet]. 2006;25(1):93–106. Available from: http://www.journals.cambridge.org/abstract_S071498080002270.
- Fast CT, Houlihan D, Buchanan JA. Developing the family involvement questionnaire-long-term care: a measure of familial involvement in the lives of residents at long-term care facilities. *Gerontologist*. 2019;59(2):E52–E55.
- Gaugler JE. Family involvement in residential long-term care: a synthesis and critical review. *Ageing Ment Heal*. 2005;9(2):105–118.
- Puurveen G, Baumbusch J, Gandhi P. From family involvement to family inclusion in nursing home settings: a critical interpretive synthesis. *J Fam Nurs*. 2018;24(1):60–85.
- Davies S, Nolan M. Making it better": self-perceived roles of family caregivers of older people living in care homes: a qualitative study. *Int J Nurs Stud*. 2006;43(3):281–291.
- Roberts AR, Ishler KJ, Adams KB. The predictors of and motivations for increased family involvement in nursing homes. *Gerontologist*. 2020;60(3):535–547.
- Lloyd J, Patterson T, Muers J. The positive aspects of caregiving in dementia: A critical review of the qualitative literature. *Dementia*. 2016;15(6):1534–1561.
- Forsund LH, Kiik R, Skovdahl K, Ytrehus S. Constructing togetherness throughout the phases of dementia: a qualitative study exploring how spouses maintain relationships with partners with dementia who live in institutional care. *J Clin Nurs*. 2016;25(19–20):3010–3025.
- McCleary L, Munro M, Jackson L, Mendelsohn L. Impact of SARS visiting restrictions on relatives of long-term care residents. *J Soc Work Long Term Care*. 2005;3(3–4):3–20.
- Hughes MC, Liu Y, Baumbach A. Impact of COVID-19 on the health and well-being of informal caregivers of people with dementia: a rapid systematic review. *Gerontol Geriatr Med*. 2021;7: 233372142110201.

29. Collins RN, Kishita N. Prevalence of depression and burden among informal caregivers of people with dementia: A meta-analysis. *Ageing Soc.* 2019;1–38.
30. Gaugler JE, Mittelman MS, Hepburn K, Newcomer R. Clinically significant changes in burden and depression among dementia caregivers following nursing home admission. *BMC Med [Internet]*. 2010;8(1):85. Available from; <http://www.biomed-central.com/1741-7015/8/85>.
31. Pinquart M, Spouses Sørensen S. Adult children, and children-in-law as caregivers of older adults: a meta-analytic comparison. *Psychol Aging.* 2011;26(1):1–14.
32. Altieri M, Santangelo G. The psychological impact of COVID-19 pandemic and lockdown on caregivers of people with dementia. *Am J Geriatr Psychiatry [Internet]*. 2021;29(1):27–34. <https://doi.org/10.1016/j.jagp.2020.10.009>. Available from.
33. de Dias R, Santos RL, Sousa MFB, Nogueira MML, Torres B, Belfort T, et al. Resilience of caregivers of people with dementia: a systematic review of biological and psychosocial determinants. *Trends Psychiatry Psychother [Internet]*. 2015;37(1):12–19. Available from; http://www.scielo.br/scielo.php?script=sci_arttext&pid=S2237-60892015000100012&lng=en&tlng=en.
34. Vektis Business Intelligence Center for Healthcare. Search the AGB register [Internet]. 2020 [cited 2020 Apr 1]. Available from: <https://www.vektis.nl/agb-register/zoeken#resultaten>.
35. Government of the Netherlands. Chronic Care Act (WLZ) [Internet]. 2021 [cited 2021 May 18]. Available from: <https://www.government.nl/topics/nursing-homes-and-residential-care/chronic-care-act-wlz>.
36. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The brief resilience scale: Assessing the ability to bounce back. *Int J Behav Med.* 2008;15(3):194–200.
37. Windle G, Bennett KM, Noyes J. A methodological review of resilience measurement: Gregory Bateson ecology mind scales. *Health Qual Life Outcomes [Internet]*. 2011;9(8). Available from; <http://www.hqlo.com/content/9/1/8>.
38. De Lange A, Hulsbosch L, Knispel A. *Impact of the coronacrisis on people with severe mental illnesses [Impact van de coronacrisis op mensen met ernstige psychische aandoeningen]*. Utrecht; 2020.
39. de Jong Gierveld J, van Tilburg T. Manual of the Loneliness Scale 1999 (Updated from the printed version: 19-05-2020) [Internet]. Amsterdam; 1999. Available from: https://home.fsw.vu.nl/tg.van.tilburg/manual_loneliness_scale_1999.html#short_version.
40. Iacobucci D, Schneider MJ, Popovich DL, Bakamitsos GA. Mean centering helps alleviate “micro” but not “macro” multicollinearity. *Behav Res Methods [Internet]*. 2016;48(4):1308–1317. <https://doi.org/10.3758/s13428-015-0624-x>. Available from.
41. Seddon K, Jones K, Boyle M. Committed to caring: Carer experiences after a relative goes into nursing or residential care. *Qual Ageing.* 2002;3(3):16–26.
42. Kovaleva M, Spangler S, Clevenger C, Hepburn K. Chronic stress, social isolation, and perceived loneliness in dementia caregivers. *J Psychosoc Nurs Ment Health Serv.* 2018;56(10):36–43.
43. Hoek LJ, van Haastregt JC, de Vries E, Backhaus R, Hamers JP, Verbeek H. Partnerships in nursing homes: how do family caregivers of residents with dementia perceive collaboration with staff? *Dementia.* 2020;0(0):1–18.
44. O’Shea E. Remembering people with dementia during the COVID-19 crisis. *HRB Open Res.* 2020;3.
45. Mills J, Li C, Fullerton S, Chapman M, Jap J, Sinclair C, et al. Staying connected and informed: Online resources and virtual communities of practice supporting palliative care during the novel coronavirus pandemic. *Prog Palliat Care [Internet]*. 2020;28(4):251–253. <https://doi.org/10.1080/09699260.2020.1759876>. Available from.
46. Unson C, Flynn D, Glendon MA, Haymes E, Sancho D. Dementia and caregiver stress: an application of the reconceptualized uncertainty in illness theory. *Issues Ment Health Nurs.* 2015;36(6):439–446.
47. Greenberg NE, Wallick A, Brown LM. Impact of COVID-19 pandemic restrictions on community-dwelling caregivers and persons with dementia. *Psychol Trauma Theory, Res Pract Policy.* 2020;12:S220–S221.
48. Gijzen M, Shields-Zeeman L, Kleinjan M, Kroon H, van der Roest H, Bolier L, et al. The bittersweet effects of COVID-19 on mental health: results of an online survey among a sample of the Dutch population five weeks after relaxation of lockdown restrictions. *Int J Environ Res Public Health.* 2020;17(23):1–11.