

EUR Research Information Portal

Novel Risk Markers for Type 2 Diabetes : Inflammation, Body Fat and Sex Hormones

Publication status and date:

Published: 25/04/2018

Document Version

Other version

Citation for the published version (APA):

Brahimaj, A. (2018). *Novel Risk Markers for Type 2 Diabetes : Inflammation, Body Fat and Sex Hormones*. [Doctoral Thesis, Erasmus University Rotterdam]. Erasmus Universiteit Rotterdam (EUR).

[Link to publication on the EUR Research Information Portal](#)

Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.

In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: openaccess.library@eur.nl. Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

Propositions of this thesis:

Novel Risk Markers for Type 2 Diabetes

Inflammation, Body Fat and Sex Hormones

1. Both pro-inflammatory and anti-inflammatory markers are associated with the progression from normoglycemia to type 2 diabetes and need for insulin therapy in a phase-specific manner. (This thesis)
2. High overall antioxidant capacity of diet may lead to lower levels of pro-inflammatory adipocytokines and higher levels of anti-inflammatory adipocytokines. (This thesis)
3. Serum apoCIII levels as well as apoCIII-to-apoA1 ratio are associated with incident type 2 diabetes, independent of known risk factors. (This thesis)
4. Combined metabolic indices that include anthropometric and laboratory measures are strong risk markers associated with incident type 2 diabetes. (This thesis)
5. Dehydroepiandrosterone may have implications in preventive interventions for type 2 diabetes. (This thesis)
6. Markers of inflammation and immunity enable differentiation between the early preclinical and clinical phases of the disease, disease complications, and progression. (Grossmann V. et al. *Diabetes Care* 2015;38:1356-1364)
7. Observationally, low levels of HDL cholesterol are consistently associated with increased risk of type 2 diabetes. (Hasse CL et al. *Diabetes* 2015;64:3328-3333)
8. Change in visceral adiposity independently predicts risk of developing type 2 diabetes over 10 years. (Wander PL. et al. *Diabetes Care* 2013;36:289-293)
9. A sufficient intake of antioxidants plays a role in type 2 diabetes prevention. (Montonen J. et al. *Diabetes Care* 2004;27:1845)
10. It is an integral part of good doctoring to ask not only, 'What is the diagnosis, and what is the treatment?' but also, 'Why did this happen, and could it have been prevented?' (Rose G. *Int J Epidemiol* 1985;14:32-38)
11. Nobody has ever measured, not even poets, how much the heart can hold. (Zelda Fitzgerald)