

EUR Research Information Portal

The pathophysiological consequences of somatostatin receptor internalization and resistance

Published in:
Endocrine Reviews

Publication status and date:
Published: 01/01/2003

DOI (link to publisher):
[10.1210/er.2000-0001](https://doi.org/10.1210/er.2000-0001)

Document Version
Publisher's PDF, also known as Version of record

Citation for the published version (APA):
Hofland, L., & Lamberts, S. W. J. (2003). The pathophysiological consequences of somatostatin receptor internalization and resistance. *Endocrine Reviews*, 24(1), 28-47. <https://doi.org/10.1210/er.2000-0001>

[Link to publication on the EUR Research Information Portal](#)

Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.

In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: openaccess.library@eur.nl. Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

The Pathophysiological Consequences of Somatostatin Receptor Internalization and Resistance

LEO J. HOFBLAND AND STEVEN W. J. LAMBERTS

Department of Internal Medicine, Erasmus Medical Center, 3015 GD Rotterdam, The Netherlands

Somatostatin receptors expressed on tumor cells form the rationale for somatostatin analog treatment of patients with somatostatin receptor-positive neuroendocrine tumors. Nevertheless, although somatostatin analogs effectively control hormonal hypersecretion by GH-secreting pituitary adenomas, islet cell tumors, and carcinoid tumors, significant differences are observed among patients with respect to the efficacy of treatment. This may be related to a differential expression of somatostatin receptor subtypes among tumors. In addition, the property of somatostatin receptor subtypes to undergo agonist-induced internalization has important con-

sequences for visualizing, as well as for therapy, of receptor-positive tumors using radioisotope- or chemotherapeutic-compound-coupled somatostatin analogs. This review covers the pathophysiological role of somatostatin receptor subtypes in determining the efficacy of treatment of patients with somatostatin receptor-positive tumors using somatostatin analogs, as well as the preclinical and clinical consequences of agonist-induced receptor internalization for somatostatin receptor-targeted radio- and chemotherapy. Herein, the development and potential role of novel somatostatin analogs is discussed. (Endocrine Reviews 24: 28–47, 2003)

- I. General Introduction
 - A. SS and sst subtypes
 - B. SS receptor subtype expression in normal and tumorous human tissues
 - C. Agonist-induced internalization of sst subtypes
- II. Consequences of sst Internalization for sst-Targeted Radiotherapy or Chemotherapy of sst-Positive Tumors
 - A. Preclinical evidence for internalization of radiolabeled SS-analogs
 - B. SS receptor-targeted radiotherapy
 - C. SS receptor-targeted chemotherapy: preclinical evidence
- III. Tachyphylaxis and Resistance to SS
 - A. Introduction
 - B. Tachyphylaxis of pathological hormone secretion
 - C. Escape from antiproliferative effects
 - D. Mechanisms of tachyphylaxis and resistance
 - E. New developments
 - F. Conclusions
- IV. Summary

I. General Introduction

SINCE ITS DISCOVERY in 1973 by Guillemin and Gerich (1), knowledge of the functional role of somatostatin (SS) in regulating neurotransmission in the brain, as well as in the regulation of secretion processes in the anterior pituitary gland, the pancreas, and the gastrointestinal tract, has increased considerably. In addition to playing an important regulatory role in neurotransmission and secretion, the peptide may control cell proliferation in normal and tumorous

tissues as well (2, 3). Between 1992 and 1994, five SS receptor (sst) subtype genes were cloned and characterized; they were code-named sst₁, sst₂, sst₃, sst₄, and sst₅ (4). The discovery of these genes initiated a large number of studies directed to elucidate their expression in SS-target tissues, their selectivity of binding of structural SS-analogs, and their coupling to the different second messenger systems known to be activated upon SS binding to its receptor. This has been reviewed extensively (5–9). The discovery of the sst subtype genes also initiated the development of a large series of novel SS-analogs that selectively bind to sst subtypes. Currently, a number of these sst subtype-selective analogs are being tested for their *in vivo* and *in vitro* potencies to modulate hormone secretion and/or cell proliferation (8, 10). The high density of SS receptors on human neuroendocrine tumors originating from normal SS target tissues has been used clinically to treat symptoms of hormonal hypersecretion in patients with GH- or TSH-secreting pituitary adenomas, as well as in patients harboring islet cell or carcinoid tumors with SS-analogs (11). However, although SS-analogs effectively control hormonal hypersecretion by neuroendocrine tumors, their effects are often transient, and considerable differences between patients harboring islet cell and carcinoid tumors exist with respect to the development of tachyphylaxis. In addition, the presence of a high density of SS receptors on human neuroendocrine tumors has allowed the development in 1989 (2, 12) of the technique of sst scintigraphy using radiolabeled SS-analogs to visualize sst-positive tumors *in vivo* (2, 13).

The above physiological and pathophysiological roles of SS and the presence of SS receptors on neuroendocrine tumors have been reviewed extensively. Much less attention has been paid to the clinical importance of sst internalization in determining the uptake of radiolabeled SS-analogs by sst-positive neuroendocrine tumors and the role of individual sst subtypes herein, as well as to the mechanisms in-

Abbreviations: CHO, Chinese hamster ovary; DOTA, tetraazacyclododecane tetraacetic acid; DTPA, diethylenetriamine pentaacetic acid; hsst, human sst; 5-HIAA, 5-hydroxyindolacetic acid; ¹⁷⁷Lu, ¹⁷⁷lutetium; MTC, medullary thyroid carcinoma; PRL, prolactin; SS, somatostatin; sst, SS receptor; wt, wild-type; ⁹⁰Y, yttrium-90.

volved in tachyphylaxis to SS-analog therapy. This manuscript gives an overview of the current knowledge on the internalization and cellular uptake of radiolabeled SS-analogs by sst-positive tumor cells and the involvement of endogenously expressed sst subtypes in this process, as well as the clinical consequences of sst internalization for sst-targeted radio- or chemotherapy. *Section III* of this review addresses the potential mechanisms involved in tachyphylaxis after long-term treatment of patients with neuroendocrine tumors with SS-analogs.

A. SS and sst subtypes

SS is a small cyclic peptide that is widely expressed throughout the central nervous system and peripheral tissues. In peripheral tissues, SS exerts predominantly inhibitory actions (14) on secretion processes, whereas the peptide acts as a neurotransmitter in both a stimulatory and inhibitory manner in the brain (15). SS is formed by proteolytic processing of larger precursor molecules, *i.e.*, prepro-SS and pro-SS. After cleavage of the pro-SS molecule, two biologically active forms of SS consisting of 14 (SS-14) or 28 (SS-28) amino acids are generated (16). SS-14 and SS-28 act via high-affinity G protein-coupled membrane receptors. Five sst subtypes have been cloned and characterized. The genes encoding the five sst subtypes are localized on different chromosomes (8). Via alternative splicing, two forms of the sst₂ receptor can be generated, *i.e.*, sst_{2A} and sst_{2B} (17, 18). The only difference between sst_{2A} and sst_{2B} is the length of their cytoplasmic tail. The five sst subtypes share a coupling to the second messenger systems known to be activated upon SS binding to its receptor. These systems include inhibition of adenylyl cyclase activity and activity of calcium channels, as well as stimulation of phosphotyrosine phosphatase or MAPK activity. This has been reviewed extensively (7–9). Although the inhibitory effects on adenylyl cyclase activity and on the influx of Ca²⁺ are linked to inhibition of secretion processes, the activation of phosphotyrosine phosphatase or MAPK activity may play a role in the regulatory effects that SS exerts on cell proliferation (2, 3, 10). The selective induction of apoptosis mediated via activation of sst₃ receptors is of particular interest in this respect. The role of the individual sst subtypes and the mechanism of action of the antiproliferative effects by SS have been reviewed recently (9). The five sst subtypes all bind SS-14 and SS-28 with high affinity but can be divided into two subclasses on their ability to bind structural octapeptide analogs of SS. The sst₁ and sst₄ receptors do not bind octapeptide SS-analogs, whereas sst_{2A}, sst₃, and sst₅ receptors display a high, low, and moderate affinity, respectively, toward octapeptide SS-analogs such as the clinically used octreotide and lanreotide (Table 1).

B. SS receptor subtype expression in normal and tumorous human tissues

Classical SS-target tissues such as the central nervous system, the anterior pituitary gland, and the pancreas express multiple sst subtypes. The expression of sst subtypes in the brain has only been studied extensively in rodent species. In the brain, mRNAs encoding for all five sst subtypes are expressed in a highly specific pattern (8). This regional, char-

TABLE 1. SS receptor subtype selectivity of binding of SS agonists

Agonist	Binding potency (IC ₅₀ values in nM)				
	sst ₁	sst ₂	sst ₃	sst ₄	sst ₅
SS-14	2.3	0.2	1.4	1.8	0.9
Octreotide	>1000	0.6	34.5	>1000	7
Lanreotide	>1000	0.8	107	>1000	5.2
Vapreotide ^a	>1000	5.4	31	45	0.7
MK-678 ^a	>1000	1.5	27	127	2
BIM-23268	18.4	15.1	61.6	16.3	0.4
BIM-23197	>1000	0.2	26.8	>1000	9.8
BIM-23244	>1000	0.3	133	>1000	0.7
SOM-230	9.3	1.0	1.5	>100	0.2

Data are derived from Refs. 8, 74, and 191.

^a Values represent K_i values in nM (Refs. 8, 187, 188, 191).

acteristic expression pattern of sst subtypes in the brain has recently been confirmed at the protein level by immunohistochemical techniques using sst subtype-specific antibodies (19). The adult human pituitary gland expresses sst₁, sst₂, sst₃, and sst₅ mRNAs, but not sst₄ mRNA (20). In addition, human pancreatic islet cells express all five sst subtype proteins, as determined by immunohistochemistry (21, 22). In human islets, sst₁, sst₂, and sst₅ receptors are the most abundantly expressed subtypes, with a high percentage of β -cells expressing sst₁ and sst₅, α -cells expressing sst₂, and δ -cells expressing sst₅ (22).

Neuroendocrine tumors, which often originate from SS-target tissues, frequently express a high density of SS receptors (23–26). The sst-expressing human tumors include pituitary adenomas, islet cell tumors, carcinoids, paragangliomas, pheochromocytomas, small cell lung cancers, and medullary thyroid carcinomas (MTCs), but also breast cancers and malignant lymphomas (24, 27). The sst subtype expression in different types of human cancers has been demonstrated at the mRNA level using *in situ* hybridization (28, 29), RNase protection assays, and RT-PCR (20, 27, 30–34). The majority of human sst-positive tumors simultaneously express multiple sst subtypes, although there is a considerable variation in sst subtype expression between the different tumor types and among tumors of the same type. Table 2 shows that sst₂ is the most abundantly expressed receptor subtype in the majority of tumors. Recent studies using antibodies raised against synthetic peptide sequences of the sst₁, sst₂, sst₃, and sst₅ receptor confirmed this variation in the expression of sst subtypes in different types of human tumors (Table 2; Refs. 35–38). The higher number of tumors expressing particular sst subtype mRNAs, compared with the number of tumors expressing sst subtype proteins, may be related to the higher sensitivity of techniques such as RT-PCR compared with immunohistochemistry. On the other hand, techniques such as RT-PCR might overestimate the real percentage of tumors expressing sst subtypes because blood vessels, immune cells, stromal and contaminating normal cells, which are present in or surround human tumors, may express sst subtypes as well (29, 39–41). The predominant expression of sst₂ receptors in human tumors forms the basis for the successful clinical application of octapeptide SS-analogs such as octreotide and lanreotide in controlling symptoms related to hormonal hypersecretion in patients with GH-secreting pituitary adenomas, islet cell tu-

TABLE 2. Expression of sst subtypes in human tumors

Tumor type	SS receptor subtype									
	sst ₁		sst ₂		sst ₃		sst ₄		sst ₅	
	mRNA	Protein	mRNA	Protein	mRNA	Protein	mRNA	Protein	mRNA	Protein
Pituitary tumor										
Somatotroph	44% (25)		96% (28)		44% (25)		5% (22)		86% (22)	
Lactotroph	84% (19)		63% (19)		35% (17)		6% (17)		71% (17)	
Nonfunctioning	38% (24)		75% (24)		43% (23)		13% (23)		48% (23)	
Corticotroph	56% (9)		67% (9)		25% (8)		0% (7)		86% (7)	
Neuroendocrine										
GEP tumors										
Carcinoid	76% (59)	88% (8)	80% (84)	78% (63)	43% (58)		68% (47)		77% (44)	
Gastrinoma	79% (28)	100% (5)	93% (28)	100% (8)	36% (28)		61% (23)		93% (28)	
Insulinoma	76% (21)		81% (21)		38% (21)		58% (19)		57% (21)	
Nonfunctioning	58% (24)		88% (24)		42% (24)		48% (21)		50% (24)	
ICT										
Renal cell cancer	85% (13)		100% (13)		0% (13)		50% (12)			
Breast cancer	33% (103)	52% (33)	99% (103)	48% (33)	38% (101)	48% (33)	23% (97)		18% (51)	
Meningioma	46% (24)		100% (24)		33% (24)		50% (12)		71% (14)	
Glioma	100% (7)		100% (7)		67% (6)		71% (7)		57% (7)	
Neuroblastoma	0% (6)		100% (15)		17% (6)					
Colorectal cancer	27% (41)		87% (41)		22% (41)		10% (41)		46% (41)	
MTC	29% (14)		79% (14)		36% (14)		0% (14)		64% (14)	
Pheochromocytoma	100% (11)	80% (5)	100% (11)	90% (20)	73% (11)		73% (11)		73% (11)	

Data on mRNA expression are derived from Refs. 20, 27, 30–34, 69, 70, 72, 192–199 and include studies using RT-PCR, Northern blotting, and *in situ* hybridization techniques. Data on protein expression include immunohistochemical studies using sst subtype-specific antibodies and are derived from Refs. 36–38 and 69. The values represent the percentage of tumors expressing the sst subtype among the tumors screened; the values between parentheses indicate the total number of tumors of the studies included. GEP, Gastroenteropancreatic; ICT, islet cell tumor; MTC, medullary thyroid carcinoma.

mors, or carcinoid tumors (2, 11), but also for the possibility to visualize sst-positive tumors using radiolabeled SS-analogs (see Section II). Knowledge of the sst subtype expression patterns in human neuroendocrine tumors may be very important for the development of the concept of sst-targeted radiotherapy or chemotherapy. As will be discussed below, sst subtypes differ in their ability to internalize receptor-bound ligand, which is a crucial step to direct a SS-analog-linked radioisotope or cytotoxic compound to the nucleus of the tumor cell.


C. Agonist-induced internalization of sst subtypes

Since the cloning of the five sst subtypes, the involvement of the individual sst subtypes in the process of receptor-mediated internalization of SS has been extensively investigated. Although differences have been reported between human and rat sst subtypes with respect to their dynamics of agonist-induced internalization, Section I.C is focused primarily on human sst subtypes and briefly summarizes their reported ability to undergo internalization after exposure to agonists. The mechanisms involved in receptor-mediated internalization of sst subtypes are not the focus of this review, and they have been reviewed elsewhere (8, 9, 42–46). In general, the mechanism and route of internalization of sst-agonist complexes follow those described for many other G protein-coupled receptors (47–50) and involve aggregation of the hormone receptor complex in specialized areas of the membrane, followed by internalization of the hormone-receptor complex via clathrin-coated, as well as uncoated, pits (47, 51). After internalization and pit formation, fusion of these vesicles with lysosomes occurs, resulting in hormone degradation or receptor recycling to the cell surface (Fig. 1; Refs. 49, 52, and 53).

The sst subtypes differentially internalize SS and SS-analogs (9). In Chinese hamster ovary (CHO)-K1 cells stably expressing one of the five human sst subtypes, sst₂, sst₃, sst₄, and sst₅ receptors displayed rapid (within minutes) agonist-dependent internalization of [¹²⁵I]LTT SS-28 ligand in a time- and temperature-dependent manner (54). Maximum internalization of the radioligand occurred within 60 min. The sst₃- and sst₅-expressing cells displayed the highest degree of internalization (78% and 66%, respectively), followed by sst₄ (29%) and sst₂ (20%). In contrast, the sst₁ subtype displayed only a very low amount (4%) of internalization. Another study using COS-7 cells transfected with the human sst₁ (hsst₁) or hsst_{2A} receptor subtypes (55) recently confirmed the low internalization rate of the sst₁ subtype. These investigators used confocal microscopy to analyze the fate of internalized novel fluorescent SS derivatives (43). In cells transfected with sst_{2A} receptors, up to 75% of specifically bound fluorescent ligand was recovered inside the cells within 60 min after agonist exposure, where it clustered into small endosome-like particles (55), whereas the capacity of internalization of SS via the sst₅ receptor was intermediate between sst₁ and sst_{2A} receptors (43). These particles increased in size over time, suggesting that the receptor-ligand complexes followed an endocytotic pathway.

Recent observations by Rocheville *et al.* (56) demonstrated that internalization of human sst subtypes can be determined by functional homo- and heterodimerization of sst subtypes as well. The hsst₁ receptors displayed no internalization of their selective ligand ¹²⁵I-SCH288, consistent with their inability to undergo agonist-induced internalization as a monofluorescent (57). However, when hsst₁ receptors were cotransfected with a c-tail deletion mutant of hsst₅, a slight but significant internalization of ¹²⁵I-SCH288 at 60 min was

FIG. 1. Schematic representation of intracellular routing of G protein-coupled receptors (GPCRs) after agonist activation. After agonist activation, GPCRs are phosphorylated (involving protein kinase A, protein kinase C, and GPCR kinases) and internalized, probably via the formation of clathrin-coated pits (involving β -arrestins). The internalized receptors are then directed to endosomes in which they are dephosphorylated. Subsequently, the receptors are recycled back to the plasma membrane as functional (resensitized) receptors. GPCR down-regulation results from lysosomal degradation of intracellular receptors, decreased mRNA and receptor protein synthesis, as well as increased degradation via mobilization of membrane receptors directly to the lysosomal compartment. L, Ligand; PP, phosphate group. [Adapted from Ref. 49.]



observed. Heterodimerization of epitope-tagged ss_{2A} and ss_3 receptors prevents agonist-induced endocytosis of ss_3 but not ss_{2A} receptors (58). Apart from changes in functionality of individual sst subtypes due to receptor dimerization, sst receptors may also form heterodimers with other G protein-coupled receptors, e.g., dopamine and opioid receptors (59, 60). Again, such heterodimers have properties different from the individual receptors. Therefore, the knowledge that homo- and heterodimerization of sst subtypes, and of sst subtypes with other G protein-coupled receptors, may influence the capacity of individual sst subtypes to undergo agonist-induced endocytosis clearly indicates the need to study internalization of sst subtypes endogenously expressed in sst-positive cells. Such studies will help to clarify the apparent discrepancies in internalization of specific sst subtypes. The above-described ability of sst subtypes to undergo agonist-induced internalization is an important characteristic of these receptors for transporting radiolabeled SS-analogs into the cell, thereby making sst-targeted radiotherapy a feasible approach to treat patients with neuroendocrine tumors expressing a high density of sst. In Section II, the preclinical evidence for internalization of radiolabeled SS-analogs, resulting in accumulation of intracellular radioactivity, by tumor cells endogenously expressing sst subtypes, is reviewed.

II. Consequences of sst Internalization for sst-Targeted Radiotherapy or Chemotherapy of sst-Positive Tumors

A. Preclinical evidence for internalization of radiolabeled SS-analogs

Human sst-positive tumors show a high uptake of [^{111}In -DTPA 0]octreotide at sst scintigraphy (13). Analysis of the uptake of [^{111}In -DTPA 0]octreotide by scintigraphy is preferably performed 24 h after the injection of the radiopharmaceutical (13). After 24 h, it is unlikely that the radioactivity in the tumors reflects cell membrane-bound ligand, but in fact,

more likely, represents internalized radioligand. Internalization of [^{111}In -DTPA 0]octreotide *in vivo* is also evidenced by our observations in rats, in which uptake of radioactivity in sst-positive organs, such as the pituitary gland and the pancreas, after the injection of the radiopharmaceutical, can be prevented by injection of excess unlabeled octreotide up to 10 min post injection, but not 20 min post injection. At that time, all radioactivity present in the sst-positive tissues probably reflects internalized radioligand (61). Direct evidence for internalization and subsequent subcellular distribution of radioisotopes delivered to the tumor cells using radiolabeled SS-analogs is presented from several *ex vivo* and *in vitro* autoradiographic studies. After incubation of human HT-29 colon carcinoma cells with the ^3H -labeled SS-analog TT-232, radioactivity was observed at the cell surface and cytoplasmic membranes, as well as the nucleus (62). Comparable observations were made in primary cultures of human carcinoid and gastrinoma cells incubated *in vitro* with [^{111}In -DTPA 0]octreotide (63). The primary cultures specifically bound and internalized this radiopharmaceutical. About 50% of the internalized radioactivity was released by the cells within 6 h, whereas the remaining radioactivity was trapped within the cells up to 42 h. Electron microscopic autoradiography demonstrated the presence of the internalized ^{111}In in the cytoplasm and the nucleus. The same processes also apply to the *in vivo* situation. From seven patients with malignant midgut carcinoid tumors, who received an iv injection of 200 MBq [^{111}In -DTPA 0]octreotide 2 d before abdominal surgery, tumor tissue was obtained and analyzed for the subcellular distribution of radioactivity using ultrastructural autoradiography (64). In all patients, the carcinoid tumor could be visualized by preoperative scintigraphy. By *ex vivo* autoradiography, silver grains were found at the plasma membrane, in the cytoplasmic areas among secretory granules and vesicular compartments, but also in the perinuclear area. This localization of ^{111}In in close proximity to the cell nucleus is especially important for this short range Auger electron-emitting radioisotope to exert its cytotoxic effect

in the form of DNA-double strand damage (see *Section II.B* and Ref. 64).

1. *Factors determining the uptake and cellular retention of radioactivity delivered via sst-mediated internalization.* [¹¹¹In-DTPA⁰]octreotide is a sst₂-preferring ligand, which suggests an important role of sst₂ in determining the accumulation of radioactivity in tumor cells after internalization of the radioligand-receptor complex. Radiolabeled octapeptide SS-analogs are internalized in a high amount by sst-positive mouse and human tumor cells (63, 65–67). Evidence for a role of the sst₂ subtype in mediating the uptake of the radiopharmaceutical [¹¹¹In-DTPA⁰]octreotide by sst-positive tumors is presented from studies showing that sst₂-expressing cells internalize SS (54), as well as octreotide (68). Moreover, sst₂ receptor expression correlates well with the relative uptake values of [¹¹¹In-DTPA⁰]octreotide in patients with carcinoid tumor (69), as well as patients with neuroblastoma (70). On the other hand, on the basis of the high SS-internalization rates of the sst₃ and sst₅ subtypes, as reported by Hukovic *et al.* (54), it cannot be excluded that sst₃ and sst₅ might play a role as well. In fact, a role of the sst₃ subtype in the uptake of [¹¹¹In-DTPA⁰]octreotide is evident from a recent study by our group demonstrating a significant uptake in a patient with a thymoma. *In vitro* studies revealed the absence of sst_{2A}, sst_{2B}, and sst₅ receptors and a predominant expression of sst₃ receptors in the tumor tissue (71). The hypothesis that sst subtypes other than sst₂ receptors may be involved in the uptake of [¹¹¹In-DTPA⁰]octreotide *in vivo* is further underlined by the observation that sst scintigraphy visualized tumor sites in three patients with thyroid tumors (one MTC, one Hürthle cell adenoma, and one Hürthle cell carcinoma), which lacked sst₂ mRNA expression but expressed the other four sst subtypes (72), as well as in a patient with a proopiomelanocortin and CRH-expressing MTC (sst₂ negative; sst₁₋₇, sst₃₋₇, and sst₅-mRNA positive; Ref. 73).

As discussed in *Section I.A* (Table 1), octapeptide SS-analogs such as octreotide bind with high affinity to sst₂ and with lower affinity to sst₃ and sst₅ (74, 75). Therefore, both the affinity of the radioligand for the receptor and the efficiency of internalization of the radioligand-receptor complex can be important factors in determining the uptake of radioactivity in sst scintigraphy of sst-positive tumors. Moreover, the differential expression of sst subtypes in tumors (Table 2), as well as the level of sst subtype expression, may play a role. Until now, data on the differential internalization of SS by sst subtypes were derived from studies using cell lines transfected to overexpress the individual sst subtypes (*Section I.C*). Data on the internalization of SS ligands by (tumor) cells endogenously expressing sst subtypes are needed to evaluate the real significance of these findings for human sst-positive tumors. Because most human tumors express multiple sst subtypes, the development of novel sst subtype-selective agonists and antagonists will also be of help to unravel this question. Receptor-mediated endocytosis of SS-analogs is especially important when radiotherapy of human sst-positive tumors using radiolabeled SS-analogs is considered. Internalization of radioligand will result in a prolonged cellular retention of radioactivity, thus resulting in a prolonged exposure of the tumor cell to radiation. Human neu-

roendocrine tumor cells internalize the radioligand [¹¹¹In-DTPA⁰]octreotide. However, this radiopharmaceutical may not be the most suitable compound to carry out radiotherapy because the Auger electrons emitter ¹¹¹In has a low tissue penetration. In addition, a stable coupling of α - or β -emitting isotopes to [DTPA⁰]octreotide could not be achieved, which initiated the development of a novel compound, [DOTA⁰,Tyr³]octreotide, in which the diethylenetriamine pentaacetic acid (DTPA) molecule is replaced by another chelator, tetraazacyclododecane tetraacetic acid (DOTA), allowing a stable binding with the β -emitter yttrium-90 (⁹⁰Y) (76). Recently, we demonstrated that iodinated [DOTA⁰,Tyr³]octreotide is internalized in a high amount by mouse AtT20 pituitary tumor cells, as well as by human insulinoma cells (77). Internalization of iodinated [DOTA⁰,Tyr³]octreotide was approximately 5-fold higher, compared with the iodinated DTPA-coupled parent molecule, [DTPA⁰,Tyr³]octreotide. Therefore, coupling of the octreotide molecule to these chelating groups does not prevent internalization of the hybrid molecules. The high internalization rate of [DOTA⁰,¹²⁵I-Tyr³]octreotide *in vitro* was also evident from the very high uptake of this radioligand *in vivo* in sst-positive organs in rats. De Jong *et al.* (78) recently showed that the amount of [⁹⁰Y-DOTA⁰,Tyr³]octreotide internalized by sst-positive pancreatic tumor cells was significantly higher than that of [¹¹¹In-DOTA⁰,Tyr³]octreotide and [¹¹¹In-DTPA⁰]octreotide (1.8- and 3.5-fold, respectively). Moreover, in eight patients with sst-positive tumors, a higher uptake value of [¹¹¹In-DOTA⁰,Tyr³]octreotide compared with that of [¹¹¹In-DTPA⁰]octreotide was found in normal sst-positive organs like the spleen and pituitary gland, as well as in most tumors (79). If [⁹⁰Y-DOTA⁰,Tyr³]octreotide has the same characteristics of uptake in sst-positive cells, it may be a suitable radiopharmaceutical for sst-targeted radiotherapy.

After internalization of the receptor-radioligand complex, an important process is the retention of radioactivity within the tumor cells. For iodinated SS ligands, it seems clear that a significant proportion of radioactivity is rapidly excreted. This may be due in part to the excretion of radioligand degradation products, although recycling of the receptor-ligand complex may play a role as well. Recycling of SS receptors after being internalized has been demonstrated for sst₂ (80) and sst₃ (44, 81) receptors. Koenig *et al.* (80) also showed that biologically active SS agonists were excreted after being internalized by sst₂-expressing CHO cells. Therefore, trapping of radioisotopes into tumor cells may be an additional important mechanism determining the amount of uptake of radioligand that is used for sst scintigraphy and targeted radiotherapy. In this respect, Duncan *et al.* (82) previously demonstrated that [¹¹¹In-DTPA⁰]octreotide is delivered *in vivo* to pancreatic tumor cell lysosomes and proposed that lysosomes play a critical role in the cellular physiology of radiolabeled SS-analogs. The internalized [¹¹¹In-DTPA⁰]octreotide was shown to be metabolized to ¹¹¹In-DTPA-D-Phe *in vivo* (83, 84). Accumulation of radioactivity in nuclear-lysosomal density gradient fractions was also found in neuroblastoma cells exposed *in vitro* to the radioligand (85). For the radioligand [⁹⁰Y-DOTA⁰,Tyr³]octreotide, it remains to be determined whether similar

processes occur. Finally, uptake of radiolabeled [DTPA⁰]-octreotide in rats and humans (61, 86), as well as that of [DOTA⁰,Tyr³]octreotide in rats (87), demonstrated a bell-shaped curve, dependent upon the amount of injected peptide. Studies to determine the optimal peptide mass for uptake of radioactivity in human tumors after the injection of radiolabeled SS-analogs are ongoing (87).

As shown in Table 1, different octapeptide SS-analogs such as octreotide, lanreotide, and vapreotide (RC-160) interact with the same subclass of sst subtypes (sst₂, sst₃, and sst₅). Nevertheless, slightly different affinities for the different sst subtypes have been found (Table 1). However, *in vivo* studies in rats (88) and humans (89) comparing uptake values of [¹¹¹In-DTPA⁰]RC-160 and [¹¹¹In-DTPA⁰]octreotide showed that [¹¹¹In-DTPA⁰]RC-160 has no additional value for scintigraphy. In fact, the use of [¹¹¹In-DTPA⁰]RC-160 is associated with higher background activity (89). Another recently developed radiopharmaceutical, ¹¹¹In- or ⁹⁰Y-labeled DOTA-lanreotide, bound with high affinity to hsst₂-hsst₅ and with low affinity to hsst₁ expressed in COS-7 cells, suggesting that this radiolabeled peptide may also be useful for sst scintigraphy or radiotherapy (90). Apart from differences in the affinity profiles of unlabeled SS-analogs due to structural differences, radiolabeling of such analogs has major effects on binding affinity for the different human sst subtypes as well (91). Yttrium labeling of [DOTA-Tyr³]octreotide, DOTA-lanreotide, and DOTA-RC-160 significantly increases binding affinities for sst₃ and sst₅ receptors. Such differences, in combination with the high internalization rates of sst₃ and sst₅ receptors (54), could very well account for the higher cellular uptake values *in vivo* and *in vitro* of [⁹⁰Y-DOTA⁰,Tyr³]octreotide, compared with [¹¹¹In-DTPA⁰,Tyr³]octreotide and [¹¹¹In-DTPA⁰]octreotide (77–79). Therefore, several characteristics of SS-analogs developed for sst scintigraphy and radiotherapy, such as small structural modifications, chelator substitution, or type of radioisotope, considerably affect binding affinity (91).

Preclinical studies have shown that down-regulation of SS receptors may occur during agonist exposure. On the other hand, agonist-induced up-regulation of sst expression has been demonstrated as well (see Section III.D.1). Agonist-induced regulation of tumoral sst expression may theoretically influence the results of sst scintigraphy and the efficacy of targeted radiotherapy. The few available clinical data on this issue add to the equivocal data regarding up-regulation and/or down-regulation of SS receptors upon exposure to SS or SS-analog treatment. In five patients with metastatic MTC who were studied before and after 3 months of therapy with a high dose of octreotide, tumor/background ratios determined by sst scintigraphy were reduced in 14 of 18 metastases, suggestive of a down-regulation of SS receptors by octreotide therapy (92). Moreover, reduced orbital uptake of octreoscan was observed in 10 patients with thyroid eye disease after 3 months of treatment with lanreotide or octreotide (93). On the other hand, in patients with a somatostatinoma, the tumors can be visualized by sst scintigraphy (94, 95), suggesting that a complete sst down-regulation does not occur in this type of tumor. Finally, Dorr *et al.* (96) reported decreased uptake of octreoscan in the liver, spleen, and kidney during continuous octreotide therapy, whereas

tumor uptake values were increased simultaneously in five patients with metastatic carcinoid disease and decreased in one patient with advanced MTC. In conclusion, homologous down-regulation of sst expression may be (tumor) cell type specific, as was already evident from experimental studies (see Section III.D.1).

In conclusion, it is well established now that radiolabeled SS-analogs, including those that are used for sst scintigraphy and sst-targeted radiotherapy, can be internalized by sst-positive tumor cells. Several mechanisms may determine the amount of uptake of radiolabeled SS-analogs. These include stability of the radioligand, the expression levels of individual sst subtypes, the affinity of the radioligand for the sst (subtype), the efficiency of receptor internalization and recycling that may be different between sst subtypes, the final trapping of the radioisotopes within the tumor cells, as well as the mass of the injected peptide (summarized in Table 3).

B. SS receptor-targeted radiotherapy

1. Preclinical evidence. Several preclinical studies have demonstrated tumor growth-inhibitory effects after treatment with radiolabeled SS-analogs. In athymic mice bearing sst-positive PC-3 prostatic adenocarcinoma, Zamora *et al.* (97) showed that intratumoral injections with seven 200- μ Ci doses of the β -emitting ¹⁸⁸Re-RC-160 SS-analog reduced tumor size by 90%. Additionally, a significantly higher proportion of survivors was observed in the group treated with ¹⁸⁸Re-RC-160. In this study, treatments were initiated 19 d after inoculation of PC-3 tumor cells when the animals carried solid tumors (500–1000 mm³). In addition, three serial treatments with regionally injected 200 μ Ci ¹⁸⁸Re-RC-160 decreased tumor burdens in experimental models of H-69 human small cell lung cancer cells or ZR-75-1 mammary adenocarcinoma cells xenografted into the pleural cavity of athymic mice (98). Tumor ablation was observed in up to 60% of the animals bearing H69 tumors and in 40% of the animals bearing ZR-75-1 tumors (98). In another study in nude mice bearing solid sst-positive AR42J pancreatic tumors, a single treatment with 500 μ Ci of another SS-analog, radiolabeled with the β -emitting isotope ⁹⁰Y, *e.g.*, [⁹⁰Y]SDZ413, decreased tumor mass by 75% 8 d after injection and prolonged survival, although tumor regrowth was observed after 2 wk of treatment (99). More recently, a significant radiotherapeutic effect of the ⁹⁰Y-labeled SS-analog [⁹⁰Y-DOTA-D-Phe¹,Tyr³]octreotide was demonstrated by the same group of investigators in rats bearing solid sst-positive pancreatic CA 20948 tumors (76). A single iv administration of 10 mCi/kg [⁹⁰Y-DOTA-D-Phe¹,Tyr³]octreotide resulted in a complete remission of

TABLE 3. Factors important in determining the amount of tumoral uptake of radiolabeled SS-analogs

1. Stability of the radioligand
2. Density of SS receptors expressed on tumors
3. Type of SS receptors expressed by tumors
4. Affinity of radioligand for sst subtype(s) expressed by tumors
5. Efficiency of receptor subtype-mediated internalization of radioligand
6. Trapping of radioisotopes within the tumor cell
7. Mass of the injected radiopharmaceutical

the tumors in five of seven (71%) tumor-bearing rats. In this study, no tumor regrowth had occurred even 8 months post injection. This pancreatic CA 20948 tumor expressed a high level of *sst*₂ mRNA and a low level of *sst*₅ mRNA, suggesting that the radiotherapeutic effect of [⁹⁰Y-DOTA-D-Phe¹,Tyr³]octreotide was mediated via targeting the *sst*₂ receptor subtype. [⁹⁰Y-DOTA-D-Phe¹,Tyr³]octreotide is the radiopharmaceutical that is currently being tested in ongoing clinical phase I and II trials (see *Section II.B.2*).

Whereas solid experimental tumor models were used in the above studies, Slooter *et al.* (100) recently studied the radiotherapeutic effect of [¹¹¹In-DTPA⁰]octreotide in a rat model of hepatic metastasis of different tumor cell lines. In this study, the development of hepatic metastases was determined 21 d after injection of *sst*-positive or *sst*-negative tumor cells into the vena porta in rats. Tumor-bearing rats were treated twice (d 1 and d 8) with 370 MBq of [¹¹¹In-DTPA⁰]octreotide. These investigators demonstrated a significant reduction in the number of liver metastases by this treatment regimen in the *sst*-positive tumor model, but not in the *sst*-negative tumors. This suggests that the presence of *sst* on the tumor cells is required for effectiveness of treatment with radiolabeled SS-analogs. This was further confirmed by their observation that the radiotherapeutic effect of [¹¹¹In-DTPA⁰]octreotide could be blocked by coinjection with a *sst*-saturating dose of unlabeled octreotide (100). As indicated above, it should be mentioned that ¹¹¹In is not a β -emitting radioisotope, but emits γ -rays, internal conversion, and Auger electrons. Internal conversion and Auger electrons have a medium- to short-range tissue penetration (200–500 μ m and 0.02–10 μ m, respectively), and it is suggested that these radiochemical properties of ¹¹¹In cause the radiotherapeutic effect of [¹¹¹In-DTPA⁰]octreotide. Because in this study the tumor load was relatively small, studies of the radiotherapeutic effect of [¹¹¹In-DTPA⁰]octreotide in experimental models of more advanced stages of tumor development are required (100).

Recently, a novel ¹⁷⁷lutetium (¹⁷⁷Lu, a low-energy β -particle and γ -emitter) radiolabeled SS-analog, *i.e.*, [¹⁷⁷Lu-DOTA,Tyr³]octreotate, has been proven to be a very promising radiopharmaceutical. Preclinical studies in rats bearing CA 20948 pancreatic tumors demonstrated even a 100% cure of small (≤ 1 cm²) tumors after two repeated doses of 277.5 MBq or one single dose of 555 MBq of [¹⁷⁷Lu-DOTA,Tyr³]octreotate (101). In rats with larger tumors (≥ 1 cm²; range, 1.4–10 cm²), cure rates between 40% and 50% were observed (102). In rats bearing AR42J pancreatic tumors, which had a more favorable uptake compared with the CA 20948 model, treatment with 555 MBq of [¹⁷⁷Lu-DOTA,Tyr³]octreotate resulted in an almost 100% cure, irrespective of the tumor size (102). On the basis of the distinct properties of the two radionuclides and the results of preclinical studies, this group of investigators proposed the use of a combination of ⁹⁰Y- and ¹⁷⁷Lu-labeled SS-analogs, because ⁹⁰Y is particularly effective in large tumors and ¹⁷⁷Lu seems most effective in small tumors (102). Preliminary results in a rat model with tumors of more than one size indeed showed longer survival rates with the combined treatment, compared with treatment with the ⁹⁰Y- or ¹⁷⁷Lu-labeled SS-analogs alone (103).

It is well known that tumor cells display various degrees

of sensitivity to radiation. Adenovirus-based transfer of wild-type (wt) p53 tumor suppressor gene sensitizes ovarian tumor cells to radiation-induced apoptosis (104). In addition, overexpression of the tumor suppressor gene Bax can sensitize tumor cells to radiation, as well as to chemotherapy-induced apoptosis (105, 106). Of particular importance in this respect are recent studies demonstrating that octreotide induces wt p53 and Bax in MCF-7 human breast cancer cells (107). SS-mediated induction of wt p53 and apoptosis is selectively induced via *sst*₃ (108), in contrast to p53-independent, retinoblastoma protein-mediated signaling of cell cycle arrest (109, 110). The majority of human *sst*-positive tumors express *sst*₃ (*Section I*). It cannot be excluded, therefore, that the therapeutic potential of internalized radionuclide may be limited by the lack of expression of a functional p53 or *sst*₃. On the basis of their observations, Sharma and Srikant (107) suggested that α - or β -emitting octreotide-tagged radionuclides should elicit maximal cytotoxic response due not only to radiation-induced damage after internalization, but also to the triggering of apoptosis via the induction of wt p53 and Bax by receptor-mediated signaling. Moreover, on the basis of these data, it is predicted that treatment with SS-analogs alone or in combination with radiation and/or chemotherapy will be most effective in treating wt p53- and *sst*-expressing tumors not only of the breast but also of other organs (107).

Taking the preclinical studies together, it can be concluded that radiotherapy using radiolabeled SS-analogs is effective in experimental *sst*-positive tumor models and that *sst*-targeted radionuclide therapy may be a feasible approach to treat patients with advanced, metastatic *sst*-positive neuroendocrine tumors.

2. Clinical evidence. In 1989, the technique of *sst* scintigraphy to visualize *sst*-positive tumors in man was developed using the radiolabeled SS-analog [¹²³I-Tyr³]octreotide (12). Because the use of this radiopharmaceutical had a number of drawbacks (*i.e.*, expensive, lack of availability, short physical half-life, and predominant hepatic clearance resulting in accumulation of radioactivity in liver, gall bladder, bile ducts, and gastrointestinal tract; Ref. 2), novel SS-analogs were developed to circumvent these disadvantages. As described above, the most widely used SS-analog for *sst* scintigraphy is a DTPA-coupled octreotide. The radioisotope ¹¹¹In binds with very high affinity to the DTPA molecule, and [¹¹¹In-DTPA⁰]octreotide has proved to be a highly suitable radiopharmaceutical for the detection of *sst*-positive tumors by γ -camera scintigraphy (13). Apart from its use in *sst* scintigraphy, [¹¹¹In-DTPA⁰]octreotide has been used for radiotherapeutic application as well (111, 112). Although no controlled trials have been performed with [¹¹¹In-DTPA⁰]octreotide, preliminary reports demonstrate a certain degree of efficacy of this radiopharmaceutical in the treatment of selected, high *sst*-expressing metastasized neuroendocrine tumors. Krenning *et al.* (111) reported treatment with [¹¹¹In-DTPA⁰]octreotide, up to maximal cumulative patient doses of 74 GBq in a phase I trial of 30 end-stage patients with neuroendocrine tumors that all demonstrated recent progression. They reported promising beneficial effects on clinical symptoms, hormone pro-

duction, and tumor size. In 21 patients who received a cumulative dose of more than 20 GBq, 8 patients showed stabilization of disease, and 6 others demonstrated a reduction in tumor size. In a few patients, a transient decline in platelet counts and lymphocyte subsets occurred (111). More recently, this group of investigators reported on 40 evaluable patients treated with doses of at least 20 GBq up to 160 GBq. In 21 patients, therapeutic effects were observed: partial remission in 1, minor remissions in 6, and stabilization of previously progressive disease in 14 patients. On the basis of the observation that three of six patients, who received more than 100 GBq of [$^{111}\text{In-DTPA}^0$]octreotide, developed myelodysplastic syndrome or leukemia, 100 GBq was considered to be the maximal tolerable dose (113). In another study, 14 patients with sst-positive malignancies of different types were treated with two monthly doses of 180-mCi iv injections of [$^{111}\text{In-DTPA}^0$]octreotide. Clinical benefit occurred in 6 of 10 patients with gastroenteropancreatic tumors, objective partial radiographic responses occurred in 2 of 14 patients, and significant tumor necrosis occurred in 6 of 10 patients. Possible treatment-related toxicity included myelosuppression (114). In a patient with a midgut carcinoid tumor, treatment with therapeutic doses of [$^{111}\text{In-DTPA}^0$]octreotide induced symptomatic relief, including a slight reduction in levels of the main tumor marker, urinary 5-hydroxyindolacetic acid (5-HIAA). Again, a slight reduction in leukocyte counts was observed as adverse reaction (112). These data suggest that radiotherapy with high doses of [$^{111}\text{In-DTPA}^0$]octreotide might be useful as a therapeutic agent in patients with sst-positive malignancies. The therapeutic effect of [$^{111}\text{In-DTPA}^0$]octreotide seems due to the emission of Auger and conversion electrons having a low tissue penetration (see Section II.B.1 and Ref. 111). ^{111}In may therefore not be the most optimal radionuclide for sst-targeted radionuclide therapy. For this, novel DOTA-chelated SS-analogs have been synthesized, which allow a fixed binding of β -emitting radionuclides, such as ^{90}Y . Recently, clinical trials using [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide as well as another DOTA-coupled SS-analog, e.g., [$^{90}\text{Y-DOTA}$]lanreotide, have been initiated. Preliminary results with [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide demonstrated promising effects in patients with different types of sst-positive neuroendocrine tumors (115). Multiple treatments with [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide resulted in stable disease in three of six patients and partial remission in the remaining patients. In two of four patients who received a single treatment with [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide, tumor glucose uptake was reduced, whereas the other two showed clinical improvement and stable disease. In a larger group of patients with advanced sst-positive tumors of different origin, a [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide inpatient dose escalation study has been performed. In this study, 29 patients received 4 or more single doses of [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide with ascending activity at intervals of approximately 6 wk. Preliminary results showed disease stabilization in 20 of the 29 patients, partial remission in 2, a reduction in tumor mass of less than 50% in 4, and progression of tumor growth in 3 patients (116). However, a significant proportion of the patients (5 of 29, or 17%) developed renal and/or hematological toxicity. Studies directed to reduce renal toxicity, i.e., amino acid infusions, are ongoing (116). Paganelli *et al.* (117, 118) reported favorable

preliminary results regarding tumor growth using [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide as well. A recent phase II study, including 41 patients with neuroendocrine, gastroenteropancreatic, and bronchial tumors and 82% of patients with therapy-resistant and progressive disease, showed an overall response rate of 24%. Side effects included grade III (NCICG) pancytopenia in 5% and vomiting shortly after injection in 23%. No grade III-IV renal toxicity was observed (119). To evaluate the clinical benefit and objective response rate of high-dose [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide treatment (4 equal iv injections totaling 7.4 GBq/m² with renal protection), a phase II study in 39 patients with progressive neuroendocrine, gastropancreatic, and bronchial tumors was performed (120). The results showed an overall objective response rate of 23% (World Health Organization criteria: complete remission in 5%, partial remission in 18%, stable disease in 69%, progressive disease in 8%). In the patients with endocrine pancreatic tumors, objective response rate was 38% (13 patients). The overall clinical benefit in this study was 63%. Side effects were grade III or IV lymphocytopenia (23%), grade III anemia (3%), and grade II renal insufficiency (3%). Finally, in a patient with metastatic gastrinoma treated with [$^{90}\text{Y-DOTA}$]lanreotide, a 25% reduction in liver metastases as indicated by computed tomography was observed (121). In conclusion, the results of sst-targeted radiotherapy with [$^{90}\text{Y-DOTA}^0, \text{Tyr}^3$]octreotide are most encouraging and extend the therapeutic options in the treatment of patients with sst-positive neuroendocrine tumors.

Very recently, Kwekkeboom *et al.* (122) introduced a novel DOTA-tagged SS-analog, e.g., [$^{177}\text{Lu-DOTA}^0, \text{Tyr}^3$]octreotate (in which the C-terminal threoninol is replaced with threonine), radiolabeled with the β - and γ -emitting ^{177}Lu , as a potential promising radiotherapeutic with a 3- to 4-fold higher tumoral uptake of radioactivity compared with [$^{111}\text{In-DTPA}^0$]octreotide. [$^{177}\text{Lu-DOTA}^0, \text{Tyr}^3$]octreotate has a 9-fold increased affinity for sst₂, compared with [$^{177}\text{Lu-DOTA}^0, \text{Tyr}^3$]octreotide, and a 6- to 7-fold increase in affinity for their yttrium-loaded counterparts (91). Preliminary promising results using this radiopharmaceutical have been reported (102, 103).

C. SS receptor-targeted chemotherapy: preclinical evidence

The wide spectrum of adverse reactions when treating patients with advanced, metastatic tumors with chemotherapeutic agents are caused by the severe toxicity of these agents to normal cells. Like peptide receptor-targeted radiotherapy, targeted chemotherapy to deliver the chemotherapeutic compounds selectively to tumor cells might be a promising approach as well. Schally and Nagy (123) pioneered this concept with the development of cytotoxic analogs of LHRH, cytotoxic bombesin analogs, and cytotoxic SS-analogs, to treat LHRH receptor-, bombesin receptor-, and sst-positive tumors, respectively. This group of investigators provided preclinical evidence for the effectiveness of cytotoxic LHRH analogs in experimental models of human ovarian, mammary, or prostatic cancer (123), as well as for the effectiveness of cytotoxic bombesin analogs in the treatment of experimental models of bombesin receptor-positive small cell lung carcinoma, colorectal, gastric, pancreatic, mammary, and prostatic cancers (123). Schally and co-work-

ers (124, 125) synthesized two different cytotoxic SS-analogs, code-named AN-51, consisting of methotrexate linked to the N terminal of the octapeptide SS-analog RC-121 and AN-238, which is the RC-121 analog linked to a highly potent derivative of doxorubicin, *e.g.*, 2-pyrrolinodoxorubicin. Both the AN-51 and AN-238 compounds had intermediate binding affinities to sst-positive tissues *in vitro*, in comparison with the RC-121 compound alone, suggesting that coupling of the chemotherapeutic compound to the SS-analog slightly reduced their binding properties (124, 125). The binding affinity of the AN-238 compound for rat pituitary membrane SS receptors was 23.8 nM, which is comparable to the binding affinities of several DTPA- and DOTA-coupled SS-analogs to hsst₂ receptors (91). In preclinical studies, it was demonstrated that both the AN-51 and AN-238 compounds inhibited tumor growth in experimental tumor models. In nude mice transplanted with the human Mia PaCa-2 pancreatic tumor, AN-51 significantly inhibited tumor growth, whereas the chemotherapeutic compound alone, methotrexate, or RC-121 alone had no significant inhibitory effect (124), and with methotrexate alone displaying a much higher toxicity compared with AN-51. Thereafter, this group tested the cytotoxic properties of the AN-238 compound, which displayed a very high toxicity in sst-positive cells *in vitro*. Potent tumor growth inhibitory properties of AN-238 were observed in many experimental mouse and rat models of human breast cancer, prostate cancer, ovarian cancer, small cell lung cancer, pancreatic cancer, renal cell cancer, as well as glioblastoma (126–133). Again, a much higher toxicity and lower or absent effectiveness on tumor growth was observed in animals treated with the cytotoxic radical alone. In these animal studies, the major side effect of treatment with cytotoxic SS-analogs was a transient fall in white blood cell counts. In conclusion, sst-targeted chemotherapy is effective in preclinical tumor models and seems a highly promising approach as well to treat sst-positive tumors. The sst-targeted chemotherapy may result in a chemotherapeutic approach using lower dosages of the chemotherapeutic compound and thus lower toxicity. Until now, however, no clinical trials have been reported using targeted LHRH, bombesin, or SS-analogs. In addition, evidence will have to be provided that cytotoxic SS-analogs can also be internalized by sst-positive tumor cells. As for the concept of sst-targeted radiotherapy, the efficacy of sst-targeted chemotherapy will be determined by the amount of uptake of the cytotoxic radicals by the tumors. Moreover, the effect of cytotoxic SS-analog treatment on the function of normal sst-expressing cells is to be determined.

III. Tachyphylaxis and Resistance to SS

A. Introduction

Concomitant with the widespread distribution of sst throughout central and peripheral tissues, the acute administration of SS or its analogs induces a large number of mainly inhibitory effects (8, 9). Nevertheless, these initially potent responses diminish with continued exposure (9, 11). The different mechanisms that are potentially involved in this adaptation or tachyphylaxis to continuous exposure to SS or

SS-analogs may be associated with processes such as receptor phosphorylation, G protein uncoupling, receptor internalization, and degradation. This has been reviewed extensively (8, 9). Different from these physiological responses to continued SS exposure is the response of neuroendocrine tumor cells. Patients with certain types of sst-positive tumors (*e.g.*, GH-secreting pituitary adenomas, islet cell tumors, and carcinoids) can be treated for many months to years with the current clinically available SS-analogs. The long-term control of hormonal hypersecretion and/or tumor growth by treatment with SS-analogs may vary considerably, however, among patients. *Section III* focuses particularly on tachyphylaxis and resistance to treatment with the different available formulations of SS-analogs, as well as the potential mechanisms involved herein. Although the direct fundamental evidence for the clinical observations of tachyphylaxis is relatively weak, several potential mechanisms determining cellular responsiveness to SS are discussed.

B. Tachyphylaxis of pathological hormone secretion

1. *Pituitary adenomas.* Whereas normal hormone secretion shows tachyphylaxis after continuous receptor activation within hours to days (9), pathological hormone secretion by sst-positive tumor cells can be inhibited during significantly prolonged periods. In about half of the patients with GH-secreting pituitary adenomas, serum GH and IGF-I levels are normalized by octreotide treatment (134). Escape from SS-analog therapy has not been observed in this type of patient, even after many years of continuous treatment (135). Figure 2A shows a typical example of the persistent suppression of serum IGF-I levels in an acromegalic patient during a period of 8 yr of treatment with three times daily injections of 50–100 µg octreotide. The persistently lowered IGF-I levels seem not to be caused by radiotherapeutic effects because drug withdrawal after 5 yr of treatment resulted in an instant rise in IGF-I levels and immediate recurrence of signs and symptoms. Moreover, although no desensitization to continuous sc treatment with octreotide is observed, acromegalic patients treated with long-acting formulations of SS-analogs, *i.e.*, long-acting repeatable octreotide administration (Refs. 135a–135c) or slow-release lanreotide (136) did not show any signs of tachyphylaxis to treatment periods up to 3 yr as well. To our knowledge, only one rare case of acromegaly showing desensitization to octreotide has been described so far (137). Partial tachyphylaxis to SS-analogs was reported in a patient with acromegaly. In this patient, previously treated with ⁹⁰Y implant, external radiotherapy, and three daily sc injections with octreotide, GH levels progressively rose after switching to lanreotide and depot octreotide (Sandostatim LAR, Novartis Pharmaceuticals Corp., Basel, Switzerland). Interestingly, there were no signs of tumor growth or alterations in sst status as determined by [¹¹¹In-DTPA⁰]octreotide scintigraphy (138). Octreotide withdrawal for 24 h in this patient resulted in a 64% increased sensitivity in terms of inhibition of GH levels by recommencing octreotide treatment, suggesting that changes in receptor function or on the receptor signal transduction cascade play a role, rather than changes in receptor expression (138).

The majority of TSH-secreting and clinically nonfunction-




FIG. 2. Absence of tachyphylaxis to octreotide therapy in a patient with a GH-secreting pituitary adenoma (A) and desensitization in a patient with metastatic carcinoid tumor (B). A, Effect of octreotide therapy on serum IGF-I level in a patient with a GH-secreting pituitary adenoma. A 77-yr-old man transsphenoidally operated (TSS) for a GH-secreting macroadenoma that had resulted in active acromegaly. Two months after incomplete surgery of the tumor, external radiotherapy (RT) was applied. Successively, octreotide therapy was started 6 months after surgery at a dose of 100 μ g three times daily. This therapy had resulted in a prolonged suppression of serum IGF-I levels ($N < 43$ nmol/liter) and disappearance of signs and symptoms of active acromegaly. Octreotide therapy has been continued for more than 8 yr. The dose could be reduced to 50 μ g three times daily. Discontinuation of therapy resulted in an increase of serum IGF-I levels and immediate recurrence of signs and symptoms of active acromegaly. Dotted line shows the upper normal limit of serum IGF-I levels. B, Effect of octreotide therapy on urinary 5-HIAA levels in a patient with a metastatic carcinoid tumor. A 66-yr-old man, operated for a metastatic carcinoid tumor of the small intestine with abdominal lymph node metastases, hepatic metastases, and the malignant carcinoid syndrome. Urinary 5-HIAA levels were greatly elevated ($N < 40$ μ mol/24 h). Therapy with octreotide was started at a dose of 100 μ g three times daily. This therapy initially resulted in a reduction of attacks of flushing and improvement of diarrhea, which was accompanied by more than 50% reduction (but not normalization) of urinary 5-HIAA levels. However, after 4–6 months of therapy, the patient developed resistance to therapy: the flushing attacks, frequency of diarrhea, and urinary 5-HIAA levels gradually increased despite increasing the dose to 500 μ g three times daily. In addition, a slight increase of tumor mass was observed. The dotted line represents the upper normal limit of urinary 5-HIAA levels.

ing pituitary adenomas also express sst (139, 140). Octreotide treatment of patients with TSH-secreting pituitary adenomas results in a lowering of TSH levels and normalization of T_4 levels in 73% of the patients. In contrast to patients with GH-secreting pituitary adenomas, in this series of 52 patients an escape from therapy was observed in 5 patients (10%). This loss of sensitivity to the inhibitory effect of octreotide on TSH levels was observed in two patients receiving short-term therapy and in three patients receiving long-term therapy (139). Overall, Beck-Peccoz *et al.* (141) reported tachyphylaxis in 22% of the patients with a response to increasing octreotide doses, whereas subsequent escape from the inhibitory effects was observed in 10% of the cases. The role of SS-analogs in the treatment of patients with clinically nonfunctioning pituitary adenomas is less well established, whereas octreotide seems not of benefit in the treatment of patients with ACTH-secreting pituitary adenomas or prolactinomas (135). This may be due to either the absence of sst on the tumor cells or the absence of expression of particular sst subtypes (135).

2. *Islet cell tumors and carcinoids.* In striking contrast to the absence of the occurrence of tachyphylaxis of inhibition of hormone secretion by octreotide in patients with GH-secreting pituitary adenomas are the observations in patients with

islet-cell tumors and carcinoids. In the majority of patients with metastatic carcinoids, VIPomas, gastrinomas, insulinomas, and glucagonomas, treatment with octreotide induces a rapid improvement of clinical symptomatology, such as diarrhea, dehydration, flushing attacks, hypokalemia, peptic ulceration, hypoglycemic attacks, and necrotic skin lesions (142–145). On the other hand, the majority of the patients show desensitization of the inhibition of the secretion of tumor-related hormones by octreotide within weeks to months. This effect may be initially reversed by increasing the dosage of octreotide, but eventually the drug becomes ineffective in all patients (11). In a series of 57 patients with the carcinoid syndrome, 23 patients escaped from octreotide therapy after periods ranging from 1 wk to 12.5 months (median, 4 months), whereas the other responding patients could be controlled for periods extending to 2.5 yr. The estimated mean duration of response to octreotide therapy in the whole group of responding patients was approximately 1 yr (146). Figure 2B shows a typical example of tachyphylaxis of the inhibitory effect of octreotide (100 μ g three times per day) on urinary 5-HIAA levels in a patient with a metastatic carcinoid tumor. An escape from octreotide treatment was seen after 3 months of therapy. Increasing the dose of

octreotide (to 500 μg three times per day) was not beneficial in this particular patient. The potential mechanisms responsible for this desensitization, as well as for the considerable variability in the duration of the responses to octreotide therapy, are not known at present. The relatively long time-frame of this escape suggests mechanisms other than G protein uncoupling or internalization are involved. It has been suggested that this loss of sensitivity of endocrine cancers to octreotide is possibly associated with the outgrowth of clones of tumor cells that lack sst rather than with a transient down-regulation of these receptors (147). Moreover, it is not known why pituitary GH-secreting pituitary adenomas do not show tachyphylaxis to octreotide or lanreotide treatment, whereas the majority of patients with metastatic carcinoids, VIPomas, gastrinomas, insulinomas, and glucagonomas eventually desensitize. Possibly, SS-analog treatment of patients with GH-secreting pituitary adenomas induces an up-regulation of sst in the GH-secreting tumor cells, whereas other types of tumors display down-regulation of sst upon prolonged agonist exposure. As will be discussed in *Section III.D.1*, up-regulation and/or down-regulation of sst expression may be sst subtype dependent. As has been discussed in *Section I*, the majority of human sst-positive tumors express multiple sst subtypes, often with overlapping patterns. Therefore, potential tissue-specific desensitization and/or down-regulation of sst subtypes or, alternatively, tissue-specific up-regulation of the octreotide-responsive sst subtypes 2, 3, and 5 induced by prolonged agonist treatment may account for continued responsiveness of GH-secreting pituitary adenomas to SS agonists.

C. Escape from antiproliferative effects

Apart from regulating neurotransmission and secretion, SS and its analogs may inhibit cell proliferation in normal and tumoral tissues as well. Evidence for inhibition of tumor cell proliferation by SS-analogs is based primarily on studies using experimental sst-positive tumor models (2, 3). However, in a number of these studies, tumor growth is only delayed, because after a certain treatment period the tumors start to grow more rapidly, resulting in growth curves that parallel tumor growth in untreated control animals, indicating escape from SS-analog therapy (148–150). In the model of the transplantable prolactin (PRL)-secreting pituitary tumor (149), we observed during the first 2 wk of treatment with the SS-analog octreotide a significant reduction in tumor growth. After 2 wk of treatment, however, tumor growth rates in untreated and octreotide-treated animals were parallel and not significantly different. One of the mechanisms underlying this tachyphylaxis may be a down-regulation of SS receptors on the tumor cells. In primary cultures of PRL-secreting 7315b cells, an incubation with octreotide for 1 wk inhibited both the growth and hormone secretion in a parallel and dose-dependent fashion. However, prolonged (5 wk) continuous exposure to octreotide (0.1 nM to 1 μM) resulted in tachyphylaxis with respect to the inhibition of PRL secretion. In a stable cell line derived from this 7315b tumor, long-term exposure to octreotide induced a loss of sensitivity with respect to both PRL secretion and cell growth. This loss of sensitivity was accompanied by a

complete down-regulation of SS binding sites on the tumor cells. In this 7315b sst-expressing tumor model, clonal selection of sst-negative cells was not the cause of desensitization, because withdrawal of treatment from desensitized cells resulted in a reappearance of sst and the sensitivity to octreotide (151). A significant reduction in sst numbers induced by octreotide treatment has also been demonstrated in Syrian hamsters bearing transplanted insulinomas. Twice-daily injections with octreotide for 3 d resulted in a dose-dependent reduction in sst numbers on the insulinomas (2). On the other hand, the occurrence of tachyphylaxis to treatment with SS-analogs can be tumor cell type specific. *In vivo* studies by other groups showed an increase in SS-binding on experimental tumors treated with SS-analogs. Treatment of human MKN45 gastric carcinoma xenografts in nude mice for 5 wk with the SS-analog RC-160 significantly inhibited tumor growth without the occurrence of an escape. Daily sc injections with RC-160 even induced a significant up-regulation of sst in these tumors after 4–5 wk, which in this particular tumor model may be beneficial in maintaining the inhibitory effects on tumor growth (152). A comparable up-regulation of sst binding sites has been demonstrated in AR4-2J pancreatic tumor-bearing mice, in which continuous treatment (7 d) with a low dose of octreotide, administered via octreotide-containing osmotic minipumps, induced an increase in the number of tumoral sst binding sites (153). In contrast to this up-regulation of sst binding sites on pancreatic AR4-2J tumors by continuous *in vivo* treatment with low doses of octreotide, discontinuous (twice daily) sc injections of octreotide resulted in a down-regulation of sst expression (153). After removal of octreotide *in vitro*, a total recovery of [^{125}I -Tyr 3]octreotide binding was observed within 24 h. This recovery was dependent on protein synthesis, making *de novo* receptor synthesis necessary for the recovery process (153). RT-PCR analysis revealed that AR4-2J cells expressed sst $_2$ receptor mRNA only. In fact, these authors concluded that continuous treatment with a low dose of octreotide might improve the efficacy of long-term octreotide therapy. These data suggest that in a single tumor model the experimental conditions may determine whether sst $_2$ receptors are either down-regulated or up-regulated. In conclusion, the escape from the tumor growth-inhibitory effects of SS-analogs suggests that prolonged exposure to agonists may be due to sst down-regulation. Moreover, in some tumor models an up-regulation of sst expression after agonist exposure has been observed, which might explain prolonged responsiveness to SS-analogs. These apparently opposite experimental results preclude making generalized conclusions with respect to the optimal SS-analog treatment modalities that might apply to patients with sst-positive neuroendocrine tumors. In addition, an escape from SS-analog treatment could, alternatively, involve an up-regulation of binding sites that do not recognize octreotide and/or an escape of tumor cells that do not express octreotide-responsive sst subtypes. In the majority of the above-mentioned studies, the precise mechanisms of changes in sst numbers were not studied in detail. Therefore, it remains to be established whether the changes in sst numbers at the cell surface are mediated via reduced sst gene transcription, decreased stability of sst mRNAs, via

an increased intracellular breakdown of preexistent cellular SS receptors, or a combination of these events.

D. Mechanisms of tachyphylaxis and resistance

1. *Homologous (down-)regulation of sst expression.* Although uncoupling from G proteins and internalization of SS receptors cannot be fully excluded as a potential cause for reduced sensitivity to long-term SS-analog treatment in patients with neuroendocrine tumors, other mechanisms are more likely to be involved. Down-regulation of cellular SS receptors may form a long-term cause of tachyphylaxis after continuous exposure of SS receptors to agonists. On the one hand, chronic exposure of cultured pituitary cells to relatively high concentrations of SS-14, SS-28, or SS-analogs reduces the number of sst on AtT20 and 7315b pituitary tumor cells (149, 151, 154, 155). On the other hand, an up-regulation of sst expression has been observed in GH₄C₁ or GH₃ rat pituitary tumor cells (156, 157). This up-regulation of SS receptors in GH₄C₁ or GH₃ was related to changes in sst gene expression, rather than changes in receptor affinity. In fact, in GH₃ cells, chronic exposure with SS induces an increase of sst₁, sst₃, sst₄, and sst₅ mRNA expression after 6–48 h of exposure, whereas sst₂ mRNA expression displayed a biphasic response, with an increase at 2 h, a decrease at 6 h, and finally normalization after 48 h (157). Therefore, agonist-induced down-regulation and/or up-regulation of sst expression is time dependent and cell type specific. In cells that do not express sst subtypes endogenously, but were transfected with the different sst subtype genes to overexpress the different sst subtypes, agonist exposure has differential effects, depending on the sst subtype investigated. Short-term (1 h) agonist exposure decreases SS-binding in CHO cells expressing the sst_{2A} receptor (158, 159), whereas prolonged exposure (22 h) to the peptide induces an increased binding (54). SS binding in cells expressing sst₃ and sst₅ receptors was not affected by SS pretreatment, whereas sst₄ and sst₁ receptors were up-regulated (54). Whether these sst subtype-specific responses to agonist exposure also occur in human sst-positive tumors, which express multiple sst subtypes, remains to be established. To our knowledge, no such data are available at present, except for clinical data on responsiveness and the induction of tachyphylaxis to SS-analog therapy (see Section III.B.2). Apart from agonist-induced changes in cell surface sst number, tachyphylaxis of responsiveness after chronic agonist exposure and/or resistance to SS-analog treatment may be induced by several other potential mechanisms as well. Such mechanisms include heterologous regulation of SS cell surface numbers, heterogeneous expression of SS receptors in human tumors, or sst gene mutations, and they are discussed in the following paragraphs.

2. *Heterologous regulation of sst expression.* Apart from homologous regulation of sst expression (Section III.D.1), heterologous up- and down-regulation of SS receptors on normal and tumorous cells has been demonstrated as well. Glucocorticoids down-regulate sst numbers in GH₄C₁ rat pituitary tumor cells. In these cells, both cortisol and dexamethasone reduce the specific binding of [¹²⁵I-Tyr¹]SS-14 by 20% and 40%, respectively (160), probably via the inhibition of *de*

novo protein synthesis. Moreover, sst subtype expression in GH₄C₁ cells is differentially regulated by glucocorticoids. Short-term incubation for 2 h with dexamethasone increases sst₁ and sst₂ mRNA levels, whereas sst₃ mRNA levels were unchanged. On the other hand, prolonged exposure (2 d) with dexamethasone induced a reduction in sst₁ and sst₂ mRNA levels and a dramatic up-regulation of sst₃ mRNA levels. Nuclear run-on assays showed that the changes in sst₁ and sst₂ mRNA levels were associated with changes in sst gene transcription rate (161). Indirect clinical evidence for the *in vivo* down-regulation of tumoral SS receptors by glucocorticoids was obtained from the observation that in five patients with untreated Cushing's disease, octreotide did not inhibit basal or CRH-stimulated ACTH levels and did not influence cortisol levels. *In vitro*, however, octreotide inhibited CRH-stimulated ACTH secretion by human corticotroph adenoma cultures, whereas this inhibitory effect was abolished by hydrocortisone pretreatment (162). Estrogens have been shown to stimulate sst expression in pituitary (tumor) cells (163–165) *in vitro* and *in vivo*. Chronic estrogen treatment up-regulates sst₂ receptor mRNA expression in the anterior pituitary gland *in vivo* (166). Considerably less information is available regarding the heterologous regulation of sst expression in nonpituitary-derived cell systems. In breast cancer cell lines, estrogen stimulates steady state mRNA levels (167). A 5.3-kilobase pairs (kb) 5'-flanking region of the *hsst₂* gene, lacking TATA and CCAAT boxes, is the active promoter in estrogen receptor-positive breast cancer cell lines (168). In agreement with these observations, Kimura *et al.* (169) recently demonstrated that estrogen regulated promoter activity of a 5-kb 5'-untranslated region of the rat sst₂ gene, lacking TATA and CCAAT boxes (169). In concordance with the findings in pituitary-derived cells, dexamethasone may cause down-regulation of sst numbers without changing receptor affinity in AR42J rat pancreatic acinar carcinoma cells (170). Thyroid hormones may regulate sst expression as well. In TtT-97 tumors, which represent an *in vivo* murine thyrotropic model not expressing any sst subtype mRNA or protein, thyroid hormone treatment induces specific up-regulation of sst₁ and sst₅ mRNAs and high affinity sst binding sites in the tumors (171). Taken together, these data demonstrate that sst subtype expression can be influenced by different steroids and hormones in a time-specific and receptor subtype-specific manner. It is not established, however, whether treatment of patients with, for example, glucocorticoids or antiestrogens may influence sst expression and thus responsiveness to SS *in vivo* as well. Apart from a regulatory effect of glucocorticoids and estrogens on sst expression, it is also likely that such agents directly influence the responsiveness of tumor cells to SS agonists. Indeed, breast cancer cells have been shown to respond better to the cytotoxic effect of octreotide in the presence of the antiestrogen tamoxifen (172).

3. Resistance to SS agonists

a. *Heterogeneity of tumoral sst expression.* Certain subgroups of human sst-positive tumors express sst subtypes on the basis of their differential binding of SS and SS-analogs. The sst autoradiographic studies showed the absence of binding of [¹²⁵I-Tyr³]octreotide in a small subgroup of human insu-

linomas, carcinoids, pituitary adenomas, and meningiomas, in 50% of MTCs, and in all sst-positive ovarian cancers, whereas in the same tumors binding sites for iodinated-[Tyr¹¹]SS-14 or [LTT]SS-28 were present (23–25). This differential binding between octreotide on the one hand and SS-14/SS-28 ligands on the other hand, in insulinomas and other subgroups of sst-positive tumors, suggests that resistance to octapeptide SS-analogs may be due to the absence of specific sst subtypes that bind these analogs with high affinity, but also indicates that novel sst subtype-selective analogs can be developed for the treatment of patients with tumors carrying sst of this particular subtype(s). Although certain human sst-positive tumors lack particular sst subtypes with high affinity for octapeptide SS-analogs (Table 1), some tumors have been demonstrated to express a nonhomogenous distribution of SS receptors (23–25). A nonhomogenous distribution of sst has been found in a subset (3 of 10) of human GH-secreting pituitary adenomas (173), as well as in rare cases of carcinoid tumors (23). Moreover, in more than 50% of breast cancer specimens, sst expression displayed a nonhomogenous distribution, *i.e.*, both sst-positive and sst-negative tumor regions within individual sst-positive tumors (174). One rare case of a human carcinoid tumor has been described in which sst₁ and sst₂ mRNA were clearly localized in different tumor regions (28). In such cases, resistance to SS-analog therapy, after an initial response, may be due to the outgrowth of sst (sst₂)-negative tumor cell clones, which in fact may still express sst, albeit of the subtype to which the current generation of octapeptide SS-analogs do not bind.

b. SS receptor gene mutations. To date, relatively few data are available with respect to sst-gene mutations leading to a loss of sst function. One study addressed this issue so far in COR-L103 small cell lung cancer cells (175). Sequence analysis of the sst₂ gene demonstrated a point mutation in codon 188 of TGG for tryptophan to TGA for a stop codon causing a loss of 182 C-terminal amino acid residues in sst₂, resulting in the absence of sst₂ expression in the plasma membrane of COR-L103 cells. The nucleotide sequences of the sst₃ and sst₄ genes, which were also expressed in these cells, were normal. In a series of 19 human GH-secreting pituitary adenomas with variable sensitivity to SS-analog treatment *in vivo*, the sst₂ and sst₃ genes were found to possess intact coding sequences (176). Moreover, no mutations affecting the sst₂ protein were detected in a series of 15 GH-secreting pituitary adenomas (177). These data suggest that mutations in these sst subtypes do not form the basis for resistance of tumoral GH secretion to SS-analogs. Ballare *et al.* (178) recently described a germ line mutation (Arg240Trp) in the sst₅ gene in an acromegalic patient resistant to SS-analog treatment. This mutation results in decreased sensitivity to the inhibitory effect of SS on adenylate cyclase activity, whereas cells expressing the mutant sst₅ displayed increased proliferation and increased MAPK activity, compared with wt cells. These data suggest that this mutation in sst₅ abrogated the anti-proliferative action by SS and activated mitogenic pathways. Nevertheless, such mutations appear to be very rare. Finally, in none of a series of 43 neuroblastoma tumors were mutations in the sst₂ gene detected by PCR-based single-stranded

conformation polymorphism/heteroduplex analysis (179). Mutations in other sst subtypes that may cause this resistance cannot be excluded, however. Moreover, other causes such as sst density and/or the above-discussed mechanisms of resistance (summarized in Table 4) may play a role as well.

c. Miscellaneous potential causes of resistance to SS-analogs. Antibodies to octreotide that develop in patients treated with this analog (138, 180–182) seem not to be an important cause of escape from therapy with SS-analogs, because continued efficacy of octreotide treatment has been documented in two acromegalic patients who had antibodies to octreotide (181). G protein mutations, particularly mutations in G_sα, have been shown to be associated with overproduction of hormones by pituitary-derived hormones, as well as with pituitary hyperplasia (183). In a subgroup of patients with GH-secreting pituitary adenomas, high basal adenylyl cyclase activity and poor responsiveness to stimulatory agents such as GH-releasing hormone suggested constitutive activation of the adenylyl cyclase cascade in the tumor cells. A considerable number of these tumors indeed contained an activating mutation in G_sα (183), which correlates with a higher sensitivity to SS agonists. An increase in sst₂ mRNA does not seem to account for this increased sensitivity (184). However, mutations in inhibitory G proteins are rare, and mutations in G_{i2}α, to which sst₂ is capable of associating (185), have only been described in small numbers of adrenal cortical tumors (27%) and ovarian tumors (30%; Ref. 186). It appears therefore, that resistance to SS-analog therapy due to a mutation in inhibitory G proteins coupled to sst is not very likely to occur.

E. New developments

As described in Section III.D.3, one of the causes for resistance to therapy with the current generation of octapeptide SS-analogs may be the absence or low expression of sst₂ receptors by the tumor cells. The question then arises: What might be the role of other sst subtypes as a target for therapy with novel SS-analogs? Functional evidence for the existence of sst subtypes comes from studies using human fetal pituitary cell cultures in which SS regulates GH and TSH secretion by both sst₂ and sst₅, and PRL secretion mainly by sst₂ (74). In recent years, many new sst selective analogs have been synthesized. Using primary cultures of human GH-secreting pituitary adenomas, Melmed and co-workers (75) demonstrated that combinations of sst₂- and sst₅-selective

TABLE 4. Potential mechanisms of tachyphylaxis and resistance to SS-analog therapy in patients with sst-positive tumors

1. Down-regulation: decrease in the number and/or affinity of SS receptors
2. Desensitization: decrease in responsiveness due to receptor uncoupling from second messenger activation
3. Nonhomogeneous expression of SS receptors in tumors: outgrowth of sst-negative cell clones
4. Resistance due to the absence of sst subtypes with high affinity for octapeptide SS-analogs
5. Resistance due to tachyphylaxis of the inhibitory effect of SS-analogs on indirect tumor growth-promoting mechanisms (*i.e.*, GH or gastrin secretion)
6. Mutations in sst genes leading to absence of functional receptor proteins

compounds decreased GH secretion significantly more than the single compounds alone. In addition, in PRL-secreting primary tumor cell cultures, PRL secretion was preferentially inhibited by sst_5 -selective analogs, whereas sst_2 -selective analogs were ineffective. Even more exciting are recent studies using a bispecific sst analog with high affinity binding to both sst_2 and sst_5 receptors. This compound, BIM-23244 (Table 1), was quite effective in inhibiting GH secretion *in vitro* by a series of five octreotide partially responsive tumors. These tumors turned out to have 9-fold lower sst_2 mRNA levels and approximately 7-fold higher sst_5 mRNA levels, compared with a group of octreotide-sensitive tumors. The same compound also inhibited PRL release by five mixed GH-PRL-secreting pituitary adenomas (187). Thus, apart from highly sst-selective analogs, there may be a place for new sst bispecific analogs in the treatment of pituitary adenomas resistant to sst_2 agonists. More recently, a SS peptidomimetic, named SOM230, with high affinity for sst_1 , sst_2 , sst_3 , and sst_5 receptors (Table 1) has been shown to have a much higher efficacy in lowering normal plasma IGF-I levels in rats, compared with the effects of the sst_2 -selective analog octreotide (188). Long-term (weeks to months) continuous, as well as discontinuous, treatment with octreotide in rats is known to result in a loss of the inhibitory effect of the drug on circulating GH and IGF-I levels (148, 149, 189). The potent inhibitory effect of SOM230 on IGF-I levels, showing no signs of loss of its inhibitory effect during a period of 126 d of continuous infusion, could be explained by a 40-fold increase in the affinity for sst_5 receptors, as compared with octreotide in combination with the key role that sst_5 plays in controlling GH release (75, 187). SOM230 has a very long terminal elimination half-life of 23 h in rats, compared with octreotide (2 h), and no obvious adverse side effects, including changes in glucose levels, over the 126-d period of treatment, and it is currently under evaluation in phase I trials (188). Moreover, sst subtypes may form homo- or heterodimers (56, 58) or may heterodimerize with other G protein-coupled receptors such as the dopamine D2 receptor (60) or the opioid receptor MOR-1 (59), resulting in a novel receptor state with properties distinct from the individual receptors in terms of enhanced internalization, reduced agonist-induced desensitization, and functional activity. These new fundamental insights into receptor function will help us to explain the observed differences in the development of tachyphylaxis not only between patients with different tumor types, but also among patients with the same type of neuroendocrine tumor but with different sst subtype expression patterns. It is a challenge to evaluate whether these new bispecific or more universal SS-analogs are indeed effective in tumors resistant to the current clinically available compounds as octreotide and lanreotide, as well as to investigate whether such new compounds can prevent neuroendocrine tumors from tachyphylaxis to treatment. Apart from new analogs with a broader sst binding profile, a hybrid SS-dopamine molecule has also been recently synthesized. This molecule, BIM-23A387, retained high affinity binding to both sst_2 and D2 receptors and had a tremendous enhanced potency on GH and PRL release by primary cultures of human pituitary adenoma cells, compared with sst_2 - and D2-specific analogs, alone or in combination (190). This significant enhanced po-

tency, however, could not be explained on the basis of the binding affinity of the compounds for sst_2 and D2 receptors (190). The mechanism by which this molecule exerts its potent action is unknown but strengthens the observations that processes like heterodimerization of receptors indeed have functional implications.

F. Conclusions

The induction of tachyphylaxis of responsiveness to SS-agonists has been demonstrated in a variety of sst-positive cell systems. The time-frame of the occurrence of tachyphylaxis *in vivo* on normal hormone secretion is relatively rapid (hours to days), whereas escape from therapy with SS-analogs in patients with sst-positive tumors or in experimental models of sst-positive tumors generally occurs after prolonged exposure to SS agonists (weeks to years). This relative late induction of tachyphylaxis of responsiveness suggests that sst down-regulation, rather than rapid processes like G protein uncoupling and/or receptor internalization are involved. Moreover, escape from SS-analog therapy could involve the outgrowth of tumor cell clones lacking the expression of sst subtypes to which the currently clinically used octapeptide analogs bind with high affinity. The development of novel sst subtype-selective and nonselective analogs, as well as chimeric compounds, could be of interest as potential new treatment modalities for resistant tumors. The only group of tumors that show no signs of desensitization to treatment with SS-analogs are GH-secreting pituitary adenomas. In SS-analog-sensitive patients with GH-secreting pituitary adenomas, circulating GH and IGF-I concentrations can be effectively suppressed, even during many years of treatment with these compounds. The underlying mechanisms for this difference in developing tachyphylaxis to SS-analog treatment between GH-secreting pituitary adenomas on the one hand, and other types of neuroendocrine tumors on the other hand, have not yet been elucidated but could involve the differential expression of sst subtypes, a tissue-specific desensitization, and/or down-regulation of sst subtypes, or alternatively, tissue-specific up-regulation of SS-analog responsive sst subtypes by prolonged agonist treatment resulting in continued responsiveness.

In conclusion, clinical observations clearly demonstrate tachyphylaxis and/or resistance to SS-analog treatment in patients with neuroendocrine tumors, but the direct fundamental evidence explaining the mechanisms involved is currently weak.

IV. Summary

During the past decade, novel insights into the physiological and pathophysiological role of SS and its receptors have been developed. Although in the mid-1980s it was debated whether or not SS was internalized by sst-expressing cells, recent studies have now clearly demonstrated that SS and SS-analogs are efficiently internalized via a rapid process of agonist-induced receptor-mediated endocytosis. Moreover, in 1989 the technique of sst scintigraphy to visualize sst-positive tumors in humans was developed, and the concept of the radiotherapeutic use of radioisotope-coupled SS-

analogs, *i.e.*, peptide receptor radionuclide therapy, was introduced. Finally, in the beginning of the 1990s, five sst subtypes were cloned and characterized, and the expression of these subtypes has been studied in both normal and tumoral sst-expressing tissues. Taking these discoveries together, several new questions can be raised. These include: 1) Which sst subtypes that are expressed in human sst-positive tumors determine (un)responsiveness to octapeptide SS-analogs such as octreotide or lanreotide, and is there a role for novel sst subtype selective SS-analogs? 2) Which sst subtypes are involved in receptor-mediated endocytosis of radiolabeled SS-analogs and form the basis for targeted radiotherapy or chemotherapy using SS-analogs coupled with radioisotopes or chemotherapeutic compounds, respectively? and 3) What is the role of the individual sst subtypes in determining responsiveness, as well as tachyphylaxis of responsiveness of sst-positive cells upon agonist exposure? In this review, the current knowledge of the clinical consequences of agonist-induced sst internalization for treatment for sst-targeted radiotherapy or chemotherapy are discussed, as well as the different mechanisms that could play a role in tachyphylaxis and/or resistance to SS-analog therapy in patients with neuroendocrine tumors.

The individual sst subtypes differentially internalize SS-analogs. The sst₁ receptors show low agonist-induced internalization, whereas sst₂, sst₃, sst₄, and sst₅ are more efficient in this respect. The predominant expression of sst₂ receptors in most human sst-positive neuroendocrine tumors and the efficiency of sst₂ receptors to undergo agonist-induced internalization is very important for the radiotherapeutic application of radiolabeled octapeptide SS-analogs. *In vitro* studies have demonstrated that sst₂-expressing tumor cell lines, as well as primary cultures of human tumors, internalize radiolabeled SS-analogs such as [¹¹¹In-DTPA⁰]octreotide, [⁹⁰Y-DOTA⁰,Tyr³]octreotide, and [⁹⁰Y-DOTA]lanreotide. Preclinical studies using experimental tumor models have now demonstrated that tumor growth can be inhibited by administration of radiopharmaceutical compounds as [¹¹¹In-DTPA⁰]octreotide and [⁹⁰Y-DOTA⁰,Tyr³]octreotide. Clinical trials have already demonstrated promising effects using these radiopharmaceuticals, as well as of [⁹⁰Y-DOTA-lanreotide], on tumor size in patients with advanced sst-positive neuroendocrine tumors. Finally, the concept of targeted chemotherapy to deliver chemotherapeutic compounds selectively to sst-positive tumor cells, thereby reducing their toxicity, has now been validated using newly developed cytotoxic SS-analogs in experimental mouse and rat models of human pancreatic, breast, prostate, ovarian, and small cell lung cancer.

The presence of sst₂ receptors in tumors is a prerequisite for sensitivity of inhibition of tumor-related hormonal hypersecretion to treatment with octapeptide SS-analogs. The successful clinical application of SS-analogs such as octreotide and lanreotide in the treatment of hormonal hypersecretion in patients with GH-secreting pituitary adenomas and islet cell or carcinoid tumors is caused by the predominant expression of sst₂ receptors in these tumors. On the other hand, novel sst subtype-selective analogs, as well as bispecific and more universal agonists, have been synthesized now and were demonstrated to be effective in the *in*

vitro inhibition of hormone secretion of sst-positive tumors that do not express sst₂ receptors. Patients with sst-positive tumors also show considerable variability in their responsiveness to treatment with SS-analogs. Patients with GH-secreting pituitary adenomas do not show desensitization to treatment with SS-analogs, whereas patients with islet cell or carcinoid tumors often demonstrate tachyphylaxis to treatment. The occurrence of tachyphylaxis upon treatment with SS-analogs is highly variable. Some patients escape very rapidly, whereas others show tachyphylaxis only after several years of treatment. Nevertheless, despite the increasing fundamental knowledge on the role of individual sst subtypes in agonist-induced internalization and/or desensitization of sst subtypes, as well as in agonist-induced, sst subtype-specific regulation of sst expression and receptor homo- and heterodimerization, the direct fundamental evidence for the observed differences between patients with neuroendocrine tumors in the development of tachyphylaxis to SS-analogs is currently weak and requires further studies.

Acknowledgments

Address all correspondence and requests for reprints to: Leo J. Hofland, Department of Internal Medicine, University Hospital Dijkzigt, Dr. Molewaterplein 40, 3015 GD Rotterdam, The Netherlands. E-mail: hofland@inw3.fgg.eur.nl

References

1. **Guillemin R, Gerich JE** 1976 Somatostatin: physiological and clinical significance. *Annu Rev Med* 27:379–388
2. **Lamberts SW, Krenning EP, Reubi JC** 1991 The role of somatostatin and its analogs in the diagnosis and treatment of tumors. *Endocr Rev* 12:450–482
3. **Schally AV** 1988 Oncological applications of somatostatin analogues [published erratum appears in *Cancer Res* 1989 Mar 15; 49(6):1618]. *Cancer Res* 48:6977–6985
4. **Hoyer D, Bell GI, Berelowitz M, Epelbaum J, Feniuk W, Humphrey PP, O'Carroll AM, Patel YC, Schonbrunn A, Taylor JE, Reisine T** 1995 Classification and nomenclature of somatostatin receptors. *Trends Pharmacol Sci* 16:86–88
5. **Hoyer D, Lubbert H, Bruns C** 1994 Molecular pharmacology of somatostatin receptors. *Naunyn Schmiedebergs Arch Pharmacol* 350:441–453
6. **Bruns C, Weckbecker G, Raulf F, Kaupmann K, Schoeffter P, Hoyer D, Lubbert H** 1994 Molecular pharmacology of somatostatin-receptor subtypes. *Ann N Y Acad Sci* 733:138–146
7. **Reisine T, Bell GI** 1995 Molecular biology of somatostatin receptors. *Endocr Rev* 16:427–442
8. **Patel YC** 1997 Molecular pharmacology of somatostatin receptor subtypes. *J Endocrinol Invest* 20:348–367
9. **Patel YC** 1999 Somatostatin and its receptor family. *Front Neuroendocrinol* 20:157–198
10. **Hofland LJ, Visser-Wisselaar HA, Lamberts SW** 1995 Somatostatin analogs: clinical application in relation to human somatostatin receptor subtypes. *Biochem Pharmacol* 50:287–297
11. **Lamberts SW, van der Lely AJ, de Herder WW, Hofland LJ** 1996 Octreotide. *N Engl J Med* 334:246–254
12. **Krenning EP, Bakker WH, Breeman WA, Koper JW, Kooij PP, Ausema L, Lameris JS, Reubi JC, Lamberts SW** 1989 Localisation of endocrine-related tumours with radioiodinated analogue of somatostatin. *Lancet* 1:242–244
13. **Krenning EP, Kwekkeboom DJ, Bakker WH, Breeman WA, Kooij PP, Oei HY, van Hagen M, Postema PT, de Jong M, Reubi JC, Visser TJ, Reys AEM, Hofland LJ, Keper JW, Lamberts SWJ** 1993 Somatostatin receptor scintigraphy with [¹¹¹In-DTPA-D-Phe¹]-

- and [123I-Tyr³]-octreotide: the Rotterdam experience with more than 1000 patients. *Eur J Nucl Med* 20:716–731
14. Reichlin S 1983 Somatostatin. *N Engl J Med* 309:1495–1501
 15. Epelbaum J 1986 Somatostatin in the central nervous system: physiology and pathological modifications. *Prog Neurobiol* 27:63–100
 16. Sevarino KA, Felix R, Banks CM, Low MJ, Montminy MR, Mandel G, Goodman RH 1987 Cell-specific processing of preprosomatostatin in cultured neuroendocrine cells. *J Biol Chem* 262:4987–4993
 17. Vanetti M, Kouba M, Wang X, Vogt G, Hollt V 1992 Cloning and expression of a novel mouse somatostatin receptor (SSTR2B). *FEBS Lett* 311:290–294
 18. Patel YC, Greenwood M, Kent G, Panetta R, Srikant CB 1993 Multiple gene transcripts of the somatostatin receptor SSTR2: tissue selective distribution and cAMP regulation. *Biochem Biophys Res Commun* 192:288–294
 19. Dournaud P, Boudin H, Schonbrunn A, Tannenbaum GS, Beaudet A 1998 Interrelationships between somatostatin sst2A receptors and somatostatin-containing axons in rat brain: evidence for regulation of cell surface receptors by endogenous somatostatin. *J Neurosci* 18:1056–1071
 20. Panetta R, Patel YC 1995 Expression of mRNA for all five human somatostatin receptors (hSSTR1–5) in pituitary tumors. *Life Sci* 56:333–342
 21. Reubi JC, Kappeler A, Waser B, Schonbrunn A, Laissue J 1998 Immunohistochemical localization of somatostatin receptor sst2A in human pancreatic islets. *J Clin Endocrinol Metab* 83:3746–3749
 22. Kumar U, Sasi R, Suresh S, Patel A, Thangaraju M, Metrakos P, Patel SC, Patel YC 1999 Subtype-selective expression of the five somatostatin receptors (hSSTR1–5) in human pancreatic islet cells: a quantitative double-label immunohistochemical analysis. *Diabetes* 48:77–85
 23. Reubi JC, Krenning E, Lamberts SW, Kvols L 1992 *In vitro* detection of somatostatin receptors in human tumors. *Metabolism* 41:104–110
 24. Reubi JC, Laissue J, Krenning E, Lamberts SW 1992 Somatostatin receptors in human cancer: incidence, characteristics, functional correlates and clinical implications. *J Steroid Biochem Mol Biol* 43:27–35
 25. Reubi JC, Laissue J, Waser B, Horisberger U, Schaer JC 1994 Expression of somatostatin receptors in normal, inflamed, and neoplastic human gastrointestinal tissues. *Ann N Y Acad Sci* 733:122–137
 26. Reubi JC, Schaer JC, Laissue JA, Waser B 1996 Somatostatin receptors and their subtypes in human tumors and in peritumoral vessels. *Metabolism* 45:39–41
 27. Vikić-Topić S, Raisch KP, Kvols LK, Vuk-Pavlović S 1995 Expression of somatostatin receptor subtypes in breast carcinoma, carcinoid tumor, and renal cell carcinoma. *J Clin Endocrinol Metab* 80:2974–2979
 28. Reubi JC, Schaer JC, Waser B, Mengod G 1994 Expression and localization of somatostatin receptor SSTR1, SSTR2, and SSTR3 messenger RNAs in primary human tumors using *in situ* hybridization. *Cancer Res* 54:3455–3459
 29. Reubi JC, Waser B, Schaer JC, Markwalder R 1995 Somatostatin receptors in human prostate and prostate cancer. *J Clin Endocrinol Metab* 80:2806–2814
 30. Greenman Y, Melmed S 1994 Expression of three somatostatin receptor subtypes in pituitary adenomas: evidence for preferential SSTR5 expression in the mammosomatotroph lineage. *J Clin Endocrinol Metab* 79:724–729
 31. Greenman Y, Melmed S 1994 Heterogeneous expression of two somatostatin receptor subtypes in pituitary tumors. *J Clin Endocrinol Metab* 78:398–403
 32. Kubota A, Yamada Y, Kagimoto S, Shimatsu A, Imamura M, Tsuda K, Imura H, Seino S, Seino Y 1994 Identification of somatostatin receptor subtypes and an implication for the efficacy of somatostatin analogue SMS 201-995 in treatment of human endocrine tumors. *J Clin Invest* 93:1321–1325
 33. Epelbaum J, Bertherat J, Prevost G, Kordon C, Meyerhof W, Wulfsen I, Richter D, Plouin P F 1995 Molecular and pharmacological characterization of somatostatin receptor subtypes in adrenal, extraadrenal, and malignant pheochromocytomas. *J Clin Endocrinol Metab* 80:1837–1844
 34. Miller GM, Alexander JM, Bikkal HA, Katznelson L, Zervas NT, Klibanski A 1995 Somatostatin receptor subtype gene expression in pituitary adenomas. *J Clin Endocrinol Metab* 80:1386–1392
 35. Reubi JC, Kappeler A, Waser B, Laissue J, Hipkin RW, Schonbrunn A 1998 Immunohistochemical localization of somatostatin receptors sst2A in human tumors. *Am J Pathol* 153:233–245
 36. Hofland LJ, Liu Q, Van Koetsveld PM, Zuijderwijk J, Van Der Ham F, De Krijger RR, Schonbrunn A, Lamberts SW 1999 Immunohistochemical detection of somatostatin receptor subtypes sst1 and sst2A in human somatostatin receptor positive tumors. *J Clin Endocrinol Metab* 84:775–780
 37. Schulz S, Schmitt J, Wiborny D, Schmidt H, Olbricht S, Weise W, Roessner A, Gramsch C, Hollt V 1998 Immunocytochemical detection of somatostatin receptors sst1, sst2A, sst2B, and sst3 in paraffin-embedded breast cancer tissue using subtype-specific antibodies. *Clin Cancer Res* 4:2047–2052
 38. Kimura N, Pilichowska M, Date F, Kimura I, Schindler M 1999 Immunohistochemical expression of somatostatin type 2A receptor in neuroendocrine tumors. *Clin Cancer Res* 5:3483–3487
 39. Reubi JC, Horisberger U, Laissue J 1994 High density of somatostatin receptors in veins surrounding human cancer tissue: role in tumor-host interaction? *Int J Cancer* 56:681–688
 40. Reubi JC, Waser B, Horisberger U, Krenning E, Lamberts SW, Gebbers JO, Gersbach P, Laissue JA 1993 *In vitro* autoradiographic and *in vivo* scintigraphic localization of somatostatin receptors in human lymphatic tissue. *Blood* 82:2143–2151
 41. Reubi JC, Waser B, Schmassmann A, Laissue JA 1999 Receptor autoradiographic evaluation of cholecystokinin, neurotensin, somatostatin and vasoactive intestinal peptide receptors in gastrointestinal adenocarcinoma samples: where are they really located? *Int J Cancer* 81:376–386
 42. Patel YC, Greenwood MT, Panetta R, Demchyshyn L, Niznik H, Srikant CB 1995 The somatostatin receptor family. *Life Sci* 57:1249–1265
 43. Beaudet A, Nouel D, Stroh T, Vandebulcke F, Dal-Farra C, Vincent JP 1998 Fluorescent ligands for studying neuropeptide receptors by confocal microscopy. *Braz J Med Biol Res* 31:1479–1489
 44. Röth A, Kreienkamp HJ, Nehring RB, Roosterman D, Meyerhof W, Richter D 1997 Endocytosis of the rat somatostatin receptors: subtype discrimination, ligand specificity, and delineation of carboxy-terminal positive and negative sequence motifs. *DNA Cell Biol* 16:111–119
 45. Csaba Z, Dournaud P 2001 Cellular biology of somatostatin receptors. *Neuropeptides* 35:1–23
 46. Schonbrunn A 1999 Somatostatin receptors present knowledge and future directions. *Ann Oncol* 10:S17–S21
 47. Koeng JA, Edwardson JM 1997 Endocytosis and recycling of G protein-coupled receptors. *Trends Pharmacol Sci* 18:276–287
 48. Hausdorff WP, Caron MG, Lefkowitz RJ 1990 Turning off the signal: desensitization of β -adrenergic receptor function [published erratum appears in *FASEB J* 1990 Sep;4(12):3049]. *FASEB J* 4:2881–2889
 49. Ferguson SS, Barak LS, Zhang J, Caron MG 1996 G-protein-coupled receptor regulation: role of G-protein-coupled receptor kinases and arrestins. *Can J Physiol Pharmacol* 74:1095–1110
 50. Yu SS, Lefkowitz RJ, Hausdorff WP 1993 β -adrenergic receptor sequestration. A potential mechanism of receptor resensitization. *J Biol Chem* 268:337–341
 51. Roettger BF, Rentsch RU, Pinon D, Holicky E, Hadac E, Larkin JM, Miller LJ 1995 Dual pathways of internalization of the cholecystokinin receptor. *J Cell Biol* 128:1029–1041
 52. Brown MS, Goldstein JL 1986 A receptor-mediated pathway for cholesterol homeostasis. *Science* 232:34–47
 53. Steinman RM, Mellman IS, Muller WA, Cohn ZA 1983 Endocytosis and the recycling of plasma membrane. *J Cell Biol* 96:1–27
 54. Hukovic N, Panetta R, Kumar U, Patel YC 1996 Agonist-dependent regulation of cloned human somatostatin receptor types 1–5 (hSSTR1–5): subtype selective internalization or upregulation. *Endocrinology* 137:4046–4049
 55. Nouel D, Gaudriault G, Houle M, Reisine T, Vincent JP, Mazella J, Beaudet A 1997 Differential internalization of somatostatin in COS-7 cells transfected with SST1 and SST2 receptor subtypes: a confocal microscopic study using novel fluorescent somatostatin derivatives. *Endocrinology* 138:296–306

56. Rocheville M, Lange DC, Kumar U, Sasi R, Patel RC, Patel YC 2000 Subtypes of the somatostatin receptor assemble as functional homo- and heterodimers. *J Biol Chem* 275:7862–7869
57. Hukovic N, Rocheville M, Kumar U, Sasi R, Khare S, Patel YC 1999 Agonist-dependent up-regulation of human somatostatin receptor type 1 requires molecular signals in the cytoplasmic C-tail. *J Biol Chem* 274:24550–24558
58. Pfeiffer M, Koch T, Schroder H, Klutzny M, Kirscht S, Kreienkamp HJ, Hollt V, Schulz S 2001 Homo- and heterodimerization of somatostatin receptor subtypes. Inactivation of sst(3) receptor function by heterodimerization with sst(2A). *J Biol Chem* 276:14027–14036
59. Pfeiffer M, Koch T, Schroder H, Laugsch M, Hollt V, Schulz S 2002 Heterodimerization of somatostatin and opioid receptors cross-modulates phosphorylation, internalization, and desensitization. *J Biol Chem* 277:19762–19772
60. Rocheville M, Lange DC, Kumar U, Patel SC, Patel RC, Patel YC 2000 Receptors for dopamine and somatostatin: formation of hetero-oligomers with enhanced functional activity. *Science* 288:154–157
61. Breeman WA, Kwekkeboom DJ, Kooij PP, Bakker WH, Hofland LJ, Visser TJ, Ensing GJ, Lamberts SW, Krenning EP 1995 Effect of dose and specific activity on tissue distribution of indium-111-pentetreotide in rats. *J Nucl Med* 36:623–627
62. Szegedi Z, Takacs J, Szende B, Vadasz Z, Horvath A, Gulyas E, Toth G, Petak I, Bocsi J, Keri G 1999 A specifically radiolabeled somatostatin analog with strong antitumor activity induces apoptosis and accumulates in the cytosol and the nucleus of HT29 human colon carcinoma cells. *Endocrine* 10:25–34
63. Andersson P, Forssell-Aronsson E, Johanson V, Wangberg B, Nilsson O, Fjalling M, Ahlman H 1996 Internalization of indium-111 into human neuroendocrine tumor cells after incubation with indium-111-DTPA-D-Phe1-octreotide. *J Nucl Med* 37:2002–2006
64. Janson ET, Westlin JE, Ohrvall U, Oberg K, Lukinius A 2000 Nuclear localization of ¹¹¹In after intravenous injection of [¹¹¹In-DTPA-D-Phe1]-octreotide in patients with neuroendocrine tumors. *J Nucl Med* 41:1514–1518
65. Hofland LJ, van Koetsveld PM, Waaijers M, Zuyderwijk J, Breeman WA, Lamberts SW 1995 Internalization of the radioiodinated somatostatin analog [¹²⁵I-Tyr3]octreotide by mouse and human pituitary tumor cells: increase by unlabeled octreotide. *Endocrinology* 136:3698–3706
66. Koenig JA, Edwardson JM, Humphrey PP 1997 Somatostatin receptors in Neuro2A neuroblastoma cells: ligand internalization. *Br J Pharmacol* 120:52–59
67. de Jong M, Breeman WA, Bakker WH, Kooij PP, Bernard BF, Hofland LJ, Visser TJ, Srinivasan A, Schmidt MA, Erion JL, Bugaj JE, Macke HR, Krenning EP 1998 Comparison of (111)In-labeled somatostatin analogues for tumor scintigraphy and radionuclide therapy. *Cancer Res* 58:437–441
68. Hipkin RW, Friedman J, Clark RB, Eppler CM, Schonbrunn A 1997 Agonist-induced desensitization, internalization, and phosphorylation of the sst2A somatostatin receptor. *J Biol Chem* 272:13869–13876
69. Janson ET, Stridsberg M, Gobl A, Westlin JE, Oberg K 1998 Determination of somatostatin receptor subtype 2 in carcinoid tumors by immunohistochemical investigation with somatostatin receptor subtype 2 antibodies. *Cancer Res* 58:2375–2378
70. Briganti V, Sestini R, Orlando C, Bernini G, La Cava G, Tamburini A, Raggi CC, Serio M, Maggi M 1997 Imaging of somatostatin receptors by indium-111-pentetreotide correlates with quantitative determination of somatostatin receptor type 2 gene expression in neuroblastoma tumors. *Clin Cancer Res* 3:2385–2391
71. Ferone D, van Hagen MP, Kwekkeboom DJ, van Koetsveld PM, Mooy DM, Lichtenauer-Kaligis E, Schonbrunn A, Colao A, Lamberts SW, Hofland LJ 2000 Somatostatin receptor subtypes in human thymoma and inhibition of cell proliferation by octreotide *in vitro*. *J Clin Endocrinol Metab* 85:1719–1726
72. Kolby L, Wangberg B, Ahlman H, Tisell LE, Fjalling M, Forssell-Aronsson E, Nilsson O 1998 Somatostatin receptor subtypes, octreotide scintigraphy, and clinical response to octreotide treatment in patients with neuroendocrine tumors. *World J Surg* 22:679–683
73. Kristiansen MT, Rasmussen LM, Olsen N, Asa SL, Jorgensen JO 2002 Ectopic ACTH syndrome: discrepancy between somatostatin receptor status *in vivo* and *ex vivo*, and between immunostaining and gene transcription for POMC and CRH. *Horm Res* 57:200–204
74. Shimon I, Taylor JE, Dong JZ, Bitonte RA, Kim S, Morgan B, Coy DH, Culler MD, Melmed S 1997 Somatostatin receptor subtype specificity in human fetal pituitary cultures. Differential role of SSTR2 and SSTR5 for growth hormone, thyroid-stimulating hormone, and prolactin regulation. *J Clin Invest* 99:789–798
75. Shimon I, Yan X, Taylor JE, Weiss MH, Culler MD, Melmed S 1997 Somatostatin receptor (SSTR) subtype-selective analogues differentially suppress *in vitro* growth hormone and prolactin in human pituitary adenomas. Novel potential therapy for functional pituitary tumors. *J Clin Invest* 100:2386–2392
76. Stolz B, Weckbecker G, Smith-Jones PM, Albert R, Raulf F, Bruns C 1998 The somatostatin receptor-targeted radiotherapeutic [90Y-DOTA-D-Phe1, Tyr3]octreotide (90Y-SMT 487) eradicates experimental rat pancreatic CA 20948 tumours. *Eur J Nucl Med* 25:668–674
77. Hofland LJ, Breeman WA, Krenning EP, de Jong M, Waaijers M, van Koetsveld PM, Macke HR, Lamberts SW 1999 Internalization of [DOTA⁰,¹²⁵I-Tyr3]octreotide by somatostatin receptor-positive cells *in vitro* and *in vivo*: implications for somatostatin receptor-targeted radio-guided surgery. *Proc Assoc Am Physicians* 111:63–69
78. De Jong M, Bernard BF, De Bruin E, Van Gameren A, Bakker WH, Visser TJ, Macke HR, Krenning EP 1998 Internalization of radiolabelled [DTPA0]octreotide and [DOTA0,Tyr3]octreotide: peptides for somatostatin receptor-targeted scintigraphy and radionuclide therapy. *Nucl Med Commun* 19:283–288
79. Kwekkeboom DJ, Kooij PP, Bakker WH, Macke HR, Krenning EP 1999 Comparison of ¹¹¹In-DOTA-Tyr3-octreotide and ¹¹¹In-DTPA-octreotide in the same patients: biodistribution, kinetics, organ and tumor uptake. *J Nucl Med* 40:762–767
80. Koenig JA, Kaur R, Dodgeon I, Edwardson JM, Humphrey PPA 1998 Fates of endocytosed somatostatin sst2 receptors and associated agonists. *Biochem J* 336:291–298
81. Roth A, Kreienkamp HJ, Meyerhof W, Richter D 1997 Phosphorylation of four amino acid residues in the carboxyl terminus of the rat somatostatin receptor subtype 3 is crucial for its desensitization and internalization. *J Biol Chem* 272:23769–23774
82. Duncan JR, Stephenson MT, Wu HP, Anderson CJ 1997 Indium-111-diethylenetriaminepentaacetic acid-octreotide is delivered *in vivo* to pancreatic, tumor cell, renal, and hepatocyte lysosomes. *Cancer Res* 57:659–671
83. Bass LA, Lanahan MV, Duncan JR, Erion JL, Srinivasan A, Schmidt MA, Anderson CJ 1998 Identification of the soluble *in vivo* metabolites of indium-111-diethylenetriaminepentaacetic acid-D-Phe1-octreotide. *Bioconjug Chem* 9:192–200
84. Akizawa H, Arano Y, Uezono T, Ono M, Fujioka Y, Uehara T, Yokoyama A, Akaji K, Kiso Y, Koizumi M, Saji H 1998 Renal metabolism of ¹¹¹In-DTPA-D-Phe1-octreotide *in vivo*. *Bioconjug Chem* 9:662–670
85. Hornick CA, Anthony CT, Hughey S, Gebhardt BM, Espenan GD, Woltering EA 2000 Progressive nuclear translocation of somatostatin analogs. *J Nucl Med* 41:1256–1263
86. Kooij PPM, Kwekkeboom DJ, Breeman WAP, Reijs AEM, Bakker WH, Lamberts SWJ, Visser TJ, Krenning EP 1994 The effects of specific activity on tissue distribution of [¹¹¹In-111-DTPA-D-Phe1]octreotide in humans. *J Nucl Med* 35:226P
87. de Jong M, Breeman WA, Bernard BF, van Gameren A, de Bruin E, Bakker WH, van der Pluijm ME, Visser TJ, Macke HR, Krenning EP 1999 Tumour uptake of the radiolabelled somatostatin analogue [DOTA0, TYR3]octreotide is dependent on the peptide amount. *Eur J Nucl Med* 26:693–698
88. Breeman WA, Hofland LJ, van der Pluijm M, van Koetsveld PM, de Jong M, Setyono-Han B, Bakker WH, Kwekkeboom DJ, Visser TJ, Lamberts SW 1994 A new radiolabelled somatostatin analogue [¹¹¹In-DTPA-D-Phe1]RC-160: preparation, biological activity, receptor scintigraphy in rats and comparison with [¹¹¹In-DTPA-D-Phe1]octreotide. *Eur J Nucl Med* 21:328–335
89. Breeman WAP, van Hagen PM, Kwekkeboom DJ, Visser TJ, Krenning EP 1998 Somatostatin receptor scintigraphy using [¹¹¹In-DTPA0]RC-160 in humans: a comparison with [¹¹¹In-DTPA0]octreotide. *Eur J Nucl Med* 25:182–186

90. Smith-Jones PM, Bischof C, Leimer M, Gludovacz D, Angelberger P, Pangerl T, Peck-Radosavljevic M, Hamilton G, Kaserer K, Kofler A, Schlangbauer-Wadl H, Traub T, Virgolini I 1999 DOTA-lanreotide: a novel somatostatin analog for tumor diagnosis and therapy. *Endocrinology* 140:5136–5148
91. Reubi JC, Schar JC, Waser B, Wenger S, Heppeler A, Schmitt JS, Macke HR 2000 Affinity profiles for human somatostatin receptor subtypes SST1–SST5 of somatostatin radiotracers selected for scintigraphic and radiotherapeutic use. *Eur J Nucl Med* 27:273–282
92. Ronga G, Salerno G, Procaccini E, Mauro L, Annovazzi A, Barone R, Mellozzi M, Tamburrano G, Signore A 1995 ¹¹¹In-octreotide scintigraphy in metastatic medullary thyroid carcinoma before and after octreotide therapy: *in vivo* evidence of the possible down-regulation of somatostatin receptors. *Q J Nucl Med* 39:134–136
93. Krassas GE, Doulas A, Kaltsas T, Halkias A, Pontikides N 1999 Somatostatin receptor scintigraphy before and after treatment with somatostatin analogues in patients with thyroid eye disease. *Thyroid* 9:47–52
94. Schillaci O, Annibale B, Scopinaro F, delle Fave G, Colella AC 1997 Somatostatin receptor scintigraphy of malignant somatostatinoma with indium-111-pentetreotide. *J Nucl Med* 38:886–887
95. Lamberts SW, Hofland LJ, van Koetsveld PM, Reubi JC, Bruining HA, Bakker WH, Krenning EP 1990 Parallel *in vivo* and *in vitro* detection of functional somatostatin receptors in human endocrine pancreatic tumors: consequences with regard to diagnosis, localization, and therapy. *J Clin Endocrinol Metab* 71:566–574
96. Dorr U, Wurm K, Horing E, Guzman G, Rath U, Bihl H 1993 Diagnostic reliability of somatostatin receptor scintigraphy during continuous treatment with different somatostatin analogs. *Horm Metab Res Suppl* 27:36–43
97. Zamora PO, Gulhke S, Bender H, Diekmann D, Rhodes BA, Biersack HJ, Knapp Jr FF 1996 Experimental radiotherapy of receptor-positive human prostate adenocarcinoma with ¹⁸⁸Re-RC-160, a directly-radiolabeled somatostatin analogue. *Int J Cancer* 65:214–220
98. Zamora PO, Bender H, Gulhke S, Marek MJ, Knapp Jr FF, Rhodes BA, Biersack HJ 1997 Pre-clinical experience with Re-188-RC-160, a radiolabeled somatostatin analog for use in peptide-targeted radiotherapy. *Anticancer Res* 17:1803–1808
99. Stolz B, Smith-Jones P, Albert R, Tolcsvai L, Briner U, Ruser G, Macke H, Weckbecker G, Bruns C 1996 Somatostatin analogues for somatostatin-receptor-mediated radiotherapy of cancer. *Digestion* 57:17–21
100. Slooter GD, Breeman WA, Marquet RL, Krenning EP, van Eijck CH 1999 Anti-proliferative effect of radiolabelled octreotide in a metastases model in rat liver. *Int J Cancer* 81:767–771
101. de Jong M, Breeman WA, Bernard BF, Bakker WH, Schaar M, van Garamen A, Bugaj JE, Erion J, Schmidt M, Srinivasan A, Krenning EP 2001 [¹⁷⁷Lu-DOTA(0),Tyr3] octreotate for somatostatin receptor-targeted radionuclide therapy. *Int J Cancer* 92:628–633
102. Krenning EP, de Jong M, Jamar F, Valkema R, Kwekkeboom DJ, Kvols LK, Smith C, Pauwels S 2002 Somatostatin receptor-targeted radiotherapy of tumors: preclinical and clinical findings. In: Lamberts SWJ, Dogliotti L, eds. *The expanding role of octreotide I: advances in oncology*. Bristol, UK: Bioscientifica Ltd.; 211–223
103. de Jong M, Krenning E 2002 New advances in peptide receptor radionuclide therapy. *J Nucl Med* 43:617–620
104. Gallardo D, Drazan KE, McBride WH 1996 Adenovirus-based transfer of wild-type p53 gene increases ovarian tumor radiosensitivity. *Cancer Res* 56:4891–4893
105. Sakakura C, Sweeney EA, Shirahama T, Igarashi Y, Hakomori S, Nakatani H, Tsujimoto H, Imanishi T, Ohgaki M, Ohyama T, Yamazaki J, Hagiwara A, Yamaguchi T, Sawai K, Takahashi T 1996 Overexpression of bax sensitizes human breast cancer MCF-7 cells to radiation-induced apoptosis. *Int J Cancer* 67:101–105
106. Wagener C, Bargou RC, Daniel PT, Bommert K, Mapara MY, Royer HD, Dorken B 1996 Induction of the death-promoting gene bax- α sensitizes cultured breast-cancer cells to drug-induced apoptosis. *Int J Cancer* 67:138–141
107. Sharma K, Srikant CB 1998 Induction of wild-type p53, Bax, and acidic endonuclease during somatostatin-signaled apoptosis in MCF-7 human breast cancer cells. *Int J Cancer* 76:259–266
108. Sharma K, Patel YC, Srikant CB 1996 Subtype-selective induction of wild-type p53 and apoptosis, but not cell cycle arrest, by human somatostatin receptor 3. *Mol Endocrinol* 10:1688–1696
109. Sharma K, Patel YC, Srikant CB 1999 C-terminal region of human somatostatin receptor 5 is required for induction of Rb and G1 cell cycle arrest. *Mol Endocrinol* 13:82–90
110. Pages P, Benali N, Saint-Laurent N, Esteve JP, Schally AV, Tkaczuk J, Vaysse N, Susini C, Buscail L 1999 sst2 Somatostatin receptor mediates cell cycle arrest and induction of p27(Kip1). Evidence for the role of SHP-1. *J Biol Chem* 274:15186–15193
111. Krenning EP, de Jong M, Kooij PP, Breeman WA, Bakker WH, de Herder WW, van Eijck CH, Kwekkeboom DJ, Jamar F, Pauwels S, Valkema R 1999 Radiolabelled somatostatin analogue(s) for peptide receptor scintigraphy and radionuclide therapy. *Ann Oncol* 10:S23–S29
112. Fjalling M, Andersson P, Forssell-Aronsson E, Gretarsdottir J, Johansson V, Tisell LE, Wangberg B, Nilsson O, Berg G, Michanek A, Lindstedt G, Ahlman H 1996 Systemic radionuclide therapy using indium-111-DTPA-D-Phe1-octreotide in midgut carcinoid syndrome. *J Nucl Med* 37:1519–1521
113. Valkema R, De Jong M, Bakker WH, Breeman WA, Kooij PP, Lugtenburg PJ, De Jong FH, Christiansen A, Kam BL, De Herder WW, Stridsberg M, Lindemans J, Ensing G, Krenning EP 2002 Phase I study of peptide receptor radionuclide therapy with [In-DTPA]octreotide: the Rotterdam experience. *Semin Nucl Med* 32:110–122
114. McCarthy KE, Woltering EA, Espenan GD, Cronin M, Maloney TJ, Anthony LB 1998 *In situ* radiotherapy with ¹¹¹In-pentetreotide: initial observations and future directions. *Cancer J Sci Am* 4:94–102
115. Otte A, Mueller-Brand J, Dellas S, Nitzsche EU, Herrmann R, Maecke HR 1998 Yttrium-90-labelled somatostatin-analogue for cancer treatment. *Lancet* 351:417–418
116. Otte A, Herrmann R, Heppeler A, Behe M, Jermann E, Powell P, Maecke HR, Muller J 1999 Yttrium-90 DOTATOC: first clinical results. *Eur J Nucl Med* 26:1439–1447
117. Paganelli G, Zoboli S, Cremonesi M, Macke HR, Chinol M 1999 Receptor-mediated radionuclide therapy with ⁹⁰Y-DOTA-D-Phe1-Tyr3-octreotide: preliminary report in cancer patients. *Cancer Biother Radiopharm* 14:477–483
118. Paganelli G, Zoboli S, Cremonesi M, Bodei L, Ferrari M, Grana C, Bartolomei M, Orsi F, De Cicco C, Macke HR, Chinol M, de Braud F 2001 Receptor-mediated radiotherapy with ⁹⁰Y-DOTA-D-Phe1-Tyr3-octreotide. *Eur J Nucl Med* 28:426–434
119. Waldherr C, Pless M, Maecke HR, Haldemann A, Mueller-Brand J 2001 The clinical value of [⁹⁰Y-DOTA]-D-Phe1-Tyr3-octreotide (⁹⁰Y-DOTATOC) in the treatment of neuroendocrine tumours: a clinical phase II study. *Ann Oncol* 12:941–945
120. Waldherr C, Pless M, Maecke HR, Schumacher T, Crazzolara A, Nitzsche EU, Haldemann A, Mueller-Brand J 2002 Tumor response and clinical benefit in neuroendocrine tumors after 7.4 GBq (⁹⁰Y-DOTATOC). *J Nucl Med* 43:610–616
121. Leimer M, Kurtaran A, Smith-Jones P, Raderer M, Havlik E, Angelberger P, Vorbeck F, Niederle B, Herold C, Virgolini I 1998 Response to treatment with yttrium 90-DOTA-lanreotide of a patient with metastatic gastrinoma. *J Nucl Med* 39:2090–2094
122. Kwekkeboom DJ, Bakker WH, Kooij PP, Konijnenberg MW, Srinivasan A, Erion JL, Schmidt MA, Bugaj JL, de Jong M, Krenning EP 2001 [¹⁷⁷Lu-DOTAOTyr3]octreotate: comparison with [¹¹¹In-DTPAO]octreotide in patients. *Eur J Nucl Med* 28:1319–1325
123. Schally AV, Nagy A 1999 Cancer chemotherapy based on targeting of cytotoxic peptide conjugates to their receptors on tumors. *Eur J Endocrinol* 141:1–14
124. Radulovic S, Nagy A, Szoke B, Schally AV 1992 Cytotoxic analog of somatostatin containing methotrexate inhibits growth of MIA PaCa-2 human pancreatic cancer xenografts in nude mice. *Cancer Lett* 62:263–271
125. Nagy A, Schally AV, Halmos G, Armatis P, Cai RZ, Csernus V, Kovacs M, Koppán M, Szepeshazi K, Kahan Z 1998 Synthesis and biological evaluation of cytotoxic analogs of somatostatin containing doxorubicin or its intensely potent derivative, 2-pyrrolinodoxorubicin. *Proc Natl Acad Sci USA* 95:1794–1799
126. Plonowski A, Schally AV, Nagy A, Sun B, Szepeshazi K 1999 Inhibition of PC-3 human androgen-independent prostate cancer and its metastases by cytotoxic somatostatin analogue AN-238. *Cancer Res* 59:1947–1953

127. **Plonowski A, Schally AV, Nagy A, Kiaris H, Hebert F, Halmos G** 2000 Inhibition of metastatic renal cell carcinomas expressing somatostatin receptors by a targeted cytotoxic analogue of somatostatin AN-238. *Cancer Res* 60:2996–3001
128. **Plonowski A, Schally AV, Koppan M, Nagy A, Arencibia JM, Csernus B, Halmos G** 2001 Inhibition of the UCI-107 human ovarian carcinoma cell line by a targeted cytotoxic analog of somatostatin, AN-238. *Cancer* 92:1168–1176
129. **Kiaris H, Schally AV, Nagy A, Sun B, Szepeshazi K, Halmos G** 2000 Regression of U-87 MG human glioblastomas in nude mice after treatment with a cytotoxic somatostatin analog AN-238. *Clin Cancer Res* 6:709–717
130. **Kiaris H, Schally AV, Nagy A, Szepeshazi K, Hebert F, Halmos G** 2001 A targeted cytotoxic somatostatin (SST) analogue, AN-238, inhibits the growth of H-69 small-cell lung carcinoma (SCLC) and H-157 non-SCLC in nude mice. *Eur J Cancer* 37:620–628
131. **Szepeshazi K, Schally AV, Halmos G, Sun B, Hebert F, Csernus B, Nagy A** 2001 Targeting of cytotoxic somatostatin analog AN-238 to somatostatin receptor subtypes 5 and/or 3 in experimental pancreatic cancers. *Clin Cancer Res* 7:2854–2861
132. **Kahan Z, Nagy A, Schally AV, Hebert F, Sun B, Groot K, Halmos G** 1999 Inhibition of growth of MX-1, MCF-7-MIII and MDA-MB-231 human breast cancer xenografts after administration of a targeted cytotoxic analog of somatostatin, AN-238. *Int J Cancer* 82:592–598
133. **Koppan M, Nagy A, Schally AV, Arencibia JM, Plonowski A, Halmos G** 1998 Targeted cytotoxic analogue of somatostatin AN-238 inhibits growth of androgen-independent Dunning R-3327-AT-1 prostate cancer in rats at nontoxic doses. *Cancer Res* 58:4132–4137
134. **Ezzat S, Snyder PJ, Young WF, Boyajy LD, Newman C, Klibanski A, Molitch ME, Boyd AE, Sheeler L, Cook DM, Malarkey WB, Jackson I, Vance ML, Thorne MO, Barkan AL, Frohman LA, Melmed S** 1992 Octreotide treatment of acromegaly. A randomized, multicenter study. *Ann Intern Med* 117:711–718
135. **Lamberts SW** 1988 The role of somatostatin in the regulation of anterior pituitary hormone secretion and the use of its analogs in the treatment of human pituitary tumors. *Endocr Rev* 9:417–436
- 135a. **Davies PH, Stewart SE, Lancranjan L, Sheppard MC, Stewart PM** 1998 Long-term therapy with long-acting octreotide (Sandostatin-LAR) for the management of acromegaly. *Clin Endocrinol (Oxf)* 48:311–316
- 135b. **Flogstad AK, Halse J, Bakke S, Lancranjan I, Marbach P, Bruns C, Jervell J** 1997 Sandostatin LAR in acromegalic patients: long-term treatment. *J Clin Endocrinol Metab* 82:23–28
- 135c. **Gillis JC, Noble S, Goa KL** 1997 Octreotide long-acting release (LAR). A review of its pharmacological properties and therapeutic use in the management of acromegaly. *Drugs* 53:681–699
136. **Caron P, Morange-Ramos I, Cogne M, Jaquet P** 1997 Three year follow-up of acromegalic patients treated with intramuscular slow-release lanreotide. *J Clin Endocrinol Metab* 82:18–22
137. **Haraguchi K, Ohtaka M, Takazawa K, Endo T, Onaya T** 1995 Desensitization to somatostatin analogue (octreotide) observed in a patient with acromegaly. *Endocr J* 42:295–300
138. **Wahid ST, Marbach P, Stolz B, Miller M, James RA, Ball SG** 2002 Partial tachyphylaxis to somatostatin (SST) analogues in a patient with acromegaly: the role of SST receptor desensitisation and circulating antibodies to SST analogues. *Eur J Endocrinol* 146:295–302
139. **Chanson P, Weintraub BD, Harris AG** 1993 Octreotide therapy for thyroid-stimulating hormone-secreting pituitary adenomas. A follow-up of 52 patients. *Ann Intern Med* 119:236–240
140. **de Bruin TW, Kwakkeboom DJ, Van't Verlaat JW, Reubi JC, Krenning EP, Lamberts SW, Croughs RJ** 1992 Clinically nonfunctioning pituitary adenoma and octreotide response to long term high dose treatment, and studies *in vitro*. *J Clin Endocrinol Metab* 75:1310–1317
141. **Beck-Peccoz P, Brucker-Davis F, Persani L, Smallridge RC, Weintraub BD** 1996 Thyrotropin-secreting pituitary tumors. *Endocr Rev* 17:610–638
142. **Kvols LK, Moertel CG, O'Connell MJ, Schutt AJ, Rubin J, Hahn RG** 1986 Treatment of the malignant carcinoid syndrome. Evaluation of a long-acting somatostatin analogue. *N Engl J Med* 315:663–666
143. **Kvols LK, Buck M, Moertel CG, Schutt AJ, Rubin J, O'Connell MJ, Hahn RG** 1987 Treatment of metastatic islet cell carcinoma with a somatostatin analogue (SMS 201-995). *Ann Intern Med* 107:162–168
144. **Wymenga AN, Eriksson B, Salmela PI, Jacobsen MB, Van Cutsem EJ, Fiase RH, Valimaki MJ, Renstrup J, de Vries EG, Oberg KE** 1999 Efficacy and safety of prolonged-release lanreotide in patients with gastrointestinal neuroendocrine tumors and hormone-related symptoms. *J Clin Oncol* 17:1111
145. **Ruszniewski P, Ducreux M, Chayvialle JA, Blumberg J, Cloarec D, Michel H, Raymond JM, Dupas JL, Gouerou H, Jian R, Genestien E, Bernades P, Rougier P** 1996 Treatment of the carcinoid syndrome with the longacting somatostatin analogue lanreotide: a prospective study in 39 patients. *Gut* 39:279–283
146. **Moertel CG** 1987 Karmofsky memorial lecture. An odyssey in the land of small tumors. *J Clin Oncol* 5:1502–1522
147. **Lamberts SW, Pieters GF, Metselaar HJ, Ong GL, Tan HS, Reubi JC** 1988 Development of resistance to a long-acting somatostatin analogue during treatment of two patients with metastatic endocrine pancreatic tumours. *Acta Endocrinol (Copenh)* 119:561–566
148. **Siegel RA, Tolcsvai L, Rudin M** 1988 Partial inhibition of the growth of transplanted dunning rat prostate tumors with the long-acting somatostatin analogue sandostatin (SMS 201-995). *Cancer Res* 48:4651–4655
149. **Lamberts SW, Reubi JC, Uiterlinden P, Zuiderwijk J, van den Werff P, van Hal P** 1986 Studies on the mechanism of action of the inhibitory effect of the somatostatin analog SMS 201-995 on the growth of the prolactin/adrenocorticotropin-secreting pituitary tumor 7315a. *Endocrinology* 118:2188–2194
150. **Klijn JGM, Setyono-Han B, Bakker GH** 1988 Prophylactic neuro-peptide-analog treatment of a transplantable pancreatic tumor in rats. *Prog Cancer Res Ther* 35:350
151. **Koper JW, Hofland LJ, van Koetsveld PM, den Holder F, Lamberts SW** 1990 Desensitization and resensitization of rat pituitary tumor cells in long-term culture to the effects of the somatostatin analogue SMS 201-995 on cell growth and prolactin secretion. *Cancer Res* 50:6238–6242
152. **Pinski J, Halmos G, Yano T, Szepeshazi K, Qin Y, Ertl T, Schally AV** 1994 Inhibition of growth of MKN45 human gastric-carcinoma xenografts in nude mice by treatment with bombesin/gastrin-releasing-peptide antagonist (RC-3095) and somatostatin analogue RC-160. *Int J Cancer* 57:574–580
153. **Froidevaux S, Hintermann E, Torok M, Macke HR, Beglinger C, Eberle AN** 1999 Differential regulation of somatostatin receptor type 2 (sst 2) expression in AR4-2J tumor cells implanted into mice during octreotide treatment. *Cancer Res* 59:3652–3657
154. **Mahy N, Woolkalis M, Manning D, Reisine T** 1988 Characteristics of somatostatin desensitization in the pituitary tumor cell line AtT-20. *J Pharmacol Exp Ther* 247:390–396
155. **Srikant CB, Heisler S** 1985 Relationship between receptor binding and biopotency of somatostatin-14 and somatostatin-28 in mouse pituitary tumor cells. *Endocrinology* 117:271–278
156. **Presky DH, Schonbrunn A** 1988 Somatostatin pretreatment increases the number of somatostatin receptors in GH4C1 pituitary cells and does not reduce cellular responsiveness to somatostatin. *J Biol Chem* 263:714–721
157. **Bruno JF, Xu Y, Berelowitz M** 1994 Somatostatin regulates somatostatin receptor subtype mRNA expression in GH3 cells. *Biochem Biophys Res Commun* 202:1738–1743
158. **Vanetti M, Vogt G, Holtt V** 1993 The two isoforms of the mouse somatostatin receptor (mSSTR2A and mSSTR2B) differ in coupling efficiency to adenylate cyclase and in agonist-induced receptor desensitization. *FEBS Lett* 331:260–266
159. **Rens-Domiano S, Law SF, Yamada Y, Seino S, Bell GI, Reisine T** 1992 Pharmacological properties of two cloned somatostatin receptors. *Mol Pharmacol* 42:28–34
160. **Schonbrunn A** 1982 Glucocorticoids down-regulate somatostatin receptors on pituitary cells in culture. *Endocrinology* 110:1147–1154
161. **Xu Y, Berelowitz M, Bruno JF** 1995 Dexamethasone regulates somatostatin receptor subtype messenger ribonucleic acid expression in rat pituitary GH4C1 cells. *Endocrinology* 136:5070–5075
162. **Stalla GK, Brockmeier SJ, Renner U, Newton C, Buchfelder M, Stalla J, Muller OA** 1994 Octreotide exerts different effects *in vivo* and *in vitro* in Cushing's disease. *Eur J Endocrinol* 130:125–131
163. **Visser-Wisselaar HA, Van Uffelen CJ, Van Koetsveld PM,**

- Lichtenauer-Kaligis EG, Waaijers AM, Uitterlinden P, Mooy DM, Lamberts SW, Hofland LJ 1997 17- β -Estradiol-dependent regulation of somatostatin receptor subtype expression in the 7315b prolactin secreting rat pituitary tumor *in vitro* and *in vivo*. *Endocrinology* 138:1180–1189
164. Kimura N, Hayafuji C, Konagaya H, Takahashi K 1986 17- β -Estradiol induces somatostatin (SRIF) inhibition of prolactin release and regulates SRIF receptors in rat anterior pituitary cells. *Endocrinology* 119:1028–1036
165. Kimura N, Hayafuji C 1989 Characterization of 17- β -estradiol-dependent and -independent somatostatin receptor subtypes in rat anterior pituitary. *J Biol Chem* 264:7033–7040
166. Kimura N, Tomizawa S, Arai KN 1998 Chronic treatment with estrogen up-regulates expression of sst2 messenger ribonucleic acid (mRNA) but down-regulates expression of sst5 mRNA in rat pituitaries. *Endocrinology* 139:1573–1580
167. Xu Y, Song J, Berelowitz M, Bruno JF 1996 Estrogen regulates somatostatin receptor subtype 2 messenger ribonucleic acid expression in human breast cancer cells. *Endocrinology* 137:5634–5640
168. Xu Y, Berelowitz M, Bruno JF 1998 Characterization of the promoter region of the human somatostatin receptor subtype 2 gene and localization of sequences required for estrogen-responsiveness. *Mol Cell Endocrinol* 139:71–77
169. Kimura N, Tomizawa S, Arai KN, Osamura RY 2001 Characterization of 5'-flanking region of rat somatostatin receptor sst2 gene: transcriptional regulatory elements and activation by Pitx1 and estrogen. *Endocrinology* 142:1427–1441
170. Viguerie N, Esteve JP, Susini C, Logsdon CD, Vaysse N, Ribet A 1987 Dexamethasone effects on somatostatin receptors in pancreatic acinar AR4-2J cells. *Biochem Biophys Res Commun* 147:942–948
171. James RA, Sarapura VD, Bruns C, Raulf F, Dowding JM, Gordon DF, Wood WM, Ridgway EC 1997 Thyroid hormone-induced expression of specific somatostatin receptor subtypes correlates with involution of the TtT-97 murine thyrotrope tumor. *Endocrinology* 138:719–724
172. Candi E, Melino G, De Laurenzi V, Piacentini M, Guerrieri P, Spinedi A, Knight RA 1995 Tamoxifen and somatostatin affect tumours by inducing apoptosis. *Cancer Lett* 96:141–145
173. Reubi JC, Heitz PU, Landolt AM 1987 Visualization of somatostatin receptors and correlation with immunoreactive growth hormone and prolactin in human pituitary adenomas: evidence for different tumor subclasses. *J Clin Endocrinol Metab* 65:65–73
174. Reubi JC, Waser B, Foekens JA, Klijn JG, Lamberts SW, Laissue J 1990 Somatostatin receptor incidence and distribution in breast cancer using receptor autoradiography: relationship to EGF receptors. *Int J Cancer* 46:416–420
175. Zhang CY, Yokogoshi Y, Yoshimoto K, Fujinaka Y, Matsumoto K, Saito S 1995 Point mutation of the somatostatin receptor 2 gene in the human small cell lung cancer cell line COR-L103. *Biochem Biophys Res Commun* 210:805–815
176. Corbetta S, Ballare E, Mantovani G, Lania A, Losa M, Di Blasio AM, Spada A 2001 Somatostatin receptor subtype 2 and 5 in human GH-secreting pituitary adenomas: analysis of gene sequence and mRNA expression. *Eur J Clin Invest* 31:208–214
177. Petersenn S, Heyens M, Ludecke DK, Beil FU, Schulte HM 2000 Absence of somatostatin receptor type 2 A mutations and gip oncogene in pituitary somatotroph adenomas. *Clin Endocrinol (Oxf)* 52:35–42
178. Ballare E, Persani L, Lania AG, Filopanti M, Giammona E, Corbetta S, Mantovani S, Arosio M, Beck-Peccoz P, Faglia G, Spada A 2001 Mutation of somatostatin receptor type 5 in an acromegalic patient resistant to somatostatin analog treatment. *J Clin Endocrinol Metab* 86:3809–3814
179. Abel F, Ejleskar K, Kogner P, Martinsson T 1999 Gain of chromosome arm 17q is associated with unfavourable prognosis in neuroblastoma, but does not involve mutations in the somatostatin receptor 2(SSTR2) gene at 17q24. *Br J Cancer* 81:1402–1409
180. Kwekkeboom DJ, Assies J, Hofland LJ, Reubi JC, Lamberts SW, Krenning EP 1993 A case of antibody formation against octreotide visualized with ¹¹¹In-octreotide scintigraphy. *Clin Endocrinol (Oxf)* 39:239–243; discussion 244
181. Orskov H, Christensen SE, Weeke J, Kaal A, Harris AG 1991 Effects of antibodies against octreotide in two patients with acromegaly. *Clin Endocrinol (Oxf)* 34:395–398
182. Kendall-Taylor P, Chatterjee S, White MC, Harris MM, Davidson K, Besser GM, Wass JA 1989 Octreotide. *Lancet* 2:859–860 (Letter, Comment)
183. Milligan G 1996 Endocrine disorders associated with mutations in guanine nucleotide binding proteins. *Baillieres Clin Endocrinol Metab* 10:177–187
184. Barlier A, Pellegrini-Bouiller I, Gunz G, Zamora AJ, Jaquet P, Enjalbert A 1999 Impact of gsp oncogene on the expression of genes coding for Gs α , Pit-1, Gi2 α , and somatostatin receptor 2 in human somatotroph adenomas: involvement in octreotide sensitivity. *J Clin Endocrinol Metab* 84:2759–2765
185. Luthin DR, Eppler CM, Linden J 1993 Identification and quantification of Gi-type GTP-binding proteins that copurify with a pituitary somatostatin receptor. *J Biol Chem* 268:5990–5996
186. Lyons J, Landis CA, Harsh G, Vallar L, Grunewald K, Feichtinger H, Duh QY, Clark OH, Kawasaki E, Bourne HR, McCormick F 1990 Two G protein oncogenes in human endocrine tumors. *Science* 249:655–659
187. Saveanu A, Gunz G, Dufour H, Caron P, Fina F, Ouafik L, Culler MD, Moreau JP, Enjalbert A, Jaquet P 2001 Bim-23244, a somatostatin receptor subtype 2- and 5-selective analog with enhanced efficacy in suppressing growth hormone (GH) from octreotide-resistant human GH-secreting adenomas. *J Clin Endocrinol Metab* 86:140–145
188. Bruns C, Lewis I, Briner U, Meno-Tetang G, Weckbecker G 2002 SOM230: a novel somatostatin peptidomimetic with broad somatotropin release inhibiting factor (SRIF) receptor binding and a unique antisecretory profile. *Eur J Endocrinol* 146:707–716
189. Lamberts SW, Verleun T, Zuiderwijk JM, Oosterom R 1987 The effect of the somatostatin analog SMS 201-995 on normal growth hormone secretion in the rat. A comparison with the effect of bromocriptine on normal prolactin secretion. *Acta Endocrinol (Copenh)* 115:196–202
190. Saveanu A, Lavaque E, Gunz G, Barlier A, Kim S, Taylor JE, Culler MD, Enjalbert A, Jaquet P 2002 Demonstration of enhanced potency of a chimeric somatostatin-dopamine molecule, BIM-23A387, in suppressing growth hormone and prolactin secretion from human pituitary somatotroph adenoma cells. *J Clin Endocrinol Metab* 87:5545–5552
191. Patel YC, Srikant CB 1994 Subtype selectivity of peptide analogs for all five cloned human somatostatin receptors (hsstr 1–5). *Endocrinology* 135:2814–2817
192. Schaer JC, Waser B, Mengod G, Reubi JC 1997 Somatostatin receptor subtypes sst1, sst2, sst3 and sst5 expression in human pituitary, gastroentero-pancreatic and mammary tumors: comparison of mRNA analysis with receptor autoradiography. *Int J Cancer* 70:530–537
193. Jais P, Terris B, Ruszniewski P, LeRomancer M, Reyl-Desmars F, Vissuzaine C, Cadiot G, Mignon M, Lewin MJ 1997 Somatostatin receptor subtype gene expression in human endocrine gastroentero-pancreatic tumours. *Eur J Clin Invest* 27:639–644
194. Wulbrand U, Wied M, Zofel P, Goke B, Arnold R, Fehmann H 1998 Growth factor receptor expression in human gastroentero-pancreatic neuroendocrine tumours. *Eur J Clin Invest* 28:1038–1049
195. Mato E, Matias-Guiu X, Chico A, Webb SM, Cabezas R, Berna L, De Leiva A 1998 Somatostatin and somatostatin receptor subtype gene expression in medullary thyroid carcinoma. *J Clin Endocrinol Metab* 83:2417–2420
196. Laws SA, Gough AC, Evans AA, Bains MA, Primrose JN 1997 Somatostatin receptor subtype mRNA expression in human colorectal cancer and normal colonic mucosae. *Br J Cancer* 75:360–366
197. Nilsson O, Kolby L, Wangberg B, Wigander A, Billig H, William-Olsson L, Fjalling M, Forsell-Aronsson E, Ahlman H 1998 Comparative studies on the expression of somatostatin receptor subtypes, outcome of octreotide scintigraphy and response to octreotide treatment in patients with carcinoid tumours. *Br J Cancer* 77:632–637
198. Evans AA, Crook T, Laws SA, Gough AC, Royle GT, Primrose JN 1997 Analysis of somatostatin receptor subtype mRNA expression in human breast cancer. *Br J Cancer* 75:798–803
199. Dutour A, Kumar U, Panetta R, Ouafik L, Fina F, Sasi R, Patel YC 1998 Expression of somatostatin receptor subtypes in human brain tumors. *Int J Cancer* 76:620–627