

Towards improving the safety and diagnostic yield of stereotactic biopsy in a single centre

Ruben Dammers · Joost W. Schouten ·
Iain K. Haitsma · Arnaud J. P. E. Vincent ·
Johan M. Kros · Clemens M. F. Dirven

Received: 8 March 2010 / Accepted: 13 July 2010 / Published online: 1 August 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract

Background Previously, we reported on our single centre results regarding the diagnostic yield of stereotactic needle biopsies of brain lesions. The yield then (1996–2006) was 89.4%. In the present study, we review and evaluate our experience with intraoperative frozen-section histopathologic diagnosis on-demand in order to improve the diagnostic yield.

Methods One hundred sixty-four consecutive frameless biopsy procedures in 160 patients (group 1, 2006–2010) were compared with the historic control group (group 2, $n=164$ frameless biopsy procedures). Diagnostic yield, as well as demographics, morbidity and mortality, was compared. Statistical analysis was performed by Student's t , Mann–Whitney U , Chi-square test and backward logistic regression when appropriate.

Results Demographics were comparable. In group 1, a non-diagnostic tissue specimen was obtained in 1.8%, compared to 11.0% in group 2 ($p=0.001$). Also, both the operating time and the number of biopsies needed were decreased significantly. Procedure-related mortality decreased from 3.7% to 0.6% ($p=0.121$). Multivariate analysis only proved operating time (odds ratio (OR), 1.012; 95% confidence interval (CI), 1.000–1.025; $p=0.043$), a right-sided lesion

(OR, 3.183; 95% CI, 1.217–8.322; $p=0.018$) and on-demand intraoperative histology (OR, 0.175; 95% CI, 0.050–0.618; $p=0.007$) important factors predicting non-diagnostic biopsies.

Conclusions The importance of a reliable pathological diagnosis as obtained by biopsy must not be underestimated. We believe that when performing stereotactic biopsy for intracranial lesions, next to minimising morbidity, one should strive for as high a positive yield as possible. In the present single centre retrospective series, we have shown that using a standardised procedure and careful on-demand intraoperative frozen-section analysis can improve the diagnostic yield of stereotactic brain biopsy procedures as compared to a historical series.

Keywords Cerebral biopsy · Diagnostic yield · On-demand histopathology

Introduction

In an earlier report, we demonstrated our single centre results with both frame-based and frameless neuronavigational image-guided techniques in the context of needle biopsies of brain lesions [7]. We reported no differences in diagnostic yield, complications and biopsy-related mortality between both techniques, as corroborated by others [36, 40, 41].

On the matter of diagnostic yield, we discussed the possible importance of intraoperative frozen-section or cytological smear preparations. Our previous results, when no intraoperative histological diagnosis was readily available, showed an overall diagnostic yield of 89.4%, which is on the lower spectrum of yield reported by others [1, 9, 11, 12, 17, 19, 34, 36, 40]. We believe that neurosurgeons should strive for as high a diagnostic yield as possible, respecting the possible morbidity or even mortality,

Financial disclosures and conflicts of interest The authors have no financial interest in any materials or equipment used in the study.

R. Dammers (✉) · J. W. Schouten · I. K. Haitsma ·
A. J. P. E. Vincent · C. M. F. Dirven
Department of Neurosurgery, Erasmus MC,
's Gravendijkwal 230, P.O. Box 2040, 3000 CA Rotterdam,
The Netherlands
e-mail: r.dammers@erasmusmc.nl

J. M. Kros
Department of Pathology, Erasmus MC,
Rotterdam, The Netherlands

although small, the patient confronts when undergoing an intracranial biopsy procedure. Since several authors have shown the benefit of intraoperative histopathologic diagnosis and/or the use of modern intraoperative imaging techniques, improving the diagnostic yield up to as high as 99.2% [4, 14, 20, 22, 31, 32], we implemented intraoperative histopathologic diagnosis into our standardised biopsy procedure in order to possibly improve on our results.

In the present study, we reviewed and evaluated our experience with the use of intraoperative frozen-section diagnosis on-demand, when the neurosurgeon performing the biopsy on macroscopic observation was hesitant about the pathologic nature of the tissue sample.

Patients and methods

We reviewed a consecutive series of patients who had undergone an image-guided stereotactic intracranial biopsy procedure at the Erasmus MC from November 2006 until January 2010. This series of patients succeeded the series of 465 uninterrupted cerebral biopsies, of which 164 were frameless biopsy procedures, published in our earlier report [7]. The data were prospectively gathered, but reviewed in a retrospective manner. One hundred sixty-four frameless biopsy procedures were performed in 160 patients.

Patient and pre-operative neuroimaging characteristics

Age, gender, post-operative biopsy-related complications (morbidity) and mortality, duration of hospital stay as well as operating time, biopsy method used, number of biopsies and complications were extracted from case notes and operative reports. The site of the targeted lesion was obtained from pre-operative computed tomographic or magnetic resonance (MR) imaging. The eventual diagnosis was obtained from the pathological report, from which the diagnostic yield was determined.

Operative technique

The techniques of frameless image-guided stereotactic biopsies applied in our centre have been described previously [7]. One hundred sixty-four biopsies were performed using the frameless stereotaxy protocol under general anaesthesia and head-fixation in a three-point Mayfield clamp. Intraoperative image guidance and the surgical plan were obtained using the Stealth Treon™ system (Medtronic Inc., Minneapolis, MN, USA). Tissue samples approximately 8-mm-long and 1-mm-thick were thus obtained.

In general, four biopsies were obtained at the pre-operatively suggested target, as well as two to four more

biopsies at a site proximal to the target on the same biopsy trajectory. The macroscopic aspect of the biopsy was assessed, and it was considered pathological when one or more of the following was observed: (1) glazy/oedematous aspect, (2) haemorrhagic component and (3) dark-greyish aspect. A brain tissue sample was regarded normal when clearly white matter was observed. Intraoperative freeze sectioning was performed, only then when the surgeon was uncertain about the pathologic nature of the tissue sample, i.e. the macroscopic appearance of the acquired tissue resembled normal brain. If diagnostic tissue was not obtained, all parameters and measurements were rechecked to ensure accurate localisation, after which, a repeat biopsy sample could then be sent, via the same or a new trajectory, if need be.

Statistical analysis

Statistical analysis was performed with SPSS 15.0 for Windows (SPSS Inc., Chicago, IL, USA). The present 164 biopsies, with the abovementioned standardised procedure and intraoperative histopathologic diagnosis on-demand (group 1), were compared to the historic control of 164 frameless biopsies that were reported previously and did not include intraoperative diagnosis (group 2). Continuous data were compared using the Student's *t* test for parametric data. For non-parametric data, the Mann–Whitney *U* test was used. These are presented as mean values±standard deviations, unless otherwise specified. Proportions were compared with Chi-square and Fisher's exact test and presented as percentages. *p* values <0.05 were considered statistically significant. In order to analyse factors relating to the risk of a non-diagnostic biopsy, the variables selected were dichotomised and initially explored univariately as described above. Variables for which there was not a high probability of association (*p*>0.10) were excluded from the final regression model. Backward stepwise logistic regression was used for this purpose, and odds ratios (OR) and 95% confidence intervals (CI) are presented.

Results

In Table 1, patient and peri-operative characteristics of stereotactic biopsies in patients of group 1 and 2 are shown. There were no differences in age and gender distribution. The peri-operative complication rate was also similar. Interestingly, the number of biopsies taken and the overall operating time both decreased significantly (*p*<0.001 and *p*<0.001, respectively) when comparing with the historical group 2.

The location of the lesion intended for biopsy was not significantly different when compared to side and anatomical site. Left- and right-sided lesions were equally divided (group 1 vs. group 2, 39.0% and 36.0% vs. 39.6% and

Table 1 Patient and peri-operative characteristics of groups 1 and 2

	Group 1 (n=164)	Group 2 (n=164)	p value
Male (%)	55.5	56.7	
Age (years±SD)	52.6±18.4	55.0±17.2	
Operating time (min±SD)	108.5±34.2	127.1±32.8	<0.001
Number of biopsies±SD	7.4±3.3	9.0±3.8	<0.001
Complication (%)			
Haemorrhage	3.0	3.0	
Technical failure	2.4	3.7	

Shown are percentages and mean values±standard deviations (SD). Group 1: period November 2006 until April 2009, with intraoperative histopathological diagnosis on-demand; group 2: period August 1996 until October 2006, without the possibility of intraoperative histopathological diagnosis

40.9%, respectively). In patients of group 1, the three most often biopsied tumours were frontal (22.5%), parietal (22.0%) and thalamic (14.0%) lesions.

Post-operative complications occurred in 14 (8.5%) and 19 (11.6%) of operations in groups 1 and 2, respectively. Of these, four symptomatic haemorrhages in each group (2.4%) were diagnosed by post-operative computed tomography. There was no statistically significant difference between groups 1 and 2 in the frequency of occurrence of post-operative complications. In the most recent patients (group 1), there was one death (0.6%) attributable to the surgical procedure, i.e. death occurring within 30 days as a result of symptomatic post-operative haemorrhage or oedema. There were six biopsy-related deaths (3.7%) in group 2, which did not reach statistical significance ($p=0.121$, Fisher's exact).

Diagnostic yield

The overall diagnostic yield of group 1 was 98.2%, i.e. a histological diagnosis was established in 161 of 164 biopsy procedures, whereas in three (1.8%), the biopsy was non-diagnostic. In one patient, this resulted in a repeated biopsy procedure that resulted in the diagnosis of a lymphoma. The second and third patient underwent a subsequent craniotomy when a glioblastoma and metastasis of lung carcinoma was diagnosed, respectively. In 15 cases (8.9%), the surgeon judged the tissue obtained to resemble normal brain and therefore requested for an intraoperative frozen-section diagnosis by the neuropathologist. When non-diagnostic, this prompted for another tissue sample in the chosen trajectory. In the period from August 1996 until October 2006, i.e. group 2, when there was no availability of intraoperative histopathologic diagnosis, the diagnostic

yield was only 89.0%, which is statistically lower compared with group 1 diagnostic yield ($p=0.001$). Table 2 shows the histological diagnoses that were made on the tissue samples taken in both groups. On average, 7.4 (range, 1–19) and 9.0 (1–24) tissue samples were taken during the biopsy procedure in groups 1 and 2, respectively ($p<0.001$, Table 1).

Table 3 shows the results of the univariate analysis performed to establish the factors related to obtaining a non-diagnostic biopsy specimen ($n=21$). In the subsequent multivariate regression analysis, only operating time (OR, 1.012; 95% CI, 1.000–1.025; $p=0.043$), a right-sided lesion (OR, 3.183; 95% CI, 1.217–8.322; $p=0.018$) and the use of on-demand intraoperative histopathologic examination (OR, 0.175; 95% CI, 0.050–0.618; $p=0.007$) showed to be important factors.

Discussion

Previously, we have shown the diagnostic yield of stereotactic biopsy either by frame-based or frameless techniques to be equal, as were the morbidity and mortality related to the procedure [7]. The overall diagnostic yield in the previous report was 89.4%, which might be regarded lying on the lower spectrum of the yield obtained by others [1, 9, 11, 12, 17, 19, 34, 36, 40]. The present study suggests that by using on-demand frozen-section analysis of tissue specimens begotten by biopsy, the diagnostic yield can be dramatically improved (98.2%).

Several other authors have corroborated these results, i.e. performing intraoperative frozen-section or cytological smear preparations will increase the rate on a reliable histopathological diagnosis up to as high as 99.2% [4, 14, 17, 29, 31, 38]. McDermott and Bernstein, in *Neuro-oncology: The Essentials* (p. 115), actually state '[...] The management of the failed biopsy begins in the operating room where intraoperative examination of tissue is mandatory to ensure that diagnostic tissue is in fact obtained [...]'. Nevertheless, the five most recent reports, illustrating diagnostic yields of 99.3% [35], 94% [30], 90.6% [5], 89.8% [1] and 83.6% [37], do not mention intraoperative pathologic confirmation. Also, in our series, we were not able to perform intraoperative cytological studies in all biopsy procedures. This was primarily due to logistic considerations, such as pressure on the operating theatre regarding time and the unavailability of adequate neuropathologic support on all biopsies. Therefore, we turned to on-demand frozen-section analysis, i.e. the surgeon performing the biopsy assessed the tissue specimen macroscopically as normal brain or pathological tissue sample and, when uncertain, requested the neuropathologist for intraoperative confirmation. Apparently, concluded from the results shown in the present study, there is no need for

Table 2 Histological diagnoses made on tissue samples acquired by stereotactic biopsy

Diagnosis (%)	Group 1 (n=164)	Group 2 (n=164)	<i>p</i> value
Non-diagnostic	1.8	11.0	0.001
Malignant glioma	53.0	51.3	
Low-grade glioma	12.2	14.6	
Lymphoma	14.0	7.9	
Metastasis	3.0	7.3	
Infection/vasculitis/ MS/abscess	12.2	6.7	
Craniopharyngioma cysts	1.8	1.2	
Others ^a	1.8	0	

Shown are percentages. Group 1: period November 2006 until April 2009, with intraoperative histopathological diagnosis on-demand; group 2: period August 1996 until October 2006, without the possibility of intraoperative histopathological diagnosis. Malignant glioma includes astrocytoma, oligoastrocytoma, oligodendroglioma WHO grade III and glioblastoma (WHO grade IV). Low-grade glioma includes astrocytoma, oligoastrocytoma and oligodendroglioma WHO grade II

MS multiple sclerosis

^a Include meningioma, medulloblastoma, teratoma, malignant peripheral nerve sheath tumour and old haematoma

an intraoperative cytological preparation, and therefore, a neuropathologist standby for every stereotactic cerebral biopsy procedure when an experienced surgeon evaluates the tissue samples obtained. These results are corroborated by the recent article of Shooman et al. who on top of this state that routine intraoperative neuropathological examination is not needed at all [35].

Table 4 shows the diagnostic yield, complication rates, deaths and methods used to ensure a greater diagnostic yield in the most recent biopsy series. Since 2001, a total of 12,038 biopsy procedures are collected in the literature, with an overall diagnostic yield ranging from 83.6% to 100%. Positron emission tomography imaging, MR perfusion and MR imaging spectroscopy integrated into the planning for stereotactic biopsy procedures to determine the trajectory and selection of the appropriate target promise to be important adjuncts to improve diagnostic yield and to decrease sampling error, especially when low-grade gliomas are involved [6, 13, 15, 24, 27, 28, 32, 33]. Recent studies show a yield of 100%, although patient numbers are relatively small [6, 13, 15, 24, 32, 33]. Additional use of metabolic imaging data acquired with MR spectroscopy, more specifically, the ratio of *N*-acetyl aspartate and choline-containing compounds, for target selection can potentially increase the diagnostic efficacy of biopsy procedures.

Another issue to be addressed is the diagnostic accuracy, i.e. the rate of sampling error as established by subsequent open craniotomy. In literature, accuracy rates ranging from 73% to 97% have been reported [2, 7, 17, 30, 40]. Chernov et al. compared the diagnostic efficacy of stereotactic brain biopsy performed with and without additional use of spectroscopic imaging and found that spectroscopy was superior, although not significantly due to the small number of patients, in reaching a diagnosis at all (100% vs. 90%) [6]. Diagnostic accuracy, however, could not be improved by using spectroscopy imaging (67% vs. 79%, MR imaging with and without spectroscopy, respectively).

In the present study, an important difference with the historical control was the use of on-demand intraoperative histological confirmation of pathological tissue, which was

Table 3 Results of univariate analysis to establish confounding factors related to obtaining a non-diagnostic biopsy specimen

	Diagnostic (n=305)	Non-diagnostic (n=21)	<i>p</i> value
Male (%)	55.7	66.6	0.371
Age (years±SD)	53.9±17.7	5.8±18.2	0.828
Tumour side			
Right	38.7	66.7	0.019
Left	38.7	19.0	0.101
Tumour location (frontal, parietal, temporal, etc.)			0.388
Surgeon			0.421
Operating time (min±SD)	116.0±33.4	138.1±39.6	0.004
Number of biopsies±SD	8.3±3.7	7.3±2.7	0.259
On-demand intraoperative frozen-section histology	52.8	14.3	0.001
Peri-operative complication	6.3	4.8	1.000
Post-operative complication	10.5	4.8	0.708
Biopsy-related death	2.3	0	1.000

Shown are percentages and mean values±standard deviations (SD)

confirmed by multivariate analysis to be an important factor to decrease the chance on a non-diagnostic tissue specimen. However, due to the retrospective nature of the present study, some confounding factors should be put forward that may influence the overall results. As a result of technological advance and ease of use, the frameless stereotactic technique has been used more frequently in recent times at our neurosurgical clinic (97.6% vs. 35.3%, respectively). Although we, in concordance with others, have previously shown that this does not influence diagnostic yield [7, 36, 40, 41], Dorward et al. found disagreeing results, concluding that frameless techniques are superior to the gold standard of frame-based biopsy [8].

To exclude this possible confounding factor, we compared the most recent frameless biopsy series (group 1) with the historical control of frameless biopsies only. Furthermore, the effect of an initial learning curve cannot be ruled out. Another, more important factor might be the number of surgeons performing the biopsy procedures. These differ significantly between the historical and the present group of patients. In group 2, a total of 20 surgeons, whereas in group 1, only 15 surgeons were involved with the stereotactic biopsy procedures. In the most recent group (group 1), two surgeons (RD and JWS) performed 51.8% of surgeries and supervised another 12.8%. One might argue that due to a surgeon's increased experience, he/she is apt to

Table 4 Diagnostic yield in recent biopsy series

Author (year)	<i>n</i>	Yield	Method to improve yield	Morbidity	Mortality
Kaakaji et al. (2001) [16]	269	97.1	None	3.0	0.4
Paleologos et al. (2001) [31]	125	97.6	Intraoperative histology	2.4	0.8
Ulm et al. (2001) [39]	200	98.5	None	2.0	0
Bhardwaj et al. (2002) [3]	76	98.7	None	2.6	0
Dorward et al. (2002) [8]	155	96.8	Intraoperative histology	16.1	3.2
Gralla et al. (2003) [10]	57	98	Intraoperative histology	3.5	0
Kim et al. (2003) [17]	308	91.7	Intraoperative histology	3.9	0.6
Maia et al. (2004) [24]	21	100	MR perfusion	NA	NA
Pirotte et al. (2004) [32]	32	100	Spectroscopy	NA	NA
Aker et al. (2005) [2]	130	94	Intraoperative histology	0.8	0
Grossman et al. (2005) [11]	355	93.8	None	3.6	0.6
Hemm et al. (2005) [13]	10	100	Spectroscopy	NA	NA
Heper et al. (2005) [14]	130	99.2	Intraoperative histology	0.7	0
McGirt et al. (2005) [26]	270	93	None	5.0	1.0
Smith et al. (2005) [36]	213	90	None	4.0	0.4
Tilgner et al. (2005) [38]	5,000	90.3	Intraoperative histology	2.7	0.7
Ferreira et al. (2006) [9]	170	92	None	2.9	1.2
Shastri-Hurst et al. (2006) [34]	207	89.3	Intraoperative histology	6.4	1.9
Woodworth et al. (2006) [41]	270	90	None	4.0	1.0
Pirotte et al. (2007) [33]	20	100	Spectroscopy	10.0	0
Dammers et al. (2008) [7]	465	89.5	None	11.7	3.9
Hermann et al. (2008) [15]	9	100	Spectroscopy	NA	NA
Kongkham et al. (2008) [18]	622	98.4	Intraoperative histology	6.9	1.3
Lunsford et al. (2008) [23]	1,664	NA	None	3.1	0.1
Landriel et al. (2008) [21]	192	90.6	Intraoperative histology	3.6	2.1
Air et al. (2009) [1]	284	89.8	None	6.7	2.0
Chernov et al. (2009) [6]	69	94	Spectroscopy (100%)	1.4	0
Owen and Linskey (2009) [30]	106	94	None	3.8	0
Teixeira et al. (2009) [37]	176	83.6	None	6.4	0.6
Chen et al. (2009) [5]	299	90.6	None	7.4	1.3
Shooman et al. (2010) [35]	134	99.3	None	2.2	1.5
Total	12,038	94.7 (83.6–100)		4.7 (0.7–16.1)	0.9 (0–3.9)
Present series	164	98.2	On-demand intraoperative histology	8.5	0.6

Diagnostic yield, morbidity and mortality are expressed in percentages

prepare a biopsy trajectory and target that ensures less chance on complications and a higher chance on a positive histological diagnosis. For this, there is no evidence in recent literature as far as we are aware, although Shastri-Hurst et al. touched upon the subject in the “Results” section and found no evidence that the individual performing the biopsy most frequently had better results in terms of complication rate or non-diagnostic rate [34]. Lastly, the recent 164 cases were all histologically examined by one experienced dedicated neuropathologist (JMK), whereas in the past, three pathologists performed the histopathologic diagnosis. A single dedicated neuropathologist might be able to provide a diagnosis more often. Furthermore, the changing tumour classification and grading by the World Health Organisation might aid to construct a more proficient diagnosis nowadays.

The possible advantages of on-demand intraoperative frozen-section analysis are its cost-effectiveness concerning the consultation of a neuropathologist and the need for repeat surgery. More importantly, a non-diagnostic biopsy has an imminent effect on a patient's further treatment. On-demand histopathology, however, demands experienced and dedicated neuro-oncologic trained neurosurgeons [35]. Nevertheless, the issue of diagnostic accuracy, i.e. biopsy sampling error, remains unresolved. As described above, newly available diagnostic techniques might add to locate the adequate biopsy site and decrease sampling error.

Although tumour verification might be warranted, the treating physician should always also consider the morbidity and mortality related to the biopsy procedure. Of the 12,038 procedures in recent literature (Table 4), the average procedure-related morbidity was 4.7% (range, 0.7–16.1%), compared to our own recent series (8.5%). These vary from minor complications to devastating neurological sequelae that can influence a patients' independence. The average mortality rate was 0.9% (range, 0–3.9%) and 0.6% in recent literature and the present series, respectively, from which one might conclude that a brain biopsy is a safe procedure.

Concluding, the importance of a reliable pathological diagnosis as obtained by biopsy, notwithstanding the possibility of sampling error [25, 28], must not be underestimated. Both with regard to the possible risks for the patients, such as (increase of) neurological deficits by oedema or haemorrhage and even death, as well as with regard to the further treatment that might be warranted for each individual patient. In the present single centre retrospective series, we have shown that using a standardised procedure and careful on-demand intraoperative frozen-section analysis, when the macroscopic aspects is not clearly pathological, can improve the diagnostic yield of stereotactic brain biopsy procedures as compared to a historical series.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Air EL, Leach JL, Warnick RE, McPherson CM (2009) Comparing the risks of frameless stereotactic biopsy in eloquent and noneloquent regions of the brain: a retrospective review of 284 cases. *J Neurosurg* 111:820–824
2. Aker FV, Hakan T, Karadereler S, Erkan M (2005) Accuracy and diagnostic yield of stereotactic biopsy in the diagnosis of brain masses: comparison of results of biopsy and resected surgical specimens. *Neuropathology* 25:207–213
3. Bhardwaj RD, Bernstein M (2002) Prospective feasibility study of outpatient stereotactic brain lesion biopsy. *Neurosurgery* 51:358–361
4. Brainard JA, Prayson RA, Barnett GH (1997) Frozen section evaluation of stereotactic brain biopsies: diagnostic yield at the stereotactic target position in 188 cases. *Arch Pathol Lab Med* 121:481–484
5. Chen CC, Hsu PW, Erich Wu TW, Lee ST, Chang CN, Wei KC, Chuang CC, Wu CT, Lui TN, Hsu YH, Lin TK, Lee SC, Huang YC (2009) Stereotactic brain biopsy: single center retrospective analysis of complications. *Clin Neurol Neurosurg* 111:835–839
6. Chernov MF, Muragaki Y, Ochiai T, Taira T, Ono Y, Usukura M, Maruyama T, Nakaya K, Nakamura R, Iseki H, Kubo O, Hori T, Takakura K (2009) Spectroscopy-supported frame-based image-guided stereotactic biopsy of parenchymal brain lesions: comparative evaluation of diagnostic yield and diagnostic accuracy. *Clin Neurol Neurosurg* 111:527–535
7. Dammers R, Haitsma IK, Schouten JW, Kros JM, Avezaat CJ, Vincent AJ (2008) Safety and efficacy of frameless and frame-based intracranial biopsy techniques. *Acta Neurochir (Wien)* 150:23–29
8. Dorward NL, Paleologos TS, Alberti O, Thomas DG (2002) The advantages of frameless stereotactic biopsy over frame-based biopsy. *Br J Neurosurg* 16:110–118
9. Ferreira MP, Ferreira NP, Pereira Filho Ade A, Pereira Filho Gde A, Franciscatto AC (2006) Stereotactic computed tomography-guided brain biopsy: diagnostic yield based on a series of 170 patients. *Surg Neurol* 65(S1):27–32
10. Gralla J, Nimsky C, Buchfelder M, Fahlbusch R, Ganslandt O (2003) Frameless stereotactic brain biopsy procedures using the Stealth Station: indications, accuracy and results. *Zentralbl Neurochir* 64:166–170
11. Grossman R, Sadetzki S, Spiegelmann R, Ram Z (2005) Haemorrhagic complications and the incidence of asymptomatic bleeding associated with stereotactic brain biopsies. *Acta Neurochir (Wien)* 147:627–631
12. Hall WA (1998) The safety and efficacy of stereotactic biopsy for intracranial lesions. *Cancer* 82:1749–1755
13. Hemm S, Rigau V, Chevalier J, Picot MC, Bauchet L, El Fertit H, Rodriguez MA, Cif L, Vayssiere N, Zanca M, Baldet P, Segnarbieux F, Coubes P (2005) Stereotactic coregistration of 201Tl SPECT and MRI applied to brain tumor biopsies. *J Nucl Med* 46:1151–1157
14. Heper AO, Erden E, Savas A, Ceyhan K, Erden I, Akyar S, Kanpolat Y (2005) An analysis of stereotactic biopsy of brain tumors and nonneoplastic lesions: a prospective clinicopathologic study. *Surg Neurol* 64(Suppl 2):S82–S88
15. Hermann EJ, Hattingen E, Krauss JK, Marquardt G, Pilatus U, Franz K, Setzer M, Gasser T, Tews DS, Zanella FE, Seifert V, Lanfermann H (2008) Stereotactic biopsy in gliomas guided by 3-

- tesla 1H-chemical-shift imaging of choline. *Stereotact Funct Neurosurg* 86:300–307
16. Kaakaji W, Barnett GH, Bernhard D, Warbel A, Valaitis K, Stamp S (2001) Clinical and economic consequences of early discharge of patients following supratentorial stereotactic brain biopsy. *J Neurosurg* 94:892–898
 17. Kim JE, Kim DG, Paek SH, Jung HW (2003) Stereotactic biopsy for intracranial lesions: reliability and its impact on the planning of treatment. *Acta Neurochir (Wien)* 145:547–554
 18. Kongkham PN, Knifed E, Tamber MS, Bernstein M (2008) Complications in 622 cases of frame-based stereotactic biopsy, a decreasing procedure. *Can J Neurol Sci* 35:79–84
 19. Krieger MD, Chandrasoma PT, Zee CS, Apuzzo ML (1998) Role of stereotactic biopsy in the diagnosis and management of brain tumors. *Semin Surg Oncol* 14:13–25
 20. Kuhn SA, Romeike B, Walter J, Kalff R, Reichart R (2009) Multiplanar MRI-CT fusion neuronavigation-guided serial stereotactic biopsy of human brain tumors: proof of a strong correlation between tumor imaging and histopathology by a new technical approach. *J Cancer Res Clin Oncol* 135:1293–1302
 21. Landriell F, Hem S, Hasdeu S, Kitroser M, Peña L, Ajler P, Ciraolo C, Knezevich F (2008) Biopsia estereotáctica cerebral guiada tomográficamente análisis de una serie de 192 casos. *Rev Argent Neuroc* 22:110–113
 22. Levivier M, Goldman S, Pirotte B, Brucher JM, Baleriaux D, Luxen A, Hildebrand J, Brotchi J (1995) Diagnostic yield of stereotactic brain biopsy guided by positron emission tomography with [¹⁸F]fluorodeoxyglucose. *J Neurosurg* 82:445–452
 23. Lunsford LD, Niranjan A, Khan AA, Kondziolka D (2008) Establishing a benchmark for complications using frame-based stereotactic surgery. *Stereotact Funct Neurosurg* 86:278–287
 24. Maia AC Jr, Malheiros SM, da Rocha AJ, Stavale JN, Guimaraes IF, Borges LR, Santos AJ, da Silva CJ, de Melo JG, Lanzoni OP, Gabbai AA, Ferraz FA (2004) Stereotactic biopsy guidance in adults with supratentorial nonenhancing gliomas: role of perfusion-weighted magnetic resonance imaging. *J Neurosurg* 101:970–976
 25. McDermott MW, Bernstein M (2008) Image-guided surgery. In: Bernstein M, Berger MS (eds) *Neuro-oncology: the essentials*. Thieme Medical Publishers, Inc, New York, pp 112–125
 26. McGirt MJ, Woodworth GF, Coon AL, Frazier JM, Amundson E, Garonzik I, Olivi A, Weingart JD (2005) Independent predictors of morbidity after image-guided stereotactic brain biopsy: a risk assessment of 270 cases. *J Neurosurg* 102:897–901
 27. Mosskin M, Ericson K, Hindmarsh T, von Holst H, Collins VP, Bergstrom M, Eriksson L, Johnstrom P (1989) Positron emission tomography compared with magnetic resonance imaging and computed tomography in supratentorial gliomas using multiple stereotactic biopsies as reference. *Acta Radiol* 30:225–232
 28. Muragaki Y, Chernov M, Maruyama T, Ochiai T, Taira T, Kubo O, Nakamura R, Iseki H, Hori T, Takakura K (2008) Low-grade glioma on stereotactic biopsy: how often is the diagnosis accurate? *Minim Invasive Neurosurg* 51:275–279
 29. O'Neill KS, Dyer PV, Bell BA, Wilkins PR, Uttley D, Marsh HT (1992) Is peroperative smear cytology necessary for CT-guided stereotactic biopsy? *Br J Neurosurg* 6:421–427
 30. Owen CM, Linskey ME (2009) Frame-based stereotaxy in a frameless era: current capabilities, relative role, and the positive- and negative predictive values of blood through the needle. *J Neurooncol* 93:139–149
 31. Paleologos TS, Dorward NL, Wadley JP, Thomas DG (2001) Clinical validation of true frameless stereotactic biopsy: analysis of the first 125 consecutive cases. *Neurosurgery* 49:830–835
 32. Pirotte B, Goldman S, Massager N, David P, Wikler D, Vandesteene A, Salmon I, Brotchi J, Levivier M (2004) Comparison of 18F-FDG and 11C-methionine for PET-guided stereotactic brain biopsy of gliomas. *J Nucl Med* 45:1293–1298
 33. Pirotte BJ, Lubansu A, Massager N, Wikler D, Goldman S, Levivier M (2007) Results of positron emission tomography guidance and reassessment of the utility of and indications for stereotactic biopsy in children with infiltrative brainstem tumors. *J Neurosurg* 107:392–399
 34. Shastri-Hurst N, Tsegaye M, Robson DK, Lowe JS, Macarthur DC (2006) Stereotactic brain biopsy: an audit of sampling reliability in a clinical case series. *Br J Neurosurg* 20:222–226
 35. Shooman D, Belli A, Grundy PL (2010) Image-guided frameless stereotactic biopsy without intraoperative neuropathological examination. *J Neurosurg*. doi:10.3171/2009.12.JNS09573
 36. Smith JS, Quinones-Hinojosa A, Barbaro NM, McDermott MW (2005) Frame-based stereotactic biopsy remains an important diagnostic tool with distinct advantages over frameless stereotactic biopsy. *J Neurooncol* 73:173–179
 37. Teixeira MJ, Fonoff ET, Mandel M, Alves HL, Rosemberg S (2009) Stereotactic biopsies of brain lesions. *Arq Neuropsiquiatr* 67:74–77
 38. Tilgner J, Herr M, Ostertag C, Volk B (2005) Validation of intraoperative diagnoses using smear preparations from stereotactic brain biopsies: intraoperative versus final diagnosis—influence of clinical factors. *Neurosurgery* 56:257–265
 39. Ulm AJ, Bova FJ, Friedman WA (2001) Stereotactic biopsy aided by a computer graphics workstation: experience with 200 consecutive cases. *Surg Neurol* 56:366–371
 40. Woodworth G, McGirt MJ, Samdani A, Garonzik I, Olivi A, Weingart JD (2005) Accuracy of frameless and frame-based image-guided stereotactic brain biopsy in the diagnosis of glioma: comparison of biopsy and open resection specimen. *Neurol Res* 27:358–362
 41. Woodworth GF, McGirt MJ, Samdani A, Garonzik I, Olivi A, Weingart JD (2006) Frameless image-guided stereotactic brain biopsy procedure: diagnostic yield, surgical morbidity, and comparison with the frame-based technique. *J Neurosurg* 104:233–237

Comment

The colleagues of Rotterdam compared the diagnostic yield of frameless stereotactic biopsies with or without the possibility of frozen section and smear preparation assessment during biopsy. The option of asking a neuropathologist to the OR clearly decreased non-diagnostic samples in the final neuropathological evaluation: 1.8% (3 of 164 frameless biopsies 2006–2010) vs. 11% (18 of 160 frameless biopsies 1996–2006). The average numbers of samples were 7.4 vs. 9.0 per patient. Neurosurgeons assessed the macroscopic appearance of the samples as pathologic (oedematous, hemorrhagic, or dark-grayish) - or as normal brain, the only indication for the intraoperative freeze-sectioning.

The risk of 11% of repeat biopsy and delayed therapy would require reduction – but how?

1. Instant and routine microscopic verification of the presence of representative samples with a putative first diagnosis in the OR – in real life? Neurosurgical units are served by a few neuropathologists at best – often by one or two. Are they willing to walk from their microscopy desks with piles of slide trays to the OR in the other end of the building or in the next one? Few units really have a neuropathologist sitting in the OR until the verification is done.

2. On-demand microscopic assessment in the OR of dubious samples only – as cleverly arranged above.

3. Transfer of fresh samples to the neuropathology unit – and waiting for tens of minutes a phone call from a neuropathologist.

Juha E Jääskeläinen
Kuopio Finland