

RESEARCH ARTICLE

Association of Cardiovascular Risk Factors with Carotid Intima Media Thickness in Patients with Rheumatoid Arthritis with Low Disease Activity Compared to Controls: A Cross-Sectional Study

Deborah F. van Breukelen—van der Stoep¹, Derkjen van Zeben¹, Boudewijn Klop², Gert-Jan M. van de Geijn³, Hans J. W. Janssen³, Mieke J. M. W. Hazes⁴, Erwin Birnie^{5,6}, Noelle van der Meulen², Marijke A. De Vries², Manuel Castro Cabezas^{2*}

1 Department of Rheumatology, Sint Franciscus Gasthuis, Rotterdam, The Netherlands, **2** Department of Internal Medicine, Sint Franciscus Gasthuis, Rotterdam, The Netherlands, **3** Department of Clinical Chemistry, Sint Franciscus Gasthuis, Rotterdam, The Netherlands, **4** Department of Rheumatology, Erasmus Medical Centre, Rotterdam, The Netherlands, **5** Institute of Health Policy and Management, Erasmus University, Rotterdam, The Netherlands, **6** Statistics and Education, St. Franciscus Gasthuis, Rotterdam, The Netherlands

* m.castrocabezas@sfgv.nl



CrossMark
click for updates

OPEN ACCESS

Citation: van Breukelen—van der Stoep DF, van Zeben D, Klop B, van de Geijn G-JM, Janssen HJW, Hazes MJMW, et al. (2015) Association of Cardiovascular Risk Factors with Carotid Intima Media Thickness in Patients with Rheumatoid Arthritis with Low Disease Activity Compared to Controls: A Cross-Sectional Study. PLoS ONE 10 (10): e0140844. doi:10.1371/journal.pone.0140844

Editor: Eliseo A Eugenin, Rutgers University, UNITED STATES

Received: June 1, 2015

Accepted: October 1, 2015

Published: October 20, 2015

Copyright: © 2015 van Breukelen—van der Stoep et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are available from Dryad (doi:10.5061/dryad.hh3j7).

Funding: This study consists of data from two studies. One study, the FRANCIS study, received financial support from the Sint Franciscus Research Fund and the Stichting Coolingsel Rotterdam. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Objectives

Rheumatoid arthritis (RA) has been identified as an independent cardiovascular risk factor. The importance of risk factors such as hypertension and hyperlipidemia in the generation of atherosclerosis in RA patients is unclear. This study analyzed clinical parameters associated with carotid intima media thickness (cIMT) in patients with RA.

Methods

Subjects with RA and healthy controls without RA, both without known cardiovascular disease, were included. Participants underwent a standard physical examination and laboratory measurements including a lipid profile. cIMT was measured semi-automatically by ultrasound.

Results

In total 243 RA patients and 117 controls were included. The median RA disease duration was 7 years (IQR 2–14 years). The median DAS28 was 2.4 (IQR 1.6–3.2) and 114 (50.4%) of the RA patients were in remission. The presence of RA and cIMT were not associated (univariate analysis). Multivariable regression analysis showed that cIMT in RA patients was associated with age ($B = 0.006$, $P < 0.001$) and systolic blood pressure ($B = 0.003$, $P = 0.003$). In controls, cIMT was associated with age ($B = 0.006$, $P < 0.001$) and smoking ($B = 0.097$, $P = 0.001$).

Competing Interests: The authors have declared that no competing interests exist.

Conclusion

cIMT values were similar between RA patients and controls. Hypertension was strongly associated with cIMT in RA patients. After adjustment, no association between cIMT and specific RA disease characteristics was found in this well treated RA cohort.

Introduction

Rheumatoid Arthritis (RA) is a chronic systemic inflammatory disease of unknown etiology, affecting ~1% of the adult general population [1, 2]. Patients with RA have a higher all cause mortality risk than the general population and the leading cause of death is cardiovascular disease (CVD) [3–5]. The evidence of the excess cardiovascular risk in rheumatoid arthritis (RA) has been well described [3, 5–9] and the presence of RA can be considered an independent risk factor for CVD [7, 10]. The prevalence of CVD in patients with RA is as high as in patients with type 2 diabetes mellitus [7, 11]. Since inflammation is a key event in the development of atherosclerosis [12–14] it has been proposed that the increased inflammatory state of patients with RA explains, at least in part, the increased cardiovascular risk [8, 10, 15–17]. Several RA specific risk factors such as disease activity, inflammatory markers and anti-CCP have been associated with an increased carotid intima media thickness (cIMT) and CVD risk. A recent study by Barbarroja et al. showed that anti-CCP antibodies act as direct inductors of the pro-oxidative status and the inflammatory and atherogenic profile of lymphocytes, monocytes and neutrophils in RA [18]. In line with this report, Vázquez et al. showed an association between anti-CCP levels and CRP levels with an increased cIMT and CVD risk [19]. Moreover, traditional risk factors such as hypertension, hyperlipidemia, smoking and overweight are also highly prevalent among RA patients [20, 21], and several studies have shown a significant underdiagnosis and undertreatment of these traditional risk factors in RA [22, 23].

The cIMT measured by ultrasound is a surrogate marker of atherosclerosis and the most widely used non invasive imaging method to assess atherosclerosis and CVD risk. A higher cIMT reflects a (pre-) atherogenic condition and is predictive for future cardiovascular events [24, 25]. Several studies have shown an increased cIMT in RA patients, even early in the course of the disease [26–30]. Traditional risk factors may also play a role besides the inflammatory state in RA. Unfortunately, the association between traditional risk factors and cIMT in RA patients is still unclear [31]. Increasing evidence suggests that a cumulative number of traditional CVD risk factors contributes to the higher CVD risk in RA [32]. The aim of this study was to investigate which factors are associated to cIMT in RA patients in comparison to controls.

Materials and Methods

Study design and subjects

A cross-sectional study was carried out in patients with RA and controls to determine the differences in the relationship between cIMT and clinical factors. The study was carried out between July 2009 and February 2013 at the Diabetes and Vascular Centre and the outpatient clinic of Rheumatology of the Sint Franciscus Gasthuis, Rotterdam, the Netherlands. All patients with RA in this report were participants in the FRANCIS study, an open label randomized clinical trial to investigate the effectiveness of strict treatment of cardiovascular risk factors in RA (The Dutch Trialregister, NTR3873; ABR no. NL32669.101.10). RA patients attending

the outpatient clinic from the Department of Rheumatology were asked to participate in the FRANCIS study. Inclusion criteria were the presence of RA and an age ≤ 70 years. Exclusion criteria were the presence of diabetes mellitus (DM) or CVD. CVD was defined as a documented history of myocardial infarction, cerebrovascular event, amputation due to peripheral artery disease, intermittent claudication, or a prior percutaneous transluminal coronary angioplasty (TCA) or coronary artery bypass graft (CABG). In addition, kidney disease defined as an MDRD < 40 was an exclusion criterium. Only patients with a CVD risk $< 10\%$ according to the 2009 version of the SCORE model were randomized. Patients aged over 65 years were classified as 65 years old in order to be able to use the SCORE model. Patients with a CVD risk $\geq 10\%$ were followed in a separate cohort ("high risk cohort"). For the current analysis, baseline data from both randomized patients and patients in the high risk cohort were used. The RA patients were treated by their own rheumatologists according to a treat-to-target principle, aiming for disease remission (DAS28 < 2.6). RA was defined by the ACR '87 criteria [33]. Unmatched control subjects were non-RA patients followed in a separate observational study (ABR no. NL29910.101.09) from our department. They were recruited from the outpatient clinic of the Diabetes and Vascular Center of the Sint Franciscus Gasthuis in Rotterdam and underwent measurements identical to the RA patients of the FRANCIS study [34]. Exclusion criteria for the control group were also the presence of DM, CVD and/or kidney disease. All controls fulfilling the age limit of ≤ 70 years were included in this analysis. In addition, RA patients and controls who used statins and/or anti-hypertensives were excluded. Controls with a CRP > 10 mg/L were excluded from the analysis. Anthropometric characteristics i.e. height, weight, waist circumference and blood pressure were obtained as well as a detailed medical history and the use of medication. All subjects provided written informed consent. The studies were approved by the independent Regional Medical Ethical Committee Rotterdam of the Maasstad Hospital, the Netherlands.

Rheumatoid arthritis disease activity

Rheumatoid arthritis disease activity was assessed by using the Disease Activity Score with 28 joints counted (DAS28). This score included swollen joint count (28), tender joint count (28), VAS (0–100 scale) indicating pain/discomfort due to RA, and the erythrocyte sedimentation rate (ESR). We also calculated disease activity with C-reactive protein (CRP) instead of ESR (DAS28CRP),

Laboratory measurements

A standardized set of measurements was performed in each subject. Laboratory parameters were determined at the Department of Clinical Chemistry, Sint Franciscus Gasthuis, according to standard procedures. Renal and liver function tests as well as glucose, CRP, total cholesterol, HDL-C and triglycerides (TG) were measured using Synchron LX-20 and DxC analyzers (Beckman Coulter, Anaheim CA, USA) (34, 35). LDL-C was calculated using the Friedewald formula if TG were below 4.60 mmol/l. Apolipoprotein (apo) AI and apo B were determined in serum by rate nephelometry using IMMAGE with commercially available kits (Beckman Coulter, Anaheim CA, USA) [31,32]. In the case of RA patients, blood samples were obtained following the strict protocol of the FRANCIS study. Samples from control subjects were drawn with less strict criteria.

Intima media thickness of the carotid arteries

Carotid ultrasound scans were carried out using the ART-LAB (Esaote, Italy) by trained and experienced sonographers as described earlier [34, 35]. Ultrasound scans were performed with

the patients lying in a supine position with the head resting comfortably and the neck slightly hyperextended and rotated in the opposite direction of the probe. The ultrasound images were obtained of the distal 1 cm of the far wall of each common carotid artery (CCA) using B-mode ultrasound producing two echogenic lines. These lines represent the combined thickness of the intima and media layers of the arterial wall. Each CCA was imaged in three different projections: CCA right side 90-120-150 and CCA left side 210-240-270 degrees. The segments were measured semi-automated in triplicate. In some patients cIMT measurements were only partly available due to technical difficulties. In these cases, the mean cIMT was only calculated if at least measurements at two different angles per side were available.

Statistics

Data are given as mean \pm standard deviation (SD) unless stated otherwise. Differences between groups were determined using the unpaired Student's t-test or Chi-square test, where appropriate. In case of skewed variables (DAS28(-CRP) and TG), the Mann-Whitney U test was used for comparisons between groups. For statistical analysis, cIMT was defined as the mean of the six individual measurements. Multivariable linear regression analysis was used to identify factors associated with cIMT in RA patients and controls. Covariables entered into the regression analysis were: age, sex, systolic blood pressure, waist circumference, smoking habit, glucose, apo AI, apo B, total cholesterol, HDL-C, LDL-C, TG, glucose, creatinin and CRP. For RA patients, the presence of rheumatoid factor and anti-CCP, erosive disease, disease duration, DAS28CRP, DAS28 and the use of methotrexate and biologicals were added to the model. Covariables showing a $P < 0.10$ in the univariate analysis were entered in the multivariable analysis. In order to create comparable models for RA patients and controls, a separate analysis with and without RA disease duration was performed for RA patients. In the case that two or more related parameters were highly correlated (multicollinearity), only one was selected for the multivariable analysis. As a result, waist circumference was selected instead of BMI; systolic blood pressure was selected instead of diastolic blood pressure; and LDL-C was chosen above apo B or TG. Significant co-variables were investigated for interaction. The multivariable analysis included age, sex, smoking, waist circumference, systolic blood pressure and LDL-C. The association between glucose and cIMT was analyzed separately because it was unknown which of the control subjects had their blood samples taken after an overnight fast. All statistical analyses were performed using PASW statistics version 18.0 (IBM SPSS Statistics, New York, United States). P-values below 0.05 (two sided) were considered statistically significant.

Results

General Characteristics

243 RA patients and 117 controls were included. General characteristics of RA patients and controls are shown in [Table 1](#). Both groups were comparable regarding age, gender, lipid profile, waist circumference, and cIMT. There were more current smokers among RA patients compared to controls. RA patients had a significantly higher body mass index (BMI) and systolic blood pressure ([Table 1](#)).

The median RA disease duration for RA patients was 7 years (IQR 2–14 years). In total, 107 patients (44.0%) used NSAIDs, 186 (76.5%) RA patients used methotrexate and 96 (39.5%) used an anti-TNF α or other biological agent. Of the 96 patients with a biological agent, 25 (26%) used no methotrexate and 71 (74%) used a combination of methotrexate and a biological.

The median DAS28 was 2.4 (IQR 1.6–3.2). In total, 114 (50.4%) of the RA patients had a DAS28 < 2.6 implying disease remission, 48 (21.2%) had low disease activity (DAS28 2.6–3.1),

Table 1. General Characteristics.

	RA-patients n = 243	Controls n = 117	P-value
Male	79 (32.5%)	46 (39.3%)	0.204
Age (yrs)	51.8±11.6	50.0±12.1	0.186
Current smoker (n,%)	54 (22.2%)	16 (13.7%)	0.051
Weight (kg)	76.3±14.4	75.3±14.5	0.520
BMI (kg/m ²)	26.1±4.3	24.9±3.7	0.006
Systolic blood pressure (mmHg)	129±18	124±13	0.004
Waist circumference (cm)	93±14	90±12	0.054
Total cholesterol (mmol/l)	5.42±1.08	5.60±1.17	0.147
LDL-C (mmol/l)	3.36±1.02	3.46±1.00	0.372
HDL-C (mmol/l)	1.54±0.51	1.55±0.43	0.894
TG (mmol/l)*	1.03 (0.72–1.42)	1.12 (0.74–1.61)	0.182
Apo A1 (g/l)	1.71±0.36	1.65±0.32	0.099
Apo B (g/l)	1.00±0.26	1.04±0.30	0.192
Glucose (mmol/l)	5.40±0.53	5.16±0.89	0.002
C-reactive protein (mg/l)*	2.00 (1.00–5.00)	1.00 (1.00–2.75)	<0.001
cIMT (mm)	0.556±0.120	0.573±0.134	0.220
Rheumatoid Factor Positive (n,%)	150 (61.7%)**	NA	
Anti-CCP positivity (n,%)	142 (58.4%)***	NA	

BMI = body mass index TG = triglycerides; Apo = apolipoprotein; cIMT = carotid intima media thickness

Data presented as mean ± standard deviation or n (%) unless stated otherwise.

* Data presented as median (interquartile range). Difference between groups tested with the Mann-Whitney U test.

** data available in 235 patients.

*** data available in 204 patients

NA, not available

doi:10.1371/journal.pone.0140844.t001

47 (20.7%) had moderate disease activity (DAS28 3.1–5.1) and 17 (7.5%) had high disease activity (DAS28 ≥ 5.1). When applying DAS28CRP, the results were similar: the median DAS28CRP was 2.2 (IQR 1.6–2.9), 113 (58.8%) had disease remission, 39 (17.3%) had low disease activity, 40 (17.7%) had moderate disease activity and 14 (6.2%) had high disease activity. cIMT did not differ between RA groups based on their disease activity (data not shown).

Univariate analysis

To identify the covariables associated with cIMT, univariate regression analysis was carried out in RA patients, controls and the combined patient and control group (Table 2). In all groups age, systolic blood pressure, LDL-C and apo B showed a significant association with cIMT. Furthermore, in controls, current smoking was significantly associated with cIMT. In RA patients male gender, BMI, waist circumference, TG, glucose and disease duration were significantly associated with cIMT. The correlation coefficient between RA disease duration and age was 0.262. A trend towards a positive association between anti-CCP positivity and cIMT was observed (B = 0.033(-0.044–0.070); P = 0.078) (Table 2). When these univariate analyses were performed only in RA patients not using biologicals, the associations between cIMT and anti-CCP positivity, disease duration, glucose and TG were no longer statistically significant (data not shown). Other parameters showed similar associations as in the total RA group.

Table 2. Univariate regression analysis identifying covariables associated with cIMT.

Variable	Controls n = 117		RA patients n = 243		Total population n = 360	
	B (95% CI)	P-value	B (95% CI)	P-value	B (95% CI)	P-value
Age (yr)	0.006 (0.004–0.007)	<0.001	0.006 (0.005–0.007)	<0.001	0.006 (0.005–0.007)	<0.001
Male sex	0.021 (-0.029–0.071)	0.243	0.050 (0.018–0.082)	0.002	0.041 (0.014–0.068)	0.003
Smoking	0.097 (0.028–0.166)	0.006	0.016 (-0.021–0.053)	0.403	0.034 (0.001–0.067)	0.044
BMI (kg/m ²)	0.005 (-0.002–0.012)	0.139	0.005 (0.001–0.008)	0.010	0.004 (0.001–0.007)	0.006
Waist (cm)	0.002 (0.000–0.004)	0.101	0.002 (0.001–0.003)	<0.001	0.002 (0.001–0.003)	<0.001
BPs (mmHg)	0.002 (0.001–0.004)	0.009	0.003 (0.002–0.004)	<0.001	0.003 (0.002–0.003)	<0.001
BPd (mmHg)	0.000 (-0.003–0.003)	0.963	0.004 (0.002–0.005)	<0.001	0.003 (0.001–0.004)	<0.001
LDL-C (mmol/l)	0.043 (0.019–0.066)	<0.001	0.034 (0.020–0.049)	<0.001	0.037 (0.025–0.050)	<0.001
HDL-C (mmol/L)	-0.010 (-0.066–0.047)	0.740	-0.010 (-0.040–0.020)	0.511	-0.010 (-0.037–0.017)	0.476
Apo AI (g/l)	0.013 (-0.065–0.091)	0.750	-0.016 (-0.059–0.027)	0.473	-0.010 (-0.048–0.028)	0.610
Apo B (g/l)	0.114 (0.034–0.194)	0.005	0.131 (0.075–0.187)	<0.001	0.126 (0.081–0.172)	<0.001
TG (mmol/l)*	0.013 (-0.020–0.045)	0.441	0.026 (0.003–0.049)	0.026	0.022 (0.003–0.040)	0.022
Glucose (mmol/l)*	0.009 (-0.019–0.036)	0.536	0.044 (0.016–0.072)	0.002	0.021 (0.001–0.040)	0.029
CRP (mg/dl)	-0.000 (-0.014–0.014)	0.964	0.001 (-0.001–0.003)	0.504	0.000 (-0.002–0.002)	0.731
Creatinin (umol/l)	0.001 (-0.001–0.003)	0.549	0.002 (0.001–0.003)	0.002	0.001 (0.001–0.003)	0.002
Absence of RA	-	-	-	-	0.017 (-0.010–0.045)	0.220
Anti-CCP	-	-	0.033 (-0.004–0.070)	0.078		
Rheumatoid factor	-	-	0.020 (-0.013–0.053)	0.231		
Erosive Disease	-	-	0.020 (-0.011–0.051)	0.212		
DAS28	-	-	0.005 (-0.009–0.019)	0.494		
DAS28CRP	-	-	-0.001 (-0.017–0.015)	0.902		
Disease duration	-	-	0.002 (0.001–0.004)	0.012		
Methotrexate	-	-	0.001 (-0.036–0.037)	0.976		
Biological agents	-	-	-0.07 (-0.039–0.024)	0.651		

RA = Rheumatoid arthritis, BPs = systolic blood pressure, BPd = diastolic blood pressure, TG = triglycerides, DAS28 = RA disease activity score 28 joints

doi:10.1371/journal.pone.0140844.t002

Multivariable analysis

Multivariable regression analysis with identified covariables associated with cIMT, was performed in RA patients, controls and in both groups combined (Table 3). In controls cIMT was significantly associated with age and current smoking and in RA patients with age and systolic blood pressure. These results remained unchanged after including glucose in the analysis. In RA patients, additional multivariable analyses including RA disease duration alone and combined with anti-CCP were performed (Table 4). The addition of RA disease activity and anti-CCP positivity, alone or combined, did not improve the predictive power of the original multivariable model (Table 4). Multivariable analysis in only those RA patients not using biological therapy showed comparable results.

Discussion

Surprisingly, cIMT in RA patients was similar to the cIMT of control subjects without RA, which is in contrast to other studies, showing increased cIMT in RA patients even early in the course of the disease [29, 30, 32, 36–38]. It is even more surprising since in our study RA patients had a higher BMI, systolic blood pressure, glucose and CRP and, a larger percentage of RA patients smoked compared to controls.

Table 3. Multivariable regression analysis with identified covariables associated with cIMT.

Variable	Controls n = 117		RA patients n = 243		Total population n = 360	
	R ² = 0.321		R ² = 0.391		R ² = 0.347	
	B (95% CI)	P-value	B (95% CI)	P-value	B (95% CI)	P-value
Age (yr)	0.005 (0.003–0.007)	<0.001	0.005 (0.004–0.006)	<0.001	0.005 (0.004–0.006)	<0.001
Male sex	0.010 (-0.037–0.056)	0.678	0.013 (-0.014–0.040)	0.336	0.015 (-0.008–0.038)	0.201
Smoking	0.101 (0.041–0.162)	0.001	0.019 (-0.011–0.049)	0.207	0.034 (0.007–0.061)	0.014
Waist (cm)	0.002 (0.000–0.004)	0.127	0.001 (0.000–0.002)	0.142	0.001 (0.000–0.002)	0.106
BPs (mmHg)	0.001 (-0.001–0.002)	0.488	0.001 (0.000–0.002)	0.003	0.001 (0.000–0.002)	0.012
LDL-C (mmol/l)	0.009 (-0.016–0.034)	0.467	0.007 (-0.006–0.020)	0.270	0.009 (-0.003–0.020)	0.143
Apo B (g/l)		**		**		**

RA = Rheumatoid arthritis, BPs = systolic blood pressure, Apo = apolipoprotein

** When using apoB instead of LDL similar results were found.

doi:10.1371/journal.pone.0140844.t003

A post hoc sample size calculation showed that our sample sizes of 243 RA-patients and 117 control patients are not sufficiently large to support the presence of a significant difference in mean cIMT between the groups with 80% power. However, considering that the difference in mean cIMT we found (0.017) cannot be considered clinically meaningful, one would need to include (at least) 270 RA and control patients (at an equivalence margin of 0.05; or 68 patients at an equivalence margin of 0.10) to support that a difference in cIMT is absent. In summary, our sample sizes are sufficiently large to support that a clinically meaningful difference in cIMT in our study is absent, with 90% power.

One of the reasons for this difference with other studies may be the relatively low disease activity in our RA patients. A higher RA disease activity results in an increased cardiovascular risk. The RA disease activity in our study was low with a total of 70% of RA patients in clinical remission or with low disease activity, which may have had a beneficial effect on cIMT. When we compared cIMT in RA patients with disease remission or low disease activity with cIMT in RA patient with moderate or high disease activity, no differences in cIMT were found. This may be due in part to the small proportion of patients with moderate to high disease activity.

Table 4. Multivariable regression analysis including RA disease characteristics.

Variable	RA patients n = 243		RA patients n = 243	
	R ² = 0.39		R ² = 0.40	
	B (95% CI)	P-value	B (95% CI)	P-value
Age (yr)	0.005 (0.004–0.006)	<0.001	0.005 (0.004–0.007)	<0.001
Male sex	0.005 (-0.016–0.039)	0.409	0.005 (-0.026–0.035)	0.786
Smoking	0.019 (-0.011–0.049)	0.212	0.020 (-0.02–0.053)	0.220
Waist (cm)	0.001 (0.000–0.002)	0.156	0.001 (0.000–0.002)	0.216
BPs (mmHg)	0.001 (0.000–0.002)	0.002	0.001 (0.000–0.000)	0.005
LDL-C (mmol/l)	0.007 (0.006–0.020)	0.282	0.007 (-0.008–0.021)	0.365
RA disease duration (yrs)	0.000 (-0.002–0.001)	0.579	0.000 (-0.002–0.002)	0.792
Anti-CCP	-	-	0.007 (-0.024–0.037)	0.661

RA = Rheumatoid Arthritis, BPs = systolic blood pressure

doi:10.1371/journal.pone.0140844.t004

Our outpatient clinic of Rheumatology is a large outpatient clinic that consists of a large and representative sample of RA patients in the Netherlands. At the outpatient clinic, a structured tight control regimen of RA disease activity is carried out reflecting the present situation in the Netherlands. Therefore, we believe that the sample of RA patients in this study is representative for the general Dutch RA population, although the results may not apply to RA patients with higher average disease activity. A meta analysis by Ambrosino et al. reported outcomes of 59 studies on cIMT in RA compared to controls [30]. Not all studies included information on disease activity, but the mean DAS28 of those studies reporting disease activity ranged from 2.6 to 6.2, which is higher than our reported median DAS28 of 2.4. Of those 59 studies, 51 reported an increased cIMT in RA patients compared to controls [30]. Only 2 studies reported DAS28 levels <3.0 showing conflicting results concerning differences in cIMT in RA compared to controls [39, 40]. The same meta-analysis by Ambrosino showed that a higher inflammatory status (DAS28, CRP, ESR) was associated with an increased cIMT [30].

Besides disease activity, several RA specific risk factors, such as the presence of rheumatoid factor, anti-CCP and erosive disease, are considered risk factors for atherosclerosis. They are all surrogate markers for the inflammatory burden since the presence of rheumatoid factor and/or anti-CCP often results in a higher disease activity. Also erosive disease occurs mostly in patients with long periods of uncontrolled disease activity. In our cohort with generally a low disease activity, rheumatoid factor and anti-CCP positivity were not associated with cIMT in the multivariable model. These results are in line with several other studies that could not demonstrate an association between these parameters and cIMT [37, 39]. A limitation of our study was that only qualitative data on rheumatoid factor and anti-CCP were available, since these characteristics were not re-tested at the time of inclusion. Recently, an *in vitro* study showed a positive correlation between the level of anti-CCPs and overexpression of thrombotic, inflammatory and pro-oxidative markers in leukocytes [18]. We did find a crude (univariate) association between disease duration and cIMT, but this association was no longer significant in the multivariable model. This is in line with a recent publication by Arts et al who found that the risk of CVD in RA patients was not increased after 10 years of disease duration when compared to the first 10 years [40]. Data on the association between cIMT and RA disease duration are conflicting since several other studies reported a positive association between cIMT and RA disease duration [37, 41, 42].

Traditional cardiovascular risk factors such as smoking, overweight, hyperlipidemia and hypertension are highly prevalent in RA patients and can be easily treated by lifestyle and pharmacological interventions. Nevertheless, underdiagnosis and undertreatment of these risk factors in RA patients have been described [22, 23, 43]. An association between cIMT and blood pressure both in the general population and in RA patients has been described previously [17, 27, 44–46]. In the present paper, regression analyses underline the association of systolic blood pressure on cIMT in RA patients. These data underscore the importance of adequate blood pressure regulation in RA.

No association between cIMT and lipid levels was found. Hyperlipidemia was strongly associated to cIMT in the univariate analysis, but it was no longer significant in the multivariable analysis after adjustment for other covariables. In contrast to the association found between smoking and cIMT in controls, no association between smoking and cIMT in RA patients was found. This is in line with earlier results reported by Gonzalez et al., who found a weaker association between smoking and CVD risk in RA patients than in non-RA patients [47].

In conclusion, in this RA cohort with relatively low disease activity, cIMT was not increased compared to controls and cIMT was not associated with RA disease characteristics. Hypertension was strongly associated with cIMT in RA patients. In light of these results and based on

findings from other studies, treatment of hypertension and other traditional CVD risk factors in RA seems warranted to reduce cardiovascular risk in RA.

Acknowledgments

The authors are grateful to René Bakker, Biomedical Scientist band 6, Department of Clinical Chemistry, Sint Franciscus Gasthuis, for the development of the study database and technical support in data management.

Author Contributions

Conceived and designed the experiments: DFvB MCC DvZ MJMWH. Performed the experiments: DFvB BK NvdM MAdV. Analyzed the data: DFvB BK GJMvdG EB DvZ MCC MJMWH. Contributed reagents/materials/analysis tools: GJMvdG HJWJ. Wrote the paper: DFvB DvZ BK GJMvdG MJMWH EB NvdM HJWJ MAdV MCC.

References

1. Gabriel SE. The epidemiology of rheumatoid arthritis. *Rheum Dis Clin North Am*. 2001; 27(2):269–81. PMID: [11396092](#)
2. Lawrence RC, Helmick CG, Arnett FC, Deyo RA, Felson DT, Giannini EH, et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum*. 1998; 41(5):778–99. PMID: [9588729](#)
3. Wolfe F, Mitchell DM, Sibley JT, Fries JF, Bloch DA, Williams CA, et al. The mortality of rheumatoid arthritis. *Arthritis Rheum*. 1994; 37(4):481–94. PMID: [8147925](#)
4. Prior P, Symmons DP, Scott DL, Brown R, Hawkins CF. Cause of death in rheumatoid arthritis. *Br J Rheumatol*. 1984; 23(2):92–9. PMID: [6722416](#)
5. Avina-Zubieta JA, Choi HK, Sadatsafavi M, Etminan M, Esdaile JM, Lacaille D. Risk of cardiovascular mortality in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Arthritis Rheum*. 2008; 59(12):1690–7. doi: [10.1002/art.24092](#) PMID: [19035419](#)
6. Steen KS, Lems WF, Visman IM, Heierman M, Dijkmans BA, Twisk JW, et al. High incidence of cardiovascular events in patients with rheumatoid arthritis. *Ann Rheum Dis*. 2009; 68(9):1509–10. doi: [10.1136/ard.2008.105023](#) PMID: [19674988](#)
7. van Halm VP, Peters MJ, Voskuyl AE, Boers M, Lems WF, Visser M, et al. Rheumatoid arthritis versus diabetes as a risk factor for cardiovascular disease: a cross-sectional study, the CARPE Investigation. *Ann Rheum Dis*. 2009; 68(9):1395–400. doi: [10.1136/ard.2008.094151](#) PMID: [18697775](#)
8. Maradit-Kremers H, Nicola PJ, Crowson CS, Ballman KV, Gabriel SE. Cardiovascular death in rheumatoid arthritis: a population-based study. *Arthritis Rheum*. 2005; 52(3):722–32. PMID: [15751097](#)
9. van Breukelen-van der Stoep DF, Klop B, van Zeven D, Hazes JM, Cabezas MC. Cardiovascular risk in rheumatoid arthritis: How to lower the risk? *Atherosclerosis*. 2013; 231(1):163–72. doi: [10.1016/j.atherosclerosis.2013.09.006](#) PMID: [24125429](#)
10. Graf J, Scherzer R, Grunfeld C, Imboden J. Levels of C-reactive protein associated with high and very high cardiovascular risk are prevalent in patients with rheumatoid arthritis. *PLoS One*. 2009; 4(7):e6242. doi: [10.1371/journal.pone.0006242](#) PMID: [19606218](#)
11. Stamatelopoulou KS, Kitas GD, Papamichael CM, Chrysoshoou E, Kyrkou K, Georgiopoulos G, et al. Atherosclerosis in rheumatoid arthritis versus diabetes: a comparative study. *Arterioscler Thromb Vasc Biol*. 2009; 29(10):1702–8. doi: [10.1161/ATVBAHA.109.190108](#) PMID: [19608975](#)
12. Libby P. Inflammation in atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2012; 32(9):2045–51. doi: [10.1161/ATVBAHA.108.179705](#) PMID: [22895665](#)
13. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med*. 1999; 340(2):115–26. PMID: [9887164](#)
14. Deanfield JE, Halcox JP, Rabelink TJ. Endothelial function and dysfunction: testing and clinical relevance. *Circulation*. 2007; 115(10):1285–95. PMID: [17353456](#)
15. van Zonneveld AJ, de Boer HC, van der Veer EP, Rabelink TJ. Inflammation, vascular injury and repair in rheumatoid arthritis. *Ann Rheum Dis*. 2009; 69 Suppl 1:i57–60.

16. del Rincon ID, Williams K, Stern MP, Freeman GL, Escalante A. High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum.* 2001; 44(12):2737–45. PMID: [11762933](#)
17. del Rincon I, Freeman GL, Haas RW, O'Leary DH, Escalante A. Relative contribution of cardiovascular risk factors and rheumatoid arthritis clinical manifestations to atherosclerosis. *Arthritis Rheum.* 2005; 52(11):3413–23. PMID: [16255018](#)
18. Barbarroja N, Perez-Sanchez C, Ruiz-Limon P, Castro-Villegas C, Aguirre MA, Carretero R, et al. Anticyclic citrullinated protein antibodies are implicated in the development of cardiovascular disease in rheumatoid arthritis. *Arterioscler Thromb Vasc Biol.* 2014; 34(12):2706–16. doi: [10.1161/ATVBAHA.114.304475](#) PMID: [25256232](#)
19. Vazquez-Del Mercado M, Nunez-Atahualpa L, Figueroa-Sanchez M, Gomez-Banuelos E, Rocha-Munoz AD, Martin-Marquez BT, et al. Serum levels of anticyclic citrullinated peptide antibodies, interleukin-6, tumor necrosis factor-alpha, and C-reactive protein are associated with increased carotid intima-media thickness: a cross-sectional analysis of a cohort of rheumatoid arthritis patients without cardiovascular risk factors. *Biomed Res Int.* 2015; 2015:342649. doi: [10.1155/2015/342649](#) PMID: [25821796](#)
20. Meek IL, Picavet HS, Vonkeman HE, Verschuren WM, van de Laar MA. Increased cardiovascular risk factors in different rheumatic diseases compared with the general population. *Rheumatology (Oxford).* 2013; 51(1):210–6.
21. Naranjo A, Sokka T, Descalzo MA, Calvo-Alen J, Horslev-Petersen K, Luukkainen RK, et al. Cardiovascular disease in patients with rheumatoid arthritis: results from the QUEST-RA study. *Arthritis Res Ther.* 2008; 10(2):R30. doi: [10.1186/ar2383](#) PMID: [18325087](#)
22. Panoulas VF, Toms TE, Metsios GS, Stavropoulos-Kalinoglou A, Kosovitsas A, Millionis HJ, et al. Target organ damage in patients with rheumatoid arthritis: the role of blood pressure and heart rate. *Atherosclerosis.* 2009; 209(1):255–60. doi: [10.1016/j.atherosclerosis.2009.08.047](#) PMID: [19781703](#)
23. Chung CP, Giles JT, Petri M, Szklo M, Post W, Blumenthal RS, et al. Prevalence of traditional modifiable cardiovascular risk factors in patients with rheumatoid arthritis: comparison with control subjects from the multi-ethnic study of atherosclerosis. *Semin Arthritis Rheum.* 2012; 41(4):535–44. doi: [10.1016/j.semarthrit.2011.07.004](#) PMID: [22340996](#)
24. Kerekes G, Soltesz P, Nurmohamed MT, Gonzalez-Gay MA, Turiel M, Vegh E, et al. Validated methods for assessment of subclinical atherosclerosis in rheumatology. *Nat Rev Rheumatol.* 2012; 8(4):224–34. doi: [10.1038/nrrheum.2012.16](#) PMID: [22349611](#)
25. Zhang Y, Guallar E, Qiao Y, Wasserman BA. Is carotid intima-media thickness as predictive as other noninvasive techniques for the detection of coronary artery disease? *Arterioscler Thromb Vasc Biol.* 2014; 34(7):1341–5. doi: [10.1161/ATVBAHA.113.302075](#) PMID: [24764454](#)
26. Hannawi S, Haluska B, Marwick TH, Thomas R. Atherosclerotic disease is increased in recent-onset rheumatoid arthritis: a critical role for inflammation. *Arthritis Res Ther.* 2007; 9(6):R116. PMID: [17986352](#)
27. Chatterjee Adhikari M, Guin A, Chakraborty S, Sinhamahapatra P, Ghosh A. Subclinical atherosclerosis and endothelial dysfunction in patients with early rheumatoid arthritis as evidenced by measurement of carotid intima-media thickness and flow-mediated vasodilatation: an observational study. *Semin Arthritis Rheum.* 2012; 41(5):669–75. doi: [10.1016/j.semarthrit.2011.08.003](#) PMID: [22035626](#)
28. van Sijl AM, Peters MJ, Knol DK, de Vet HC, Gonzalez-Gay MA, Smulders YM, et al. Carotid intima media thickness in rheumatoid arthritis as compared to control subjects: a meta-analysis. *Semin Arthritis Rheum.* 2011; 40(5):389–97. doi: [10.1016/j.semarthrit.2010.06.006](#) PMID: [20889191](#)
29. Sodergren A, Karp K, Boman K, Eriksson C, Lundstrom E, Smedby T, et al. Atherosclerosis in early rheumatoid arthritis: very early endothelial activation and rapid progression of intima media thickness. *Arthritis Res Ther.* 2010; 12(4):R158. doi: [10.1186/ar3116](#) PMID: [20712865](#)
30. Ambrosino P, Lupoli R, Di Minno A, Tasso M, Peluso R, Di Minno MN. Subclinical atherosclerosis in patients with rheumatoid arthritis. A meta-analysis of literature studies. *Thromb Haemost.* 2015; 113(5):916–30. doi: [10.1160/TH14-11-0921](#) PMID: [25716931](#)
31. Myasoedova E, Gabriel SE. Cardiovascular disease in rheumatoid arthritis: a step forward. *Curr Opin Rheumatol.* 2010; 22(3):342–7. doi: [10.1097/BOR.0b013e3283379b91](#) PMID: [20164773](#)
32. Im CH, Kim NR, Kang JW, Kim JH, Kang JY, Bae GB, et al. Inflammatory burden interacts with conventional cardiovascular risk factors for carotid plaque formation in rheumatoid arthritis. *Rheumatology (Oxford).* 2015; 54(5):808–15.
33. Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum.* 1988; 31(3):315–24. PMID: [3358796](#)

34. Klop B, van de Geijn GJ, Bovenberg SA, van der Meulen N, Elte JW, Birnie E, et al. Erythrocyte-bound apolipoprotein B in relation to atherosclerosis, serum lipids and ABO blood group. *PLoS One*. 2013; 8(9):e75573. doi: [10.1371/journal.pone.0075573](https://doi.org/10.1371/journal.pone.0075573) PMID: [24069429](https://pubmed.ncbi.nlm.nih.gov/24069429/)
35. Bovenberg SA, Klop B, Alipour A, Martinez-Hervas S, Westzaan A, van de Geijn GJ, et al. Erythrocyte-associated apolipoprotein B and its relationship with clinical and subclinical atherosclerosis. *Eur J Clin Invest*. 2012; 42(4):365–70. doi: [10.1111/j.1365-2362.2011.02591.x](https://doi.org/10.1111/j.1365-2362.2011.02591.x) PMID: [21913916](https://pubmed.ncbi.nlm.nih.gov/21913916/)
36. Mahajan V, Handa R, Kumar U, Sharma S, Gulati G, Pandey RM, et al. Assessment of atherosclerosis by carotid intima-media thickness in patients with rheumatoid arthritis. *J Assoc Physicians India*. 2008; 56:587–90. PMID: [19051702](https://pubmed.ncbi.nlm.nih.gov/19051702/)
37. Kumeda Y, Inaba M, Goto H, Nagata M, Henmi Y, Furumitsu Y, et al. Increased thickness of the arterial intima-media detected by ultrasonography in patients with rheumatoid arthritis. *Arthritis Rheum*. 2002; 46(6):1489–97. PMID: [12115178](https://pubmed.ncbi.nlm.nih.gov/12115178/)
38. Del Rincon I, Williams K, Stern MP, Freeman GL, O'Leary DH, Escalante A. Association between carotid atherosclerosis and markers of inflammation in rheumatoid arthritis patients and healthy subjects. *Arthritis Rheum*. 2003; 48(7):1833–40. PMID: [12847676](https://pubmed.ncbi.nlm.nih.gov/12847676/)
39. Ristic GG, Lepic T, Glisic B, Stanisavljevic D, Vojvodic D, Petronijevic M, et al. Rheumatoid arthritis is an independent risk factor for increased carotid intima-media thickness: impact of anti-inflammatory treatment. *Rheumatology (Oxford)*. 2010; 49(6):1076–81.
40. Arts EE, Fransen J, den Broeder AA, Popa CD, van Riel PL. The effect of disease duration and disease activity on the risk of cardiovascular disease in rheumatoid arthritis patients. *Ann Rheum Dis*. 2015; 74(6):998–1003. doi: [10.1136/annrheumdis-2013-204531](https://doi.org/10.1136/annrheumdis-2013-204531) PMID: [24458537](https://pubmed.ncbi.nlm.nih.gov/24458537/)
41. Grover S, Sinha RP, Singh U, Tewari S, Aggarwal A, Misra R. Subclinical atherosclerosis in rheumatoid arthritis in India. *J Rheumatol*. 2006; 33(2):244–7. PMID: [16465654](https://pubmed.ncbi.nlm.nih.gov/16465654/)
42. Gonzalez-Juanatey C, Llorca J, Testa A, Revuelta J, Garcia-Porrua C, Gonzalez-Gay MA. Increased prevalence of severe subclinical atherosclerotic findings in long-term treated rheumatoid arthritis patients without clinically evident atherosclerotic disease. *Medicine (Baltimore)*. 2003; 82(6):407–13.
43. Rollefstad S, Kvien TK, Holme I, Eirheim AS, Pedersen TR, Semb AG. Treatment to lipid targets in patients with inflammatory joint diseases in a preventive cardio-rheuma clinic. *Ann Rheum Dis*. 2012; 72(12):1968–74. doi: [10.1136/annrheumdis-2012-202789](https://doi.org/10.1136/annrheumdis-2012-202789) PMID: [23264359](https://pubmed.ncbi.nlm.nih.gov/23264359/)
44. Solomon A, Woodiwiss AJ, Abdool-Carrim AT, Stevens BA, Norton GR, Dessein PH. The carotid artery atherosclerosis burden and its relation to cardiovascular risk factors in black and white Africans with established rheumatoid arthritis: a cross-sectional study. *J Rheumatol*. 2012; 39(9):1798–806. doi: [10.3899/jrheum.120073](https://doi.org/10.3899/jrheum.120073) PMID: [22753659](https://pubmed.ncbi.nlm.nih.gov/22753659/)
45. Bots ML, Hofman A, de Bruyn AM, de Jong PT, Grobbee DE. Isolated systolic hypertension and vessel wall thickness of the carotid artery. The Rotterdam Elderly Study. *Arterioscler Thromb*. 1993; 13(1):64–9. PMID: [8422340](https://pubmed.ncbi.nlm.nih.gov/8422340/)
46. Mitrovic J, Morovic-Vergles J, Horvatic I, Badzak J, Stojic M, Gamulin S. Ambulatory arterial stiffness index and carotid intima-media thickness in hypertensive rheumatoid patients: a comparative cross-sectional study. *Int J Rheum Dis*. 2015.
47. Gonzalez A, Maradit Kremers H, Crowson CS, Ballman KV, Roger VL, Jacobsen SJ, et al. Do cardiovascular risk factors confer the same risk for cardiovascular outcomes in rheumatoid arthritis patients as in non-rheumatoid arthritis patients? *Ann Rheum Dis*. 2008; 67(1):64–9. PMID: [17517756](https://pubmed.ncbi.nlm.nih.gov/17517756/)