

# EUR Research Information Portal

Parental education, health literacy and children's adult body height

**Published in:**

Journal of Biosocial Science

**Publication status and date:**

Published: 15/11/2019

**DOI (link to publisher):**

[10.1017/S0021932019000737](https://doi.org/10.1017/S0021932019000737)

**Document Version**

Publisher's PDF, also known as Version of record

**Document License/Available under:**

Article 25fa Dutch Copyright Act

**Citation for the published version (APA):**

Jarosz, E., & Gugushvili, A. (2019). Parental education, health literacy and children's adult body height. *Journal of Biosocial Science*, 52(5), 696-718. <https://doi.org/10.1017/S0021932019000737>

[Link to publication on the EUR Research Information Portal](#)

**Terms and Conditions of Use**

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.


In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

**Take-down policy**

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: [openaccess.library@eur.nl](mailto:openaccess.library@eur.nl). Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

RESEARCH ARTICLE

# Parental education, health literacy and children's adult body height

Ewa Jarosz<sup>1,2\*</sup>  and Alexi Gugushvili<sup>3,4</sup>

<sup>1</sup>Institute of Philosophy and Sociology, Polish Academy of Sciences, Warsaw, Poland, <sup>2</sup>Department of Sociology, Centre for Time Use Research, University of Oxford, Oxford, UK, <sup>3</sup>Department of Public Administration and Sociology, Erasmus University Rotterdam, the Netherlands and <sup>4</sup>Department of Social Policy and Intervention, University of Oxford, Oxford, UK  
\*Corresponding author. Email: [ejarosz@ifispan.waw.pl](mailto:ejarosz@ifispan.waw.pl)

(Received 18 May 2019; revised 11 September 2019; accepted 11 September 2019)

## Abstract

Human anthropometric traits, while significantly determined by genetic factors, are also affected by an individual's early life environment. An adult's body height is a valid indicator of their living conditions in childhood. Parental education has been shown to be one of the key covariates of individuals' health and height, both in childhood and adulthood. Parental functional literacy has been demonstrated to be another important determinant of child health, but this has largely been overlooked in studies on height. The objective of this study was to analyse the associations between parents' education, their functional literacy and their children's adult body height. The study used data for 39,240 individuals from the 2016 wave of the nationally representative Life in Transition Survey (LITS) conducted in 34 countries in Southern and Eastern Europe, the Middle East and Central Asia. Using linear and Poisson models, regression adjustment treatment estimators and multilevel mixed-effects linear regressions, the study analysed the links between mother's and father's educational attainment, parental functional literacy, measured by the number of books in the childhood home, and children's adult height. The models also included other individual and contextual covariates of height. The results demonstrated that mother's educational attainment and parental functional literacy have independent associations with children's adult body height. Sufficient literacy skills of the parent may have a positive effect on children's growth even if parental education is low. These associations remained significant across time. The study also provides evidence of a widening of the height gap for men born in the period just before and after systemic transition in post-socialist societies, which may suggest an increase in social differences in early living standards.

**Keywords:** Body height; Parental education; Functional literacy

## Introduction

Among the many facets of persisting social inequality, those that shape human anthropometric dimensions have recently attracted much scholarly attention (Deaton, 2008; Batty *et al.*, 2009; Zurawiecka *et al.*, 2019). Adult body height is a highly heritable trait, but it has also been shown to be associated with early life environment across a set of developed and developing countries (Meyer & Selmar, 1999; Silventoinen *et al.*, 1999; Pawlowski *et al.*, 2000; Chen & Li, 2009). Furthermore, it may be linked to inter-generational transmission of existing social inequalities. Taller people are more likely to report better health (Silventoinen *et al.*, 1999), find a mating partner (Tao & Yin, 2016), experience fewer adverse health events in their lifetime (Smith, 2000; Jousilahti *et al.*, 2000), be better educated (Silventoinen, 2003) and earn a higher wage (Schultz, 2002).

Twin studies attribute between 0.70 and 0.90 of variance in height to heritability (Silventoinen *et al.*, 2003; Jelenkovic *et al.*, 2016). Evidence from genome-wide association studies (GWAS)

further support these estimates (Visscher *et al.*, 2006). Such studies have allowed the identification of specific genes associated with height, but a significant part of the genetic variation remains unexplained (Visscher *et al.*, 2010; Perola, 2011). With regard to environmental influences, the majority of factors affecting height operate at the pre-puberty stage (Gunnell, 2002). Adult body height is linked to their height in childhood, with the height-for-age z-score (HAZ) being a cross-nationally validated indicator of children's level of nutrition (Chen & Li, 2009). Early living conditions are influenced primarily by parents' and grandparents' socioeconomic characteristics (Peck & Lundberg, 1995; Moltchanova & Eriksson, 2015), and in particular their educational attainment. Parental education has been shown to have a significant association with both children's HAZ and their nutritional status (Moestue & Huttly, 2008). Studies on adoptees have suggested that in the case of height the impact of nurture – that is, everyday childrearing practices – may be as strong as that of genetic makeup (Chen & Li, 2009).

Parental education may affect a child's development through different channels. Father's education is more likely to represent the social status of the family, and may give access to higher social positions and resources that can be invested in the child's health and well-being. Mother's education may be indicative of the quality of everyday childcare (Silventoinen, 2003). Educated mothers are also more likely to immunize their children and engage in more effective everyday health-promoting behaviours (Desai & Alva, 1998). Education has also been related to individuals' fertility choices (Lindeboom *et al.*, 2009). Maternal age, number of siblings, self-care during pregnancy and socioeconomic position before birth have all been shown to be associated with anthropometric traits and the life-long health of children (Hack *et al.*, 2003; Black *et al.*, 2007; Barker, 2012), with babies born to mothers in the lowest educational categories being, on average, the shortest (Howe *et al.*, 2012).

Education has also been linked with certain lifestyle choices. As such, it is often assumed to act as a proxy for an individual's ability to access and utilize health information (Currie & Moretti, 2003). While formal education may indeed increase these abilities (Woods-Townsend *et al.*, 2018), an individual's educational attainment should not be seen as equivalent to their level of functional literacy. Functional literacy and, specifically, functional health literacy, is defined as a skill that is necessary for acquiring knowledge on good health practices, and for using this knowledge to make appropriate health-related decisions (Driessnack *et al.*, 2014). Sufficient health literacy means that an individual is able to understand medical information and to act upon that information (Sanders *et al.*, 2004). Research shows that even in countries with high levels of formal literacy, a significant proportion of the population is functionally illiterate (Kirsch, 1993; Sanders *et al.*, 2009), which means that they are able to read but not to understand written information. The inability to comprehend and process health-related information has been seen as one of the contributors to existing health and social inequalities, including those starting in early childhood (Braveman & Barclay 2009; Sikora *et al.*, 2019).

The health literacy of parents has been linked to important health outcomes for their children (DeWalt & Hink, 2009), including their nutrition status and body mass index (Chari *et al.*, 2014). Low parental health literacy has been found to be associated with a substantially higher probability of exhibiting behaviours negatively affecting child health (Sanders *et al.*, 2009). Both education and health literacy rank among the key independent determinants of health (Kickbusch, 2001). However, in studies analysing parental influence on children's anthropometric traits, the impact of health literacy is either implicitly assumed to be accounted for by educational attainment, or is not acknowledged at all. Since positive links between education, health literacy and children's health have been reported across different societies and in many social contexts (DeWalt & Hink, 2009), it is likely that both parental education and their health literacy will be independently associated with children's attained height.

Adult height has also been shown to be associated with other factors operating at the individual level. To account for those characteristics, together with parental education and health literacy, this study used a set of potential covariates of height: type of settlement where the respondent and

their parents were born, family's socioeconomic position and whether they belonged to an ethnic majority in the country. Urban and rural settlements are likely to differ in terms of access to information and health care services. Urbanization status is associated with significant height and weight differences in children, with urban children in low- and middle-income categories being on average taller and heavier than their rural counterparts (Paciorek *et al.*, 2013). Furthermore, selective migration may affect height distribution between rural and urban areas as more socially mobile individuals are also likely to be taller (Zielińska, 1991; Silventoinen *et al.*, 1999). The socioeconomic position of the father has been used as a key indicator of a family's socioeconomic situation, but the evidence of its impact on individual height is inconclusive. Some studies have reported significant links between father's position in the socioeconomic hierarchy and child height (Silventoinen, 2003), while others have reported no such association (Hasle & Boldsen, 1991). Lastly, ethnic majority status may account for possible inequalities related to the usual disadvantaged position of ethnic minorities (Nazroo, 1998). Ethnic minority groups are more likely to concentrate in deprived residential areas (Karlsen & Nazroo, 2002) with poorer access to health care facilities. Potential hostility, and different forms of abuse from majority groups, are likely to affect an individual's mental and physical health (Karlsen & Nazroo 2002). Ethnicity-related risks are very relevant for many countries covered in the Life in Transition Survey (LITS). In the late 1980s and early to mid-1990s most of the countries included in the LITS dataset underwent an abrupt systemic transition. Many country borders have appeared relatively recently, and the quality of life and living conditions of ethnic minorities have been compromised, in particular for the cohorts whose infancy and childhood fell in the times of increased ethnic violence.

Adult height has also been linked to broader contextual factors, including year of birth and the general living conditions in a country. Human height increased consistently throughout the 20<sup>th</sup> century due to improvements in living conditions and child nutrition, and the quality of health care (Bielicki, 1986; Howe *et al.*, 2012; Moltchanova & Eriksson, 2015). In countries covered by the LITS, an individual's birth cohort may be important not only because of the general trends in height dynamics, but also the region-specific history. Firstly, the impact of the Second World War might be seen for cohorts born during or immediately after the war. Earlier studies have demonstrated that individuals born around the time of the war were significantly shorter, and that this result could not be explained by cohort effects alone (Moltchanova & Eriksson, 2015). Furthermore, systemic transition during the 1990s resulted in major cuts to social and health spending, and a substantial increase in social inequality (King *et al.*, 2009; Azarova *et al.* 2017).

The other important contextual covariate – a country's level of economic development at birth – represents general standards of living. Together with other environmental factors they are estimated to account for around 20% of the variation in a population's height, with this number being even higher in poorer societies (Silventoinen, 2003; Quintana-Domeque *et al.*, 2011).

## Methods

The study used data from the LITS commissioned by the European Bank for Reconstruction and Development (EBRD, 2016). The LITS was conducted in 2016 in the following 34 countries in Central, Southern and Eastern Europe, the Middle East and Central Asia: Albania, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Bulgaria, Croatia, the Czech Republic, Cyprus, Estonia, North Macedonia, Georgia, Germany, Greece, Hungary, Italy, Kazakhstan, Kosovo, the Kyrgyz Republic, Latvia, Lithuania, Moldova, Mongolia, Montenegro, Poland, Romania, Russia, Serbia, Slovakia, Slovenia, Tajikistan, Turkey, Ukraine and Uzbekistan.

Respondents were drawn randomly, using a two-stage sampling procedure. Primary sampling units (PSU) were electoral districts, polling station territories, census enumeration districts or other administrative areas. Secondary sampling units (SSU) were households. Each country

had a minimum of 50 PSUs with each PSU containing at least 20 SSUs. After deletion of missing information and censoring individuals aged below 20 (those who had not yet achieved their full height), 17,331 men and 21,909 women were included in the analyses. The LITS has been used in comparative social and health research (Gugushvili, 2019; Gugushvili *et al.*, 2019), but post-socialist countries have generally been under-represented in multi-country surveys (Slomczynski & Tomescu-Dubrow, 2006). Also, there are no other comprehensive accounts of the association between parental socioeconomic characteristics and children's anthropometric dimensions in the majority of the analysed countries.

### **Individuals' height**

All LITS respondents were asked to report their height with the following question: What is your height in centimetres without shoes? Empirical reports on the accuracy of self-reported height are not conclusive, with height often being overestimated by both men and women (Engstrom *et al.*, 2003; Gorber *et al.*, 2007). The LITS allows how respondents' self-reported heights correspond to their objectively measured heights to be checked. In one randomly selected PSU in each country respondents' height was measured using a portable stadiometer. This took place after the declarations of height had been made by the respondents. The results of the bivariate analysis of reported and measured height showed that the mean difference between reported (170.2 cm) and measured (170.3 cm) height was  $-0.05$  cm, and it was not statistically significant (95% confidence interval (CI)  $-0.27$  to  $0.18$ ) (Gugushvili & Jarosz, 2019).

### **Individual-level independent variables**

As mother's and father's education could affect the child's height via different channels, both were included in the models. The following categories were used for mother's and father's educational attainment: (1) no education; (2) primary and lower secondary education; (3) secondary and post-secondary education; and (4) tertiary education.

The preferred measure of parental functional health literacy would be a version of the Test of Functional Health Literacy in Adults (TOFHLA) (Parker *et al.*, 1995). To the authors' knowledge, the TOFHLA is not available in any of the cross-national datasets open to public use. However, earlier research has demonstrated that the number of books in the parental home is a good proxy for parents' functional health literacy (Sanders *et al.*, 2004; Driessnack *et al.*, 2014). Specifically, having ten children's books, or more than ten adults' books, at home has been shown to be a valid independent indicator that a parent has adequate health literacy (Sanders *et al.*, 2004). In this study, the number of books in the respondent's childhood home was used as indicative of parental functional health literacy. The original LITS question asked about the approximate number of books in the respondent's childhood home, not counting magazines, newspapers and school books. Four possible answer options were given: (1) none or very few (0–10 books); (2) enough to fill one shelf (11–25 books); (3) enough to fill one bookcase (26–100 books); and (4) enough to fill two or more bookcases (101–200 books). In the current study, the terms 'health literacy' and 'functional literacy' are used interchangeably as they both refer to the ability to understand written information.

Mother's and father's sectors of employment were used as proxies for the socioeconomic status of the household. The LITS offers no information on parental social position or household income, but using both of these variables may give a picture of an individual's childhood living conditions. This variable included the following categories (same for both parents): (a) employed in agriculture; (b) employed in manufacture; (c) employed in public administration; (d) employed in other not classified sector; and (e) parent never worked.

Birth cohort was coded based on the respondent's year of birth in the following categories: (a) born before 1946; (b) 1946–1955; (c) 1956–1965; (d) 1966–1975; (e) 1976–1985; and

(f) 1986–1995. Type of settlement at birth used an original binary variable asking whether a respondent was born in a rural or an urban settlement. The LITS allows differentiation between rural and urban type of settlement at birth for both respondents and their parents, and all relevant variables were used in the study. To account for the respondent's minority status, a binary variable was created with a value of 1 for respondents belonging to the ethnic majority in their country of residence.

### **Country's level of development at birth**

To derive this contextual variable the measure of Gross Domestic Product (GDP) *per capita* by Purchasing Power Parities (PPP) in international dollars with fixed 2011 prices was used. This measure considers inflation and differences in cost of living between countries. The source of this variable is Gapminder (2018), which has compiled the data from numerous sources, such as official statistics, historical sources and their own estimates. Gapminder is arguably the only comparative dataset that provides information on the level of economic development for most years of the 20th century for each country included in the LITS. This allowed information on the level of economic development for individuals who were born as early as in the first decades of the 20<sup>th</sup> century to be derived.

### **Analytical strategy**

The analytical strategy consisted of a two-stage approach. First, baseline and fully adjusted ordinary least squares (OLS) models were fitted with individual height as a continuous dependent variable. In this approach unobserved country characteristics were accounted for by the country fixed effects. However, the focus of the study was not only on an individual's absolute height, but also on how parental education, parental functional literacy and other covariates were linked with the probability of them appearing at the lower or higher end of the height distribution. To derive these relative measures, in the second step, quartiles of height in all cohorts were calculated separately for each country. Next, binary variables were created for those whose height fell within the bottom 25% of the height distribution, and for those whose height appeared in the top 25% of the distribution by sex in a given country. These variables were used as outcome variables in the Poisson regression models. Conventional logistic regressions with corresponding odds ratios are likely to overestimate the actual associations of the independent variables in the model. On the other hand, the Poisson regressions, used in this study for binary outcome variables, allowed prevalence ratios with corresponding 95% CIs to be derived, which are more appropriate measures of association with high prevalence of positive outcomes in the binary dependent variables (Barros & Hirakata, 2003).

To assess the robustness of the findings, the treatment effects were estimated via regression adjustment of linear and Poisson models. This approach uses contrasts of averages of treatment-specific predicted outcomes to estimate treatment effects. In other words, regression adjustment is based on a two-step approach to estimating treatment effects. First, separate regression models of the outcome on a set of covariates were fitted for each reference group used for social comparison. Second, the averages of the predicted outcomes were computed for each subject and reference group. The contrasts of these averages provided the estimates of average treatment effects.

In order to identify if the broader social environment moderated the associations between mother's education, parental functional literacy and an individual's adult height, aggregate mean values of both of these variables were generated by each country and by cohort. Next, they were fitted into the multilevel fixed effects linear regressions. Models were estimated separately by sex using Stata 15 statistical software.



## Results

### *Descriptive statistics and bivariate associations*

Table 1 presents the descriptive statistics for the independent variables, and distributions of these variables by groups of individuals in the bottom and top height quartiles. Regarding parental educational attainment, less than 10% of fathers had no formal education, while this share was just above 10% for mothers. In all instances, levels of parental education were higher among individuals in the top height quartile with corresponding significant differences in proportions when compared with individuals in the bottom height quartile. Regarding parental functional health literacy, represented by the number of books in the childhood home, a significant proportion of individuals (up to 30%) grew up in homes with ten or fewer books. Individuals who reported having a larger number of books in the parental home were more likely to fall within the top quartile of the height distribution.

### *Multivariate analyses*

Tables 2 and 3 show the associations between parental education, parental functional literacy and individual's height controlling for a set of individual and contextual characteristics. In the fully adjusted models (Models 1 in Table 2 and Models 1 and 2 in Table 3), father's education mattered for individual's height only in certain instances: for instance, having a father with tertiary education compared with having a father with no education was associated with a 1.0 cm increase in height for men and a 0.8 cm increase for women. In contrast, mother's education was positively associated with height in all models for both men and women. Children of better-educated mothers had a much lower chance of being in the bottom of the height distribution within respective cohorts, and much a higher chance of appearing in the top height quartile. For instance, compared with individuals having mothers with no education, the sons of tertiary-educated mothers had a risk prevalence of 0.70 for being in the bottom height quartile, while the daughters of tertiary-educated mothers had a risk prevalence of 1.34 for being in the top height quartile.

Parental functional literacy was positively and significantly associated with both men's and women's height, net of other variables in the models. Individuals who reported having more than ten books in their childhood home were taller than those who reported having fewer than ten books. More specifically, in the adjusted Model 2 in Table 2, men and women growing up in households with more than 100 books were, on average, 1.3 and 1.2 cm taller than those who had only a few books in their parental home. The number of books mattered for both men and women in terms of the likelihood of being in the lowest and highest quartiles of the height distribution. In Table 3, both men and women were about 20% less likely to appear in the bottom height quartile within their cohorts if they grew up in homes with enough books to fill two or more bookcases.

Both men's and women's mean heights increased over the 20<sup>th</sup> century. Men and women born in 1986–1995 were, on average, 4.2 and 3.0 cm taller, respectively, than those who were born before 1946. Furthermore, the type of settlement where respondents were born was associated with their height. Men born in rural areas were, on average, 0.7 cm shorter than men born in urban areas. Furthermore, men whose fathers were born in rural areas were significantly taller compared with men whose fathers were born in urban areas.

In some instances, mother's sector of employment was associated with individual's height. Men whose mothers were employed in manufacturing, public administration or in other, unspecified sectors had, respectively, 18%, 22% and 26% higher chances of being in the top quartile of the height distribution compared with men whose mothers worked in agriculture. Father's sector of employment was not associated with individuals' height. No evidence was found for a significant association between ethnic minority status or level of economic development at birth and individual's height.

**Table 1.** Descriptive statistics (% and means) of study individuals and bivariate associations

	Men				Women			
	Full sample (%)	% in bottom height quartile	% in top height quartile	<i>p</i> -value	Full sample (%)	% in bottom height quartile	% in top height quartile	<i>p</i> -value
<b>Father's education</b>								
No education	7.1	8.1	5.4		7.4	8.4	5.6	
Primary/lower secondary	45.5	49.2	41.0		44.9	46.5	43.8	
Secondary/post-secondary	36.2	34.0	38.8		36.2	35.0	37.0	
Tertiary	11.2	8.8	14.9	<0.001	11.4	10.1	13.6	<0.001
<b>Mother's education</b>								
No education	10.6	12.2	8.3		11.0	12.8	8.3	
Primary/lower secondary	46.4	48.8	43.2		45.7	46.6	45.5	
Secondary/post-secondary	34.2	32.3	35.7		34.3	32.8	35.2	
Tertiary	8.8	6.7	12.9	<0.001	9.0	7.8	11.0	<0.001
<b>Number of books</b>								
0–10	29.2	33.2	23.9		27.5	30.4	23.3	
11–25	27.7	29.1	25.5		26.5	26.3	26.6	
26–100	26.6	23.5	30.4	<0.001	28.5	27.5	29.8	<0.001
101+	16.5	14.1	20.2		17.5	15.8	20.2	
<b>Birth cohort</b>								
<1946	10.1	10.8	9.8		14.0	14.1	13.2	
1946–1955	14.7	14.1	15.0		16.0	16.8	16.4	
1956–1965	18.7	18.7	18.5		17.7	17.4	18.0	
1966–1975	19.2	19.0	19.5		17.1	17.1	17.0	
1976–1985	20.1	20.0	20.3	0.513	18.7	18.0	18.7	0.598
1986–1995	17.2	17.3	16.9		16.5	16.7	16.7	
<b>Type of settlement</b>								
Born in urban area	54.2	51.0	58.1	<0.001	54.4	52.5	56.9	<0.001
Born in rural area	45.8	49.0	41.9		45.6	47.5	43.1	
Father born in urban area	34.3	32.1	35.9	<0.001	35.7	33.2	37.8	<0.001
Father born in rural area	65.7	67.9	64.1		64.3	66.8	62.2	

(Continued)



Table 1. (Continued)

	Men				Women			
	Full sample (%)	% in bottom height quartile	% in top height quartile	p-value	Full sample (%)	% in bottom height quartile	% in top height quartile	p-value
Mother born in urban area	35.2	33.0	38.1	<0.001	36.3	34.1	37.8	<0.001
Mother born in rural area	64.8	67.0	61.9		63.7	65.9	62.2	
Father's occupation								
Agriculture	51.3	55.0	46.7		52.0	53.2	51.1	
Manufacturing	33.5	31.1	37.0		32.9	32.2	33.3	
Public administration	7.5	6.5	9.1		7.7	7.2	8.31	
Not classified	4.6	4.9	5.0	<0.001	4.9	5.1	5.20	0.082
Never worked	3.1	2.5	2.5		2.5	2.3	2.1	
Mother's occupation								
Agriculture	32.8	36.5	27.4		34.0	34.6	33.0	
Manufacturing	31.7	29.2	35.0		31.2	30.2	33.7	
Public administration	8.4	7.0	10.9		8.2	7.4	9.1	
Not classified	5.1	4.9	5.9	<0.001	5.3	5.7	5.6	<0.000
Never worked	21.9	22.4	20.8		21.2	22.1	18.6	
Ethnicity								
Ethnic majority	16.2	17.5	15.9	0.033	15.8	17.1	15.4	0.010
Ethnic minority	83.8	82.5	84.1		84.2	82.9	84.6	
Economic development at birth	8.24 (6.05)	8.13 (0.08)	8.33 (0.09)	0.096	7.72 (5.66)	7.61 (0.06)	7.84 (0.08)	0.021

The p-values are from bivariate tests of differences in means or proportions for each covariate across the bottom and top height quartiles. Source: authors' analysis of data from EBRD (2016).

**Interaction between mother's education and parental functional literacy**

Figure 1 shows the predicted values for height in centimetres and the predicted likelihood of appearing in the bottom and top quartiles of the height distribution by mother's level of education and parental functional literacy. In most cases the differences between individuals having mothers with no education and those having mothers with tertiary education were significant regardless of the number of books at home. At the same time, regardless of the level of maternal education, the number of books at home showed a strong and consistent association with individuals' height. For instance, among men with tertiary-educated mothers who also had 100 or more books in the childhood home the probability of appearing in the lowest quartile of the height distribution was around 0.20, whereas for individuals with low-educated mothers but the same number of books at home it was around 0.30. Furthermore, if mothers had tertiary education but the number of books at home was low, the probability of appearing in the bottom quartile increased to 0.37. In short, there was no interaction effect between maternal education and the number of books at home, but rather these two variables maintained an independent association with individuals' height.

**Table 2.** Parental education, parental health literacy and individuals' adult height: point estimates from OLS regression models ( $\beta$  [95% CI])

	Men		Women	
	Model 1 Height in cm	Model 2 Height in cm	Model 1 Height in cm	Model 2 Height in cm
Father's education (Ref.: no education)				
Primary/lower secondary	0.30 [-0.37, 0.96]	-0.08 [-0.76, 0.59]	<b>0.71</b> [0.19, 1.22]	0.39 [-0.09, 0.87]
Secondary/post-secondary	<b>1.32</b> [0.46, 2.18]	0.49 [-0.38, 1.35]	<b>1.19</b> [0.62, 1.75]	<b>0.66</b> [0.10, 1.22]
Tertiary	<b>1.54</b> [0.67, 2.42]	<b>0.99</b> [0.04, 1.93]	<b>1.28</b> [0.64, 1.93]	<b>0.81</b> [0.18, 1.44]
Mother's education (Ref.: no education)				
Primary/lower secondary	<b>1.71</b> [1.24, 2.18]	<b>0.67</b> [0.21, 1.14]	<b>1.42</b> [0.97, 1.86]	<b>0.62</b> [0.16, 1.07]
Secondary/post-secondary	<b>2.95</b> [2.22, 3.69]	<b>1.06</b> [0.31, 1.80]	<b>2.38</b> [1.75, 3.00]	<b>0.86</b> [0.25, 1.46]
Tertiary	<b>4.40</b> [3.48, 5.32]	<b>1.98</b> [1.13, 2.83]	<b>3.19</b> [2.35, 4.02]	<b>1.33</b> [0.56, 2.10]
Number of books (Ref.: 0–10)				
11–25	<b>0.51</b> [0.20, 0.83]	<b>0.36</b> [0.06, 0.65]	<b>0.97</b> [0.68, 1.26]	<b>0.76</b> [0.50, 1.02]
26–100	<b>1.49</b> [1.08, 1.89]	<b>1.17</b> [0.80, 1.54]	<b>1.11</b> [0.83, 1.39]	<b>0.87</b> [0.58, 1.15]
101+	<b>1.55</b> [1.00, 2.10]	<b>1.28</b> [0.73, 1.83]	<b>1.46</b> [1.09, 1.82]	<b>1.24</b> [0.91, 1.57]
Birth cohort (Ref.: <1946)				
1946–1955		<b>1.60</b> [1.22, 1.99]		<b>1.84</b> [1.50, 2.18]
1956–1965		<b>2.73</b> [2.25, 3.22]		<b>2.75</b> [2.39, 3.10]
1966–1975		<b>4.06</b> [3.42, 4.71]		<b>3.73</b> [3.30, 4.17]
1976–1985		<b>4.32</b> [3.59, 5.06]		<b>3.78</b> [3.21, 4.34]
1986–1995		<b>4.42</b> [3.65, 5.18]		<b>3.98</b> [3.43, 4.54]
Type of settlement (Ref.: urban)				
Born in rural area		<b>-0.69</b> [-1.10, -0.29]		-0.22 [-0.49, 0.05]
Father born in rural area		<b>0.56</b> [0.15, 0.96]		-0.25 [-0.51, 0.00]
Mother born in rural area		0.00 [-0.38, 0.38]		-0.02 [-0.34, 0.29]
Father's occupation (Ref.: agriculture)				
Manufacturing		0.20 [-0.04, 0.44]		-0.28 [-0.63, 0.06]
Public administration		-0.02 [-0.47, 0.44]		-0.30 [-0.81, 0.20]
Not classified		-0.28 [-1.03, 0.47]		-0.33 [-0.91, 0.26]
Never worked		-0.36 [-1.64, 0.93]		0.19 [-0.86, 1.24]
Mother's occupation (Ref.: agriculture)				
Manufacturing		<b>0.53</b> [0.13, 0.94]		-0.00 [-0.36, 0.35]
Public administration		0.54 [-0.13, 1.20]		0.00 [-0.50, 0.50]

(Continued)

Table 2. (Continued)

	Men		Women	
	Model 1 Height in cm	Model 2 Height in cm	Model 1 Height in cm	Model 2 Height in cm
Not classified		<b>0.70</b> [0.11, 1.30]		-0.38 [-0.89, 0.12]
Never worked		0.12 [-0.36, 0.60]		-0.42 [-0.81, -0.03]
Ethnic majority		0.19 [-0.42, 0.80]		0.29 [-0.35, 0.94]
Economic development at birth		0.00 [-0.06, 0.06]		0.03 [-0.03, 0.08]
Akaike Information Criterion (AIC)	129,449.1	118,131.6	158,728.0	146,508.6
Bayesian Information Criterion (BIC)	129,535.4	118,372.2	158,816.7	146,756.5
$R^2$	0.19	0.22	0.13	0.16
Observations	17,331	17,331	21, 909	21,909

All models account for country fixed effects; 95% CIs in parentheses; significant associations are shown in bold.  
Source: authors' analysis of data from EBRD (2016).

Table 3. Parental education, parental health literacy and individuals' adult height: prevalence ratios from Poisson regression models (PR [95%CI])

	Men		Women	
	Model 1 Height in bottom quartile	Model 2 Height in top quartile	Model 1 Height in bottom quartile	Model 2 Height in top quartile
Father's education (Ref.: no education)				
Primary/lower secondary	1.03 [0.92, 1.16]	1.02 [0.87, 1.19]	0.97 [0.89, 1.07]	1.04 [0.87, 1.23]
Secondary/post-secondary	0.93 [0.82, 1.06]	1.17 [0.94, 1.44]	0.93 [0.83, 1.03]	1.07 [0.88, 1.28]
Tertiary	<b>0.84</b> [0.71, 0.99]	1.24 [0.99, 1.54]	0.90 [0.79, 1.02]	1.12 [0.91, 1.39]
Mother's education (Ref.: no education)				
Primary/lower secondary	<b>0.86</b> [0.79, 0.94]	<b>1.11</b> [1.03, 1.19]	<b>0.86</b> [0.78, 0.95]	<b>1.20</b> [1.03, 1.39]
Secondary/post-secondary	0.82 [0.72, 0.95]	1.12 [0.98, 1.29]	<b>0.82</b> [0.72, 0.94]	<b>1.25</b> [1.05, 1.49]
Tertiary	<b>0.70</b> [0.58, 0.83]	<b>1.40</b> [1.20, 1.62]	<b>0.73</b> [0.62, 0.86]	<b>1.34</b> [1.09, 1.64]
Number of books (Ref.: 0–10)				
11–25	<b>0.93</b> [0.87, 0.99]	1.05 [0.97, 1.12]	<b>0.90</b> [0.85, 0.94]	<b>1.13</b> [1.06, 1.21]
26–100	<b>0.80</b> [0.74, 0.86]	<b>1.23</b> [1.13, 1.33]	<b>0.88</b> [0.82, 0.94]	<b>1.15</b> [1.07, 1.24]
101+	<b>0.80</b> [0.72, 0.88]	<b>1.23</b> [1.12, 1.36]	<b>0.82</b> [0.75, 0.89]	<b>1.22</b> [1.12, 1.33]
Birth cohort (Ref.: <1946)				
1946–1955	0.95 [0.88, 1.02]	0.94 [0.86, 1.03]	1.06 [0.98, 1.16]	0.99 [0.89, 1.09]
1956–1965	1.07 [0.98, 1.17]	<b>0.85</b> [0.76, 0.96]	1.02 [0.93, 1.12]	0.94 [0.86, 1.02]
1966–1975	<b>1.14</b> [1.03, 1.27]	<b>0.82</b> [0.73, 0.92]	1.11 [0.98, 1.27]	<b>0.89</b> [0.82, 0.97]
1976–1985	<b>1.22</b> [1.07, 1.39]	<b>0.75</b> [0.66, 0.85]	1.11 [0.97, 1.27]	<b>0.87</b> [0.77, 0.99]
1986–1995	<b>1.26</b> [1.10, 1.44]	<b>0.70</b> [0.60, 0.82]	<b>1.16</b> [1.00, 1.35]	<b>0.87</b> [0.76, 0.99]

(Continued)


Table 3. (Continued)

	Men		Women	
	Model 1 Height in bottom quartile	Model 2 Height in top quartile	Model 1 Height in bottom quartile	Model 2 Height in top quartile
Type of settlement (Ref.: urban)				
Born in rural areas	<b>1.15</b> [1.07, 1.24]	0.94 [0.86, 1.03]	1.04 [0.97, 1.10]	0.98 [0.89, 1.07]
Father born in rural area	<b>0.90</b> [0.82, 0.98]	<b>1.15</b> [1.06, 1.24]	1.04 [0.97, 1.11]	<b>0.92</b> [0.85, 1.00]
Mother born in rural area	0.98 [0.91, 1.06]	<b>0.90</b> [0.83, 0.98]	1.02 [0.95, 1.10]	1.03 [0.96, 1.11]
Father's occupation (Ref.: agriculture)				
Manufacturing	0.98 [0.91, 1.04]	1.05 [0.99, 1.13]	1.03 [0.97, 1.10]	0.92 [0.84, 1.01]
Public administration	1.00 [0.89, 1.13]	1.00 [0.89, 1.13]	1.04 [0.95, 1.15]	0.92 [0.82, 1.03]
Not classified	1.13 [0.99, 1.31]	1.00 [0.86, 1.16]	1.05 [0.92, 1.21]	0.97 [0.83, 1.13]
Never worked	1.10 [0.88, 1.37]	1.21 [0.97, 1.52]	0.95 [0.86, 1.04]	0.99 [0.70, 1.39]
Mother's occupation (Ref.: agriculture)				
Manufacturing	<b>0.92</b> [0.85, 1.00]	<b>1.18</b> [1.06, 1.30]	1.03 [0.95, 1.10]	1.04 [0.94, 1.14]
Public administration	0.93 [0.81, 1.08]	<b>1.22</b> [1.06, 1.39]	1.01 [0.92, 1.10]	1.02 [0.91, 1.14]
Not classified	0.93 [0.81, 1.07]	<b>1.26</b> [1.09, 1.45]	1.18 [1.06, 1.31]	1.01 [0.90, 1.13]
Never worked	0.98 [0.91, 1.06]	1.07 [0.98, 1.17]	1.10 [1.03, 1.18]	0.90 [0.80, 1.01]
Ethnic majority	0.92 [0.85, 1.01]	0.97 [0.87, 1.09]	0.94 [0.82, 1.08]	1.06 [0.94, 1.19]
Economic development at birth	1.00 [0.99, 1.00]	1.00 [0.99, 1.01]	1.00 [0.99, 1.00]	1.00 [0.99, 1.01]
Akaike Information Criterion (AIC)	22,351.3	18,861.8	28,634.8	23,968.8
Bayesian Information Criterion (BIC)	22,591.8	19,102.4	28,882.6	24,216.7
R <sup>2</sup>	0.010	0.012	0.006	0.005
Observations	17, 331	17, 331	21, 909	21, 909

All models account for country fixed effects; 95% CIs are in parentheses; significant associations are shown in bold. Source: authors' analysis of data from EBRD (2016).

### Regression adjustment estimators

Table 4 presents the treatment estimators from regression adjustment models. For men's height the results were very similar to the estimates using conventional OLS regression models in Table 2. Those who had mothers with tertiary education were 1.94 (CI 0.17 to 3.70) cm taller than men with the lowest educated mothers. For the women's sample, using the more robust treatment effect estimators removed the associations with maternal education found in the conventional regression models. Nonetheless, parental health literacy, measured by the number of books in the childhood home, was consistently associated with higher height, lower likelihood of appearing in the bottom height quartile, and higher probability of appearing at the top of the height distribution. For instance, after regression adjustment those who grew up in a home with more than 100 books were 6% less likely to be in the bottom quartile of the height distribution in their respective cohorts.



**Figure 1.** Predicted values of respondents' height conditioned by maternal education and the number of books during childhood. Results are based on the models in Tables 2 and 3. Error bars represent 95% CIs. Source: authors' analysis of data from EBRD (2016).

**Table 4.** Average effects of mother's education and number of books in the parental home on individuals' adult height from treatment estimators using regression adjustment models

	Height in cm (OLS)	In bottom height quartile (Poisson)	In top height quartile (Poisson)
<b>Men</b>			
Mother's education			
Primary/lower secondary vs no education	<b>1.71</b> [0.48, 2.95]	<b>-0.11</b> [-0.19, -0.03]	0.04 [-0.01, 0.10]
Secondary/post-secondary vs no education	<b>2.26</b> [0.99, 3.53]	<b>-0.12</b> [-0.21, -0.04]	<b>0.06</b> [0.01, 0.11]
Tertiary vs no education	<b>1.94</b> [0.17, 3.70]	-0.08 [-.20, 0.03]	0.07 [-0.01, 0.14]
Number of books			
11–25 vs 0–10	<b>0.57</b> [0.18, 0.96]	<b>-0.04</b> [-0.06, -0.01]	0.01 [-0.02, 0.02]
26–100 vs 0–10	<b>1.40</b> [1.00, 1.79]	<b>-0.08</b> [-0.10, -0.05]	<b>0.04</b> [0.02, 0.06]
101+ vs 0–10	<b>1.29</b> [0.80, 1.78]	<b>-0.07</b> [-0.10, -0.03]	0.02 [-0.01, 0.05]
<b>Women</b>			
Mother's education			
Primary/lower vs no education	0.23 [-0.61, 1.07]	-0.02 [-0.07, 0.04]	0.03 [-0.02, 0.07]
Secondary/post-secondary vs no education	0.62 [-0.26, 1.50]	-0.03 [-0.09, 0.02]	0.03 [-0.02, 0.09]
Tertiary vs no education	0.96 [-0.23, 2.15]	-0.04 [-0.13, 0.04]	0.04 [-0.03, 0.11]
Number of books			
11–25 vs 0–10	<b>0.72</b> [0.40, 1.04]	<b>-0.03</b> [-0.05, -0.01]	<b>0.03</b> [0.01, 0.05]
26–100 vs 0–10	<b>0.82</b> [0.50, 1.13]	<b>-0.04</b> [-0.06, -0.02]	<b>0.03</b> [0.01, 0.05]
101+ vs 0–10	<b>1.07</b> [0.65, 1.49]	<b>-0.06</b> [-0.08, -0.03]	<b>0.03</b> [0.01, 0.05]

Models account for all covariates shown in Tables 1–3 and country fixed effects; 95% CIs are in parentheses; significant associations are shown in bold.

Source: authors' analysis of data from EBRD (2016).


### **Change over time in the association between maternal education and number of books with individual's height**

Figure 2 shows the associations between maternal education, parental functional literacy and individual's height for individuals born in different time periods from pre-1945 until 1995. The first observed trend is that, regardless of level of maternal education and number of books, individuals' height increased throughout the entire period. The most significant change was observed for men born in 1986–1995, when the role of maternal education also became more salient. More specifically, the lowest level of maternal education was associated with 174 cm of predicted height, while the highest level of education was associated with 178 cm of height. The same pattern was also repeated in terms of attaining the highest and the lowest position in the relative height distribution. The difference between individuals born to the highest- and lowest-educated mothers increased in both instances. For women, there were no systemic or significant differences across time in terms of the effects of maternal education and the number of books in the childhood home.

### **Contextual factors**

In Table 5, the mean values of maternal education and number of books in the childhood home, calculated for specific countries and cohorts, were first introduced in multilevel mixed-effects





**Figure 2.** Predicted values of individuals' height and position in the height distribution by birth cohort of individuals. Results are based on the models in Tables 2 and 3. Source: authors' analysis of data from EBRD (2016).

**Table 5.** Parental education, parental health literacy and their interactions with contextual environment, and point estimates from multilevel mixed-effects linear regressions ( $\beta$  [95% CI])

	Men		Women	
	Model 1 Main effects	Model 2 Interactions	Model 1 Main effects	Model 2 Interactions
Mother's education (Ref.: no education)				
Primary/lower secondary	<b>0.66</b> [0.21, 1.11]	0.15 [-1.39, 1.69]	<b>0.62</b> [0.19, 1.05]	<b>1.37</b> [0.11, 2.62]
Secondary/post-secondary	<b>1.04</b> [0.31, 1.76]	0.73 [-1.44, 2.90]	<b>0.86</b> [0.28, 1.44]	<b>2.03</b> [0.01, 4.05]
Tertiary	<b>1.97</b> [1.13, 2.80]	0.22 [-3.23, 3.67]	<b>1.36</b> [0.63, 2.08]	0.76 [-2.28, 3.80]
Mean level of education within cohorts	-0.10 [-0.90, 0.70]	-0.30 [-1.31, 0.71]	-0.34 [-0.86, 0.17]	-0.15 [-0.85, 0.55]
Cross-level interactions				
Mean education $\times$ primary/lower secondary		0.18 [-0.42, 0.79]		-0.26 [-0.71, 0.18]
Mean education $\times$ secondary/post-secondary		0.14 [-0.59, 0.86]		-0.36 [-0.95, 0.23]
Mean education $\times$ tertiary		0.46 [-0.47, 1.40]		0.05 [-0.82, 0.92]
Akaike Information Criterion (AIC)	118,325.7	118,323.7	146,692.8	146, 683.7
Bayesian Information Criterion (BIC)	118,589.6	118,587.6	146,964.6	146, 947.5
Observations	17,331	17,331	21,909	21, 909
Number of books (Ref.: 0–10)				
11–25	<b>0.35</b> [0.06, 0.64]	-0.46 [-1.27, 2.18]	<b>0.76</b> [0.51, 1.01]	0.61 [-1.28, 2.49]
26–100	<b>1.16</b> [0.79, 1.52]	-1.31 [-3.13, 0.51]	<b>0.86</b> [0.60, 1.13]	0.59 [-1.73, 2.90]
101+	<b>1.27</b> [0.74, 1.79]	-1.93 [-4.54, 0.69]	<b>1.23</b> [0.92, 1.54]	-0.60 [-3.26, 2.06]
Mean number of books within cohorts	0.63 [-0.33, 1.59]	0.06 [-1.00, 1.12]	0.13 [-0.73, 0.99]	-0.10 [-1.22, 1.03]
Cross-level interactions				
Mean number of books $\times$ books: 11–25		-0.00 [-0.85, 0.85]		0.08 [-0.79, 0.95]
Mean number of books $\times$ books: 26–100		<b>1.08</b> [0.25, 1.91]		0.14 [-0.86, 1.14]
Mean number of books $\times$ books: 101+		<b>1.33</b> [0.23, 2.44]		0.74 [-0.39, 1.87]
Akaike Information Criterion (AIC)	118,321.9	118,306.9	146,695.7	146,688.9
Bayesian Information Criterion (BIC)	118,578.0	118,570.8	146,967.5	146,952.7
Observations	17,331	17,331	21,909	21,909

Models account for all covariates shown in Tables 1–3. 95% CIs are in parentheses; significant associations are shown in bold. Source: authors' analysis of data from EBRD (2016).

linear models as macro-level variables, and then interacted with individual-level variables describing mother's education and parental functional literacy. The results suggest that the aggregate level of maternal education and the number of books do not explain the variation in adult height. The cross-level interaction terms in most cases were not statistically significant, but for men the

number of books had a greater effect on height in the context when the mean number of books reported in the cohort was higher.

### **Further analyses: individual's own educational attainment and health**

The additional models in Table 6 were run to account for two types of associations frequently reported in other studies on height: those between individuals' height and self-reported health and between individuals' height and their own educational attainment. The main analyses were re-run, this time also including respondents' attained level of education and self-reported health, which were expected to moderate, at least to a certain extent, the association between parental characteristics and individuals' height. The models showed that respondent's education was a statistically significant covariate of their height. Better-educated individuals were taller than their lower-educated counterparts. Accounting for own education reduced the association of maternal education and height but it remained significant in most instances. Regarding parental functional literacy, represented by the number of books in the childhood home, this variable remained one of the most significant explanations of individuals' own height.

Lastly, individuals' self-reported health was negatively and significantly related to their height. Including this variable in the models did not change the substantial and statistically significant associations between mother's education, parental functional literacy and individuals' height.

## **Discussion**

In line with earlier reports, this study found that parental educational attainment was positively associated with their offspring's attained height. Studies to date have been inconclusive regarding comparisons between maternal and paternal effects, with some finding that the impacts of father's and mother's education are similar (Moestue & Huttly, 2008), and other reporting that the effect of father's education is substantially weaker (Chen & Li, 2009). The present study found a consistent and positive association between mother's education and children's height, net of other socio-demographic and contextual characteristics. In the fully adjusted models, the association of father's education with individual's height was significant only in some instances and only for individuals with tertiary-educated fathers. It was also substantially weaker than the association with mother's education.

The impact of maternal education on adult height was stronger for men, in particular in more robust regression adjustment models, which is somewhat contradictory to results reported for the United States, Brazil and Ghana (Thomas, 1994), and possibly due to the specific cultural settings. The present findings are likely to reflect differences in the child rearing of boys and girls occurring primarily during early childhood (Case & Paxson, 2008). An important factor that could not be accounted for in the analyses is that mother's education has also been found to be associated with the child's risk of dying by the age of two (Kickbusch, 2001). As infant mortality rates, in particular for the older cohorts covered in LITS, were substantial (Ksenofontova, 1994), it may be that the effect of maternal education would be even stronger had these children survived and been included in the survey.

The number of books in the parental home, indicative of parental functional health literacy, was consistently found to be associated with individuals' height for both men and women. The association with parental functional literacy was independent of other factors included in the models and was as strong as the association with mother's education. These results were largely confirmed using regression adjustment treatment estimators from linear and Poisson models. In this study, the number of any books, apart from school books, was used as a proxy for health literacy, but earlier research has shown that the number of children's books at home may also be indicative of parental care and time dedicated to children early in life (Brunello *et al.*, 2016), with

**Table 6.** Parental education, parental health literacy and individuals' adult height accounting for individuals' own education and self-reported health

	Men			Women		
	Model 1 Height in cm (OLS)	Model 2 Height in bottom quartile (Poisson)	Model 3 Height in top quartile (Poisson)	Model 1 Height in cm (OLS)	Model 2 Height in bottom quartile (Poisson)	Model 3 Height in top quartile (Poisson)
<b>Models with respondents' own education</b>						
Father's education (Ref.: no education)						
Primary/lower secondary	-0.36 [-1.05, 0.33]	1.08 [0.96, 1.21]	0.99 [0.84, 1.15]	0.28 [-0.21, 0.78]	1.00 [0.91, 1.11]	<b>1.40</b> [1.06, 1.85]
Secondary/post-secondary	-0.10 [-1.01, 0.81]	1.04 [0.91, 1.19]	1.07 [0.87, 1.32]	0.49 [-0.05, 1.03]	0.97 [0.87, 1.08]	<b>1.44</b> [1.08, 1.91]
Tertiary	0.19 [-0.78, 1.16]	0.98 [0.82, 1.16]	1.10 [0.88, 1.37]	<b>0.62</b> [0.02, 1.21]	0.94 [0.83, 1.08]	<b>1.52</b> [1.14, 2.03]
Mother's education (Ref.: no education)						
Primary/lower secondary	<b>0.47</b> [0.01, 0.93]	<b>0.90</b> [0.82, 0.98]	1.07 [1.00, 1.16]	<b>0.54</b> [0.07, 1.00]	<b>0.87</b> [0.79, 0.97]	0.99 [0.83, 1.19]
Secondary/post-secondary	<b>0.73</b> [0.00, 1.46]	0.88 [0.76, 1.01]	1.07 [0.94, 1.23]	<b>0.74</b> [0.12, 1.36]	<b>0.85</b> [0.74, 0.97]	1.01 [0.84, 1.22]
Tertiary	<b>1.57</b> [0.73, 2.42]	<b>0.75</b> [0.63, 0.90]	<b>1.31</b> [1.12, 1.54]	<b>1.20</b> [0.39, 2.01]	<b>0.76</b> [0.64, 0.90]	1.05 [0.85, 1.30]
Number of books (Ref.: 0–10)						
11–25	0.19 [-0.11, 0.48]	0.96 [0.90, 1.02]	1.02 [0.95, 1.09]	<b>0.70</b> [0.45, 0.95]	<b>0.91</b> [0.86, 0.96]	<b>1.17</b> [1.02, 1.35]
26–100	<b>0.87</b> [0.48, 1.25]	<b>0.85</b> [0.78, 0.91]	<b>1.17</b> [1.08, 1.27]	<b>0.79</b> [0.51, 1.06]	<b>0.89</b> [0.83, 0.96]	<b>1.22</b> [1.03, 1.45]
101+	<b>0.86</b> [0.27, 1.44]	<b>0.87</b> [0.78, 0.96]	<b>1.16</b> [1.05, 1.28]	<b>1.14</b> [0.80, 1.47]	<b>0.84</b> [0.76, 0.92]	<b>1.30</b> [1.06, 1.59]
Own education (Ref.: no education)						
Primary/lower secondary	1.16 [-0.49, 2.82]	0.90 [0.71, 1.13]	1.11 [0.72, 1.72]	0.49 [-0.63, 1.61]	0.88 [0.74, 1.06]	<b>1.12</b> [1.04, 1.20]
Secondary/post-secondary	<b>2.38</b> [0.68, 4.09]	<b>0.71</b> [0.55, 0.91]	1.34 [0.86, 2.07]	0.80 [-0.33, 1.94]	0.82 [0.67, 1.00]	<b>1.14</b> [1.06, 1.22]
Tertiary	<b>3.29</b> [1.56, 5.02]	<b>0.59</b> [0.46, 0.75]	1.54 [0.99, 2.41]	0.96 [-0.15, 2.08]	<b>0.79</b> [0.64, 0.96]	<b>1.19</b> [1.09, 1.30]
<b>Models with respondents' self-reported health</b>						
Father's education (Ref.: no education)						
Primary/lower secondary	-0.17 [-0.85, 0.51]	1.05 [0.93, 1.18]	1.01 [0.86, 1.18]	0.35 [-0.14, 0.83]	0.98 [0.90, 1.08]	1.03 [0.87, 1.22]
Secondary/post-secondary	0.38 [-0.49, 1.26]	0.95 [0.84, 1.08]	1.15 [0.93, 1.42]	0.57 [-0.00, 1.14]	0.94 [0.85, 1.05]	1.05 [0.87, 1.26]

(Continued)

Table 6. (Continued)

	Men			Women		
	Model 1 Height in cm (OLS)	Model 2 Height in bottom quartile (Poisson)	Model 3 Height in top quartile (Poisson)	Model 1 Height in cm (OLS)	Model 2 Height in bottom quartile (Poisson)	Model 3 Height in top quartile (Poisson)
Tertiary	0.83 [-0.12, 1.78]	0.86 [0.73, 1.02]	1.21 [0.96, 1.51]	<b>0.71</b> [0.06, 1.35]	0.92 [0.81, 1.04]	1.10 [0.88, 1.36]
Mother's education (Ref.: no education)						
Primary/lower secondary	<b>0.66</b> [0.20, 1.12]	<b>0.87</b> [0.79, 0.95]	<b>1.11</b> [1.03, 1.19]	<b>0.55</b> [0.10, 1.00]	<b>0.87</b> [0.79, 0.96]	<b>1.19</b> [1.03, 1.37]
Secondary/post-secondary	<b>0.98</b> [0.23, 1.73]	<b>0.83</b> [0.72, 0.96]	1.11 [0.97, 1.27]	<b>0.77</b> [0.15, 1.38]	<b>0.84</b> [0.73, 0.95]	<b>1.24</b> [1.04, 1.47]
Tertiary	<b>1.89</b> [1.04, 2.74]	<b>0.70</b> [0.59, 0.84]	<b>1.38</b> [1.18, 1.61]	<b>1.21</b> [0.44, 1.98]	<b>0.75</b> [0.64, 0.88]	<b>1.31</b> [1.07, 1.61]
Number of books (Ref.: 0–10)						
11–25	<b>0.29</b> [0.00, 0.58]	<b>0.94</b> [0.89, 1.00]	1.04 [0.96, 1.11]	<b>0.69</b> [0.44, 0.95]	<b>0.91</b> [0.86, 0.95]	<b>1.12</b> [1.04, 1.20]
26–100	<b>1.06</b> [0.70, 1.42]	<b>0.81</b> [0.76, 0.87]	<b>1.21</b> [1.12, 1.32]	<b>0.78</b> [0.49, 1.07]	<b>0.89</b> [0.83, 0.95]	<b>1.14</b> [1.05, 1.23]
101+	<b>1.17</b> [0.63, 1.71]	<b>0.81</b> [0.74, 0.90]	<b>1.21</b> [1.10, 1.34]	<b>1.14</b> [0.80, 1.48]	<b>0.83</b> [0.76, 0.91]	<b>1.20</b> [1.10, 1.30]
Own health (Ref.: Medium)						
Very bad	<b>-1.20</b> [-2.20, -0.20]	1.16 [0.97, 1.39]	1.04 [0.81, 1.35]	<b>-1.09</b> [-1.85, -0.33]	<b>1.32</b> [1.19, 1.46]	0.93 [0.79, 1.10]
Bad	<b>-0.73</b> [-1.28, -0.19]	<b>1.12</b> [1.01, 1.24]	0.89 [0.78, 1.02]	<b>-0.47</b> [-0.80, -0.14]	<b>1.04</b> [0.96, 1.12]	<b>0.90</b> [0.82, 0.99]
Good	<b>0.48</b> [0.15, 0.81]	<b>0.91</b> [0.86, 0.97]	1.07 [0.99, 1.16]	<b>0.83</b> [0.57, 1.10]	<b>0.89</b> [0.84, 0.94]	<b>1.15</b> [1.08, 1.23]
Very good	<b>1.55</b> [1.11, 1.99]	<b>0.78</b> [0.71, 0.85]	<b>1.25</b> [1.13, 1.38]	<b>1.50</b> [0.96, 2.04]	<b>0.76</b> [0.67, 0.86]	<b>1.31</b> [1.19, 1.45]

Our multilevel model account for all covariate shown in Tables 1 and 2, including country fixed effects; 95% CIs in parentheses; significant associations are shown in bold.

Source: authors' analysis of data from EBRD (2016).

such resources playing an even more important role in low-income families (Tomopoulos *et al.*, 2006).

The available data did not allow for more detailed analysis of different aspects of health literacy, but it has been argued that health literacy includes the following three components: access to health care, connection between the patient and health provider and self-care (Paasche-Orlow & Wolf, 2007). The relationship between these three aspects is not straightforward, as even individuals' high literacy levels do not guarantee that they will respond in a desired way to the health information they receive (Nutbeam, 2000). The implication of this knowledge for the present study is twofold. Firstly, a higher number of books in the childhood home may be seen as a factor that, overall, positively contributes to children's physical and cognitive development, but it may also be indicative of both parental care and time spent with children, and the likelihood that a parent would understand and follow written medical advice. Secondly, the effect of an individual's functional literacy on health should be analysed in relation to a broader social context, in particular that of systemic limitations (such as significant cuts in health care provisions in post-socialist countries after the transition to a market economy), or access to different sources of health information.

After accounting for the independent effects of parental education and health literacy, there were no interaction effects between the number of books in the parental home and mother's level of education, but predicted mean height for both men and women was significantly higher in each mother educational category if there were more than 100 books in the parental home. Furthermore, having a high number of books lowered the probability of falling within the bottom quartile of the height distribution, even for those whose mothers were low-educated. The level of education and related level of health literacy had been treated as a single variable in earlier studies describing competences involved in health care decision-making (Smith *et al.*, 2009). The present findings, together with other reports (Kalichman *et al.*, 2000), show that an individual's level of education and their functional literacy should be analysed separately as these characteristics may have independent effects on the variable of interest.

As expected, birth cohort mattered for an individual's height, and there was a gradual increase in average height for the entire population over the 20<sup>th</sup> century. There was also a consistent height gap between individuals with tertiary-educated and low-educated mothers, as well as between those who were brought up in homes with a low and a high number of books. This gap persisted across the entire time covered in the analyses, i.e. it appeared in each cohort. A novel finding not reported previously is that of different trajectories of height differences for men and women across time. In the case of men, among the youngest cohorts there was a divergence between the mean height of individuals whose mothers had no education and those with tertiary-educated mothers. In other words, the height gap related to mother's education was higher for individuals born just before and soon after the collapse of socialism. Conversely, the height gap between women whose mothers had no education and those whose mothers had tertiary education did not widen, which is in line with the claims that boys are more vulnerable to socioeconomic changes (Petranović *et al.*, 2014). It can be assumed that this drift in height was caused by the negative social and health consequences of systemic transition, with individuals in low social positions, including those with low educational qualifications, being affected most negatively (Cockerham, 2002; Doniec *et al.*, 2018). Similar findings have been reported in other studies from Central Eastern Europe (Petranović *et al.*, 2014).

The association identified between type of settlement and individuals' height may be indicative of mobility patterns. Migrants are generally healthier than non-migrants (Lu, 2008), which is the first selection mechanism. There is also evidence that taller individuals are more likely to move from rural to urban areas compared with shorter individuals in equivalent social positions (Zielińska, 1991). The findings of this study show that men whose fathers were born in rural areas, but who themselves were born in urban areas, were more likely to fall within the top quartile of the height distribution. This may suggest that more spatially mobile fathers were taller and, for this reason, had taller sons. No such effect was found for women.



This study comes with a set of limitations. Firstly, it did not account for genetic determinants of height due to a lack of information on parental height in the survey. Secondly, indicators of material deprivation during childhood, or, alternatively, of household income, and associated quality of nutrition, were not available. Earlier studies reporting a relationship between an individual's height and the material situation in the household found that the association between these variables, although significant, did not decrease the importance of parental education (Silventoinen, 2003). Because household income was not controlled for, it might be that some of the variance explained by the number of books could be attributed to economic differences between households. However, disparities in height related to the number of books in the parental home did not increase over time, including after the systemic transition in post-socialist countries, which suggests that the association between the number of books and the financial situation of the household may be weak, if any.

Despite these limitations, this study offers a unique contribution to the existing research by tracking the long-term trends in height and its childhood-related explanations in the large number of countries in Central, Southern and Eastern Europe, the Middle East and Central Asia. It has demonstrated that the main explanatory variables – mother's education and parental health literacy – maintained independent associations with individuals' height. On the macro level, individuals' height was associated with being born in a particular cohort, but not with the general economic conditions at birth, which suggests that there is a secular increase of individuals' height not accounted for by economic growth. However, the substantial increase in the height gap for the cohort born around the times of the systemic transition in post-socialist societies suggests that the newly emerged social inequalities may play a detrimental role for health equity, including shaping individuals' early living conditions and consequently may affect disparities in individuals' height in adulthood.

**Funding.** This research received no specific grant from any funding agency, commercial entity or not-for-profit organization.

**Conflicts of Interest.** The authors have no conflicts of interest to declare.

**Ethical Approval.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

## References

- Azarova A, Irdam D, Gugushvili A, Fazekas M, Scheiring G, Horvat P, Stefler D *et al.* (2017) The effect of rapid privatisation on mortality in mono-industrial towns in post-Soviet Russia: a retrospective cohort study. *The Lancet Public Health* 2(5), e231–e238.
- Barker DJP (2012) Developmental origins of chronic disease. *Public Health* 126(3), 185–189.
- Barros AJ and Hirakata VN (2003) Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Medical Research Methodology* 3(1), 21.
- Batty GD, Shipley MJ, Gunnell D, Huxley R, Kivimaki M, Woodward M *et al.* (2009) Height, wealth, and health: an overview with new data from three longitudinal studies. *Economics & Human Biology* 7(2), 137–152.
- Bielicki T (1986) Physical growth as a measure of the economic well-being of populations: the twentieth century. In Falkner F and Tanner JM (eds) *Human Growth: A Comprehensive Treatise. Volume 3: Methodology and Ecological, Genetic, and Nutritional Effects on Growth*. Plenum Press, New York, pp. 283–305.
- Black S, Devereux P and Salvanes K (2007) From the cradle to the labour market? The effect of birth weight on adult outcomes. *Quarterly Journal of Economics* 122(1), 409–439.
- Braveman P and Barclay C (2009) health disparities beginning in childhood: a life-course perspective. *Pediatrics* 124 (Supplement 3), S163–S175.
- Brunello G, Weber G and Weiss C (2016) *Books are Forever: Early Life Conditions, Education and Lifetime Earnings in Europe*. *Economic Journal* 127(600), 271–296.
- Case A and Paxson C (2008) Stature and status: height, ability, and labor market outcomes. *Journal of Political Economy* 116(3), 499–532.

- Chari R, Warsh J, Ketterer T, Hossain J and Sharif I** (2014) Association between health literacy and child and adolescent obesity. *Patient Education and Counseling* **94**(1), 61–66.
- Chen Y and Li H** (2009) Mother's education and child health: is there a nurturing effect? *Journal of Health Economics* **28**(2), 413–426.
- Cockerham W** (2002) *Health and Social Change in Russia and Eastern Europe*. Routledge, New York.
- Currie J and Moretti E** (2003) Mother's education and the intergenerational transmission of human capital: evidence from college openings. *Quarterly Journal of Economics* **118**(4), 1495–1532.
- Deaton A** (2008) Height, health, and inequality: the distribution of adult heights in India. *American Economic Review* **98**(2), 468–474.
- Desai S and Alva S** (1998) Maternal education and child health: is there a strong causal relationship? *Demography* **35**(1), 71–81.
- DeWalt DA and Hink A** (2009) Health literacy and child health outcomes: a systematic review of the literature. *Pediatrics* **124** (Supplement 3), S265–S274.
- Donic K, Stefler D, Murphy M, Gugushvili A, McKee M, Marmot M et al.** (2018) Education and mortality in three Eastern European populations: findings from the PrivMort retrospective cohort study. *European Journal of Public Health* **29**(3), 549–554.
- Driessnack M, Chung S, Perkhounkova E and Hein M** (2014) Using the “newest vital sign” to assess health literacy in children. *Journal of Pediatric Health Care* **28**(2), 165–171.
- EBRD** (2016) *Life in Transition Survey III*. European Bank for Reconstruction and Development, London.
- Engstrom J, Paterson S, Doherty A, Trabulsi M and Speer K** (2003) Accuracy of self-reported height and weight in women: an integrative review of the literature. *Journal of Midwifery & Women's Health* **48**(5), 338–345.
- Gapminder** (2018) *GDP Per Capita, PPP Constant 2011 International \$*. Gapminder, Stockholm.
- Gorber SC, Tremblay M, Moher D and Gorber B** (2007) A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obesity Reviews* **8**(4), 307–326.
- Gugushvili A** (2019) A multilevel analysis of perceived intergenerational mobility and welfare state preferences. *International Journal of Social Welfare* **28**(1), 16–30.
- Gugushvili A and Jarosz E** (2019) Inequality, validity of self-reported height, and its implications for BMI estimates: An analysis of randomly selected primary sampling units' data. *Preventive Medicine Reports*, **16**(100974), 1–8.
- Gugushvili A, Jarosz E and McKee M** (2019) Compared with whom? Reference groups in socio-economic comparisons and self-reported health in 34 countries. *International Journal of Epidemiology*, doi: [10.1093/ije/dyz122](https://doi.org/10.1093/ije/dyz122)
- Gunnell D** (2002) Can adult anthropometry be used as a “biomarker” for prenatal and childhood exposures? *International Journal of Epidemiology* **31**(2), 390–394.
- Hack M, Schluchter M, Cartar L, Rahman M, Cuttler L and Borawski E** (2003) Growth of very low birth weight infants to age 20 years. *Pediatrics* **112**(1), e30–e38.
- Hasle H and Boldsen JL** (1991) Childhood conditions and adult height. *Journal of Biosocial Science* **23**(01), 107–112.
- Howe LD, Tilling K, Galobardes B, Smith GD, Gunnell D and Lawlor DA** (2012) Socioeconomic differences in childhood growth trajectories: at what age do height inequalities emerge? *Journal of Epidemiology and Community Health* **66**(2), 143–148.
- Jelenkovic A, Sund R, Hur Y-M, Yokoyama Y, Hjelmborg J, Möller S et al.** (2016) Genetic and environmental influences on height from infancy to early adulthood: an individual-based pooled analysis of 45 twin cohorts. *Scientific Reports* **6**(1), 28496.
- Jousilahti P, Tuomilehto J, Vartiainen E, Eriksson J and Puska P** (2000). Relation of adult height to cause-specific and total mortality: a prospective follow-up study of 31, 199 middle-aged men and women in Finland. *American Journal of Epidemiology* **151**(11), 1112–1120.
- Kalichman SC, Benotsch E, Suarez T, Catz S, Miller J and Rompa D** (2000) Health literacy and health-related knowledge among persons living with HIV/AIDS. *American Journal of Preventive Medicine* **18**(4), 325–331.
- Karlsen S and Nazroo JY** (2002) Relation between racial discrimination, social class, and health among ethnic minority groups. *American Journal of Public Health* **92**(4), 624–631.
- Kickbusch IS** (2001) Health literacy: addressing the health and education divide. *Health Promotion International* **16**(3), 289–297.
- King L, Hamm P and Stuckler D** (2009) Rapid large-scale privatization and death rates in ex-communist countries: an analysis of stress-related and health system mechanisms. *International Journal of Health Services* **39**(3), 461–489.
- Kirsch I** (1993) *Adult Literacy in America: A First Look at the Results of the National Adult Literacy Survey*. US Government Printing Office, Superintendent of Documents, Washington, DC, 20402 (Stock No. 065-000-00588-3).
- Ksenofontova N** (1994) Trends in infant mortality in the USSR. In Lutz W, Scherbov S and Volkov A (eds) *Demographic Trends and Patterns in the Soviet Union Before 1991*. Routledge, London/New York, pp. 359–378.
- Lindeboom M, Llena-Nozal A and van der Klaauw B** (2009) Parental education and child health: evidence from a schooling reform. *Journal of Health Economics* **28**(1), 109–131.

- Lu Y (2008) Test of the “healthy migrant hypothesis”: a longitudinal analysis of health selectivity of internal migration in Indonesia. *Social Science & Medicine* 67(8), 1331–1339.
- Meyer HE and Selmer R (1999) Income, educational level and body height. *Annals of Human Biology* 26(3), 219–227.
- Moestue H and Huttly S (2008) Adult education and child nutrition: the role of family and community. *Journal of Epidemiology & Community Health* 62(2), 153–159.
- Moltchanova E and Eriksson JG (2015) Longitudinal changes in maternal and neonatal anthropometrics: a case study of the Helsinki Birth Cohort, 1934–1944. *Journal of Developmental Origins of Health and Disease* 6(4), 285–290.
- Nazroo JY (1998) Genetic, cultural or socio-economic vulnerability? Explaining ethnic inequalities in health. *Sociology of Health & Illness* 20(5), 710–730.
- Nutbeam D (2000) Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promotion International* 15(3), 259–267.
- Paasche-Orlow M and Wolf M (2007) The causal pathways linking health literacy to health outcomes. *American Journal of Health Behavior* 31 (Supplement 1), S19–S26.
- Paciorek CJ, Stevens GA, Finucane MM and Ezzati M (2013) Children’s height and weight in rural and urban populations in low-income and middle-income countries: a systematic analysis of population-representative data. *The Lancet Global Health* 1(5), e300–e309.
- Parker RM, Baker DW, Williams MV and Nurss JR (1995) The test of functional health literacy in adults. *Journal of General Internal Medicine* 10(10), 537–541.
- Pawlowski B, Dunbar M and Lipowicz A (2000) Tall men have more reproductive success. *Nature* 403, 156–157.
- Peck MN and Lundberg O (1995) Short stature as an effect of economic and social conditions in childhood. *Social Science & Medicine* 41(5), 733–738.
- Perola M (2011) Genome-wide association approaches for identifying loci for human height genes. *Best Practice & Research Clinical Endocrinology & Metabolism* 25(1), 19–23.
- Petranović MZ, Tomas Ž, Narančić SN, Škarić-Jurić T, Veček A and Miličić J (2014) A six decades long follow-up on body size in adolescents from Zagreb, Croatia (1951–2010). *Economics & Human Biology* 13, 155–164.
- Quintana-Domeque C, Bozzoli C and Bosch M (2011) Infant mortality and adult stature in Spain. *Social Science & Medicine* 72(11), 1893–1903.
- Sanders LM, Federico S, Klass P, Abrams MA and Dreyer B (2009) Literacy and child health. *Archives of Pediatrics & Adolescent Medicine* 163(2), 131–140.
- Sanders LM, Zacur G, Haecker T and Klass P (2004) Number of children’s books in the home: an indicator of parent health literacy. *Ambulatory Pediatrics* 4(5), 424–428.
- Schultz P (2002) Wage gains associated with height as a form of health human capital. *American Economic Review* 92(2), 349–353.
- Sikora J, Evans MDR and Kelley J (2019) Scholarly culture: how books in adolescence enhance adult literacy, numeracy and technology skills in 31 societies. *Social Science Research* 77, 1–15.
- Silventoinen K (2003) Determinants of variation in adult body height. *Journal of Biosocial Science* 35(2), 263–285.
- Silventoinen K, Lahelma E and Rahkonen O (1999) Social background, adult body-height and health. *International Journal of Epidemiology* 28(5), 911–918.
- Silventoinen K, Sammalisto S, Perola M, Boomsma DI, Cornes BK, Davis C *et al.* (2003) Heritability of adult body height: a comparative study of twin cohorts in eight countries. *Twin Research* 6(5), 399–408.
- Slomczynski KM and Tomescu-Dubrow I (2006) Representation of Post-Communist European countries in cross-national public opinion surveys. *Problems of Post-Communism* 53(4), 42–52.
- Smith GD, Hart C, Upton M, Hole D, Gillis C, Watt G and Hawthorne V (2000) Height and risk of death among men and women: aetiological implications of associations with cardiorespiratory disease and cancer mortality. *Journal of Epidemiology & Community Health* 54(2), 97–103.
- Smith SK, Dixon A, Trevena L, Nutbeam D and McCaffery KJ (2009) Exploring patient involvement in healthcare decision making across different education and functional health literacy groups. *Social Science & Medicine* 69(12), 1805–1812.
- Tao H-L, Yin C-C (2016) Height, partners and offspring: evidence from Taiwan. *Journal of Biosocial Science* 48(5), 593–615.
- Thomas D (1994) Like father, like son; like mother, like daughter: parental resources and child height. *Journal of Human Resources* 29, 950–988.
- Tomopoulos S, Dreyer BP, Tamis-LeMonda C, Flynn V, Rovira I, Tineo W and Mendelsohn AL (2006) Books, toys, parent–child interaction, and development in young Latino children. *Ambulatory Pediatrics* 6(2), 72–78.
- Visscher PM, McEvoy B and Yang J (2010) From Galton to GWAS: quantitative genetics of human height. *Genetics Research* 92(5–6), 371–379.
- Visscher PM, Medland SE, Ferreira MAR, Morley KI, Zhu G, Cornes BK *et al.* (2006) Assumption-free estimation of heritability from genome-wide identity-by-descent sharing between full siblings. *PLoS Genetics* 2(3), e41.

- Woods-Townsend K, Leat H, Bay J, Bagust L, Davey H, Lovelock D et al.** (2018) LifeLab Southampton: a programme to engage adolescents with DOHaD concepts as a tool for increasing health literacy in teenagers – a pilot cluster-randomized control trial. *Journal of Developmental Origins of Health and Disease* **9**(5), 475–480.
- Zielińska D** (1991) Is there a selective rural–urban migration in respect to height and weight? *American Journal of Human Biology* **3**(4), 363–368.
- Zurawiecka M, Marchewka J and Wronka I** (2019) Fluctuating body asymmetry in young Polish women in relation to childhood socioeconomic status. *Journal of Biosocial Science* **51**(5), 775–783.