

# EUR Research Information Portal

## The global epidemiology of hepatitis E virus infection: A systematic review and meta-analysis

**Published in:**  
Liver International

**Publication status and date:**  
Published: 01/01/2020

**DOI (link to publisher):**  
[10.1111/liv.14468](https://doi.org/10.1111/liv.14468)

**Document Version**  
Publisher's PDF, also known as Version of record

### Citation for the published version (APA):

Li, P., Liu, J., Li, Y., Su, J., Ma, ZR., Bramer, WM., Cao, W., de Man, R., Peppelenbosch, M., & Pan, Q. (2020). The global epidemiology of hepatitis E virus infection: A systematic review and meta-analysis. *Liver International*, *40*(7), 1516-1528.  
<https://doi.org/10.1111/liv.14468>

[Link to publication on the EUR Research Information Portal](#)

### Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.


In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

### Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: [openaccess.library@eur.nl](mailto:openaccess.library@eur.nl). Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.



# The global epidemiology of hepatitis E virus infection: A systematic review and meta-analysis

Pengfei Li<sup>1</sup> | Jiaye Liu<sup>1</sup> | Yang Li<sup>1</sup> | Junhong Su<sup>1,2</sup> | Zhongren Ma<sup>2</sup> |  
Wichor M. Bramer<sup>3</sup> | Wanlu Cao<sup>1</sup> | Robert A. de Man<sup>1</sup> | Maikel P. Peppelenbosch<sup>1</sup> |  
Qiuwei Pan<sup>1,2</sup> 

<sup>1</sup>Department of Gastroenterology and Hepatology, Erasmus MC-University Medical Center, Rotterdam, The Netherlands

<sup>2</sup>Biomedical Research Center, Northwest Minzu University, Lanzhou, China

<sup>3</sup>Medical Library, Erasmus MC-University Medical Center, Rotterdam, The Netherlands

## Correspondence

Jiaye Liu, Department of Gastroenterology and Hepatology, Erasmus MC-University Medical Center, Rotterdam, The Netherlands.

Email: j.liu.2@erasmusmc.nl

Qiuwei Pan, Department of Gastroenterology and Hepatology, Erasmus MC, room Na-1011, 'sGravendijkwal 230, NL-3015 CE Rotterdam, The Netherlands. Email: q.pan@erasmusmc.nl

## Funding information

This research is supported by the KWF (Dutch Cancer Society) Young Investigator grant (No. 10140) and a VIDJ grant (No. 91719300) from the Netherlands Organisation for Scientific Research (NWO) to Q. Pan, the Changjiang Scholars and Innovative Research Team in University (No. IRT\_17R88) to Z. Ma, and the China Scholarship Council for funding PhD fellowship to P. Li (201808370170), J. Liu (201606240079), Y. Li (201703250073), J. Su (201708535017), W. Cao (201307060013).

**Handling Editor:** Benjamin Maasoumy

## Abstract

**Background and aims:** Hepatitis E virus (HEV), as an emerging zoonotic pathogen, is a leading cause of acute viral hepatitis worldwide, with a high risk of developing chronic infection in immunocompromised patients. However, the global epidemiology of HEV infection has not been comprehensively assessed. This study aims to map the global prevalence and identify the risk factors of HEV infection by performing a systematic review and meta-analysis.

**Methods:** A systematic searching of articles published in Medline, Embase, Web of science, Cochrane and Google scholar databases till July 2019 was conducted to identify studies with HEV prevalence data. Pooled prevalence among different countries and continents was estimated. HEV IgG seroprevalence of subgroups was compared and risk factors for HEV infection were evaluated using odd ratios (OR).

**Results:** We identified 419 related studies which comprised of 1 519 872 individuals. A total of 1 099 717 participants pooled from 287 studies of general population estimated a global anti-HEV IgG seroprevalence of 12.47% (95% CI 10.42-14.67;  $I^2 = 100\%$ ). Notably, the use of ELISA kits from different manufacturers has a substantial impact on the global estimation of anti-HEV IgG seroprevalence. The pooled estimate of anti-HEV IgM seroprevalence based on 98 studies is 1.47% (95% CI 1.14-1.85;  $I^2 = 99\%$ ). The overall estimate of HEV viral RNA-positive rate in general population is 0.20% (95% CI 0.15-0.25;  $I^2 = 98\%$ ). Consumption of raw meat ( $P = .0001$ ), exposure to soil ( $P < .0001$ ), blood transfusion ( $P = .0138$ ), travelling to endemic areas ( $P = .0244$ ), contacting with dogs ( $P = .0416$ ), living in rural areas ( $P = .0349$ ) and receiving education less than elementary school ( $P < .0001$ ) were identified as risk factors for anti-HEV IgG positivity.

**Conclusions:** Globally, approximately 939 million corresponding to 1 in 8 individuals have ever experienced HEV infection. 15-110 million individuals have recent or

**Abbreviations:** 95% CI, 95% confidential interval; GT, genotype; HEV, hepatitis E virus; HIV, human immunodeficiency virus; IDU, intravenous drug use; MSM, man having sex with man; OR, odd ratio.

Pengfei Li and Jiaye Liu contributed equally.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 John Wiley & Sons A/S. Published by John Wiley & Sons Ltd

ongoing HEV infection. Our study highlights the substantial burden of HEV infection and calls for increasing routine screening and preventive measures.

#### KEYWORDS

epidemiology, hepatitis E virus, prevalence, risk factors, seroprevalence

## 1 | INTRODUCTION

Hepatitis E virus (HEV) as a positive-sense single-stranded RNA virus is a leading cause of acute viral hepatitis worldwide. The infection is usually asymptomatic or self-limiting in the general population. However, acute infection in pregnant women may cause severe clinical outcomes, including fulminant hepatic failure with high mortality rate reaching up to 20%-30%.<sup>1</sup> These patients are mostly from resource-limited regions. In European countries, HEV infection has been frequently reported to bear high risk of developing into chronic hepatitis in immunocompromised individuals, in particular organ transplant patients.<sup>2,3</sup> Thus, HEV is truly imposing a global health burden in both developing and developed countries.

Currently, eight distinct genotypes (GTs) of HEV have been classified.<sup>4</sup> GT 1-4 are known to be the main threat to humans. GT 1 and GT 2 are restricted to human and mainly transmit through contaminated water causing acute hepatitis. GT 3 and GT 4 are zoonotic and have been identified in a wide spectrum of hosts, including human, swine, wild boar, goat, cattle, deer, camel and yak.<sup>5</sup> Both GT 3 and GT 4 can cause chronic infection in organ transplant patients,<sup>2,6</sup> and consumption of raw or undercooked animal meat has been recognized as the main routes of causing sporadic cases in developed countries.<sup>7</sup> In fact, the host range of HEV is ever expanding and the implications of the rare GTs and the newly discovered strains in human health remain largely uncertain.<sup>7</sup> This further complicates the transmission and the risk of HEV infection. In addition to the classical waterborne and foodborne transmission routes, blood transfusion-mediated transmission has been reported in organ transplant patients.<sup>8</sup> Person-to-person transmission has also been proposed.<sup>9</sup> Intriguingly, recent evidence has indicated that pet animals including dogs, cats, rabbits and horses might be accidental hosts for HEV and constitute a potential source for transmitting to human.<sup>10,11</sup> Thus, there is an urgent need to comprehensively understand the risks for HEV infection, in order to devise preventive measures.

Globally, it has been roughly estimated that one-third of the population are living in HEV endemic areas.<sup>12</sup> More recently, substantial efforts have been dedicated to systematically evaluate HEV prevalence in different continents (eg the Americas and Europe),<sup>13,14</sup> different countries (eg industrialized countries, China, Iran, Brazil and Somalia)<sup>15-17</sup> and special populations or settings (eg blood donors, swine workers and outbreak setting).<sup>18-20</sup> Most of these studies are based on seropositivity of anti-HEV IgG antibody. Anti-HEV IgG antibody developed post-infection usually persists for many years, and is thus regarded as a marker for past infection.<sup>21,22</sup> In contrast, anti-HEV IgM antibody is short-lived up to a few months, thus

#### Key points

- This meta-analysis reports the latest estimation that approximately 939 million of the global population have ever experienced hepatitis E virus (HEV) infection, and 15-110 million individuals have recent or ongoing infection.
- These findings indicate that HEV infection has emerged as a global health burden requiring implementation of specific control measures.

considered as evidence of recent or current infection. Detection of HEV RNA is a bona fide marker for active ongoing infection. In this study, we aimed to systematically estimate the global burden of HEV infection. More specifically, we have mapped the global prevalence of past, recent and ongoing HEV infection and evaluated the key risk factors of infection.

## 2 | MATERIALS AND METHODS

### 2.1 | Data sources and searches

A systematic search was conducted in Medline, Embase, Web of science, Cochrane CENTRAL and Google scholar. Databases were searched for articles in the English language from inception until July 2019. All searches from database were performed by a biomedical information specialist of the medical library, with an exhaustive set of search terms related to hepatitis E virus infection and epidemiology (the full search strategies are provided in the Supporting Information S1). This study is reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis.<sup>23</sup> No institutional review board approval was required for this meta-analysis because our study only included data which had been published previously.

### 2.2 | Study selection

Studies were included if they met the following criteria: (a) Studies which contained data about seroprevalence of anti-HEV IgG, anti-HEV IgM or HEV RNA positivity, (b) Studies contained mixed population were excluded unless they clearly and explicitly reported the

prevalence for each group, (c) Studies contained information of risk factors related to HEV infection and (d) Studies which focused on the HEV prevalence in human beings.

Studies were excluded if they met the following criteria: (a) Studies are systematic review, meta-analysis, case reports, perspectives and abstracts, (b) None human studies, (c) No primary data or incomplete data, (d) Duplicate data, (e) Studies with <50 individuals were excluded in order to decrease bias caused by small population and (f) Studies concerning about HEV outbreaks, since the prevalence and outcome of HEV infection in these studies would dramatically differ from those of the general population.

Two reviewers (PL and JL) worked independently to determine whether a study met inclusion criteria, abstracted information to assess the methodological validity of each candidate study and extracted data with structured data collection forms. The reviewers resolved discrepancies by jointly reviewing the study in question. If no consensus was reached, a third reviewer (QP), unaware of prior determinations, functioned as an arbiter.

### 2.3 | Data extraction and quality assessment

Eligible studies were further divided into three study populations: general population, occupational population and special population. General population included people without apparent risk factors and could be comprised of blood donors, pregnant women, healthy individuals and hospital attendants. For general population, individuals were further divided into subgroups by gender, different age ranges, study period (1993-2006 or 2007-2019), country development classification (developing and developed countries), gross national income classification of each country (high, upper middle, lower middle and low income) and ELISA kit manufacturers. More importantly, OR analysis of anti-HEV IgG seropositivity was conducted in possible risk factors including living area (urban or rural), consumption of raw meat, exposure to soil, contacting with cat or dog, education level (elementary school or above elementary school), intravascular drug use (IDU), water source (tap, well or river), man having sex with man (MSM), transfusion history and traveling history to endemic areas. Occupational population represents people who had been in frequent contact with pigs or pig products, including veterinarians, swine workers, slaughterhouse workers and pork sellers. Special populations are further categorized into four groups as followings: patients with acute hepatitis (caused by hepatitis B virus, hepatitis C virus or other unknown hepatitis), individuals with human immunodeficiency virus (HIV) infection, people who underwent haemodialysis and organ transplant recipients. Two independent reviewers (PL and JY) extracted data, with discrepancies and disagreements resolved by discussion. We extracted data on first author, country, continent, publication date, anti-HEV IgG prevalence, anti-HEV IgM prevalence, HEV RNA positivity, subgroup information of anti-HEV IgG prevalence and HEV-related risk factors using data extracting forms. When multiple publications were identified that reported on the same populations and outcomes, only

the most representative and comprehensive study was included for further meta-analysis in order to avoid duplicate data. The quality of studies was assessed using the Joanna Briggs Institute checklist for prevalence studies, which enabled assessment of included studies in relation to risk of bias, rigour and transparency.<sup>24</sup> Studies scoring 1-3 were defined as low quality, 4-6 as average quality and 7-9 as high quality (Table S1). Studies were not excluded on the basis of their quality score to increase transparency and to ensure all available evidence in this area was reported.

### 2.4 | Statistics analysis

After checking for consistency, the Metaprop module in the R-3.5.3 statistical software package was used for meta-analysis. A 95% confidence interval (95% CI) was estimated using Wilson score method, and pooled seroprevalence was calculated with the DerSimonian-Laird random effects model with Freeman-Tukey double arcsine transformation. Heterogeneity across the included studies was assessed using the Cochran Q statistics and  $I^2$  statistics, with  $I^2$  statistics 25%-50%, 50%-75% and >75% considered as mild, moderate and severe heterogeneity respectively. When heterogeneity was higher than 50%, a random effect model will be used. ORs were used to report the risk factors for HEV infection. ORs and their 95% CI were extracted directly from studies when available, with adjusted ORs extracted preferentially over unadjusted ORs. If included studies did not report ORs, crude ORs were calculated from extracted data. We then pooled the ORs using the DerSimonian and Laird random effect models, with the heterogeneity estimated from the Mantel-Haenszel model. Funnel plots and Egger regression test were used to assess potential publication biases. Additionally, we performed sensitivity analyses using "metainf" in a random model to investigate the effects of population source and potentially unrepresentative samples. The estimated prevalence of anti-HEV IgG, IgM and HEV RNA infection was based on the global population of 7 530 000 000 on 20 July 2019 (<https://population.io>).

## 3 | RESULTS

### 3.1 | Global prevalence of HEV infection

Our search returned 8153 records, of which 419 met the inclusion criteria (Figure 1). In total, participants from 302 studies related to general population, and 287 studies were pooled to estimate a global anti-HEV IgG seroprevalence of 12.47% (1 099 717 individuals included; 95% CI 10.42-14.67;  $I^2 = 100%$ ; Figure 2A; Figure S1). The pooled estimate of anti-HEV IgM seroprevalence based on 98 studies is 1.47% (479 001 individuals; 95% CI 1.14-1.85;  $I^2 = 99%$ ; Figure 2B; Figure S2). The overall estimate of HEV RNA-positive rate in the general population is 0.20% (3 444 752; 95% CI 0.15-0.25;  $I^2 = 98%$ ; Figure 2C; Figure S3). We also stratified data to estimate the HEV prevalence in 75 countries among six continents (excluding





FIGURE 1 Study selection

Antarctica). The highest anti-HEV IgG seropositivity rate was found in Africa (22 377; 21.76%, 95% CI 13.05-31.98;  $I^2 = 100\%$ ), followed by Asia (681 373; 15.80%, 95% CI 13.29-18.49;  $I^2 = 100\%$ ), Europe (132 419; 9.31%, 95% CI 7.35-11.48;  $I^2 = 99\%$ ), North America (71 989; 8.05%, 95% CI 5.47-11.09;  $I^2 = 99\%$ ), South America (14 586; 7.28%, 95% CI 4.83-10.19;  $I^2 = 97\%$ ) and Oceania (1563; 5.99%, 95% CI 1.22-14.03;  $I^2 = 96\%$ ; Figure S4). Besides, the anti-HEV IgM seroprevalence was 3.09% (5001; 95% CI 1.49-5.24;  $I^2 = 93\%$ ), 1.86% (141 565; 95% CI 1.34-2.46;  $I^2 = 98\%$ ), 0.79% (146 322; 95% CI 0.30-1.51;  $I^2 = 99\%$ ), 0.22% (12 197; 95% CI 0.00-0.74;  $I^2 = 91\%$ ) and 2.43% (2680; 95% CI 0.43-6.00;  $I^2 = 96\%$ ) for Africa, Asia, Europe, North America and South America respectively (Figure S5). In addition, the HEV RNA positivity rate was 0.00% (278; 95% CI 0.00-0.35), 0.93% (727 744; 95% CI 0.48-1.52;  $I^2 = 99\%$ ), 0.08% (2 441 774; 95% CI 0.05-0.11;  $I^2 = 95\%$ ), 0.00% (34 761; 95% CI 0.00-0.02;  $I^2 = 45\%$ ), 0.00% (74 131; 95% CI 0.00-0.01) and 0.18% (1054; 95% CI 0.00-1.36;  $I^2 = 81\%$ ) for Africa, Asia, Europe, North America, Oceania and South America respectively (Figure S6). HEV prevalence varies substantially among countries, from 0.25% (Tanzania, 95% CI 0.00-0.97) to 74.76% (South Sudan, 95% CI 68.61-80.44) of anti-HEV IgG, from 0.00% (Mongolia, 95% CI 0.00-0.08; Bulgaria, 95% CI 0.00-0.13) to 19.83% (United Arab, 95% CI 16.35-23.56) of anti-HEV IgM and from 0.00% (Benin, Malawi, Australia, Canada, Brazil) to 6.75% (France, 95% CI 0.14-22.04) of HEV RNA positivity (Table 1; Figures S1-S3).

We also collected data of HEV GTs, with the finding that HEV GT 1 infection occasionally occurred in China and frequently in India, and GT 3 was widely distributed in European countries. GT 3 was also prevalent in Japan and Korea, whereas GT 4 infection mainly emerged in China (Figure 3; Table S2). Based on our comprehensive estimates, approximately 938 991 000 individuals corresponding to 1/8 of the global population have ever experienced HEV infection based on anti-HEV IgG positivity. Importantly, we estimated approximately 110 691 000 global individuals with current or recent HEV infection and 15 060 000 individuals with ongoing HEV infection based on anti-HEV IgM or viral RNA positivity respectively.

We next performed subgroup analysis of anti-HEV IgG positivity rate in general population. General population of six different continents were further divided into seven age groups, including age range of 0-9, 10-19, 20-29, 30-39, 40-49, 50-59 and above 60-year-old. The corresponding pooled anti-HEV IgG-positive rates are 7.73% (6977 individuals included; 95% CI 2.29-16.02;  $I^2 = 99\%$ ), 9.03% (14 452 individuals; 95% CI 3.78-16.25;  $I^2 = 99\%$ ), 10.78% (33 365; 95% CI 7.44-14.64;  $I^2 = 99\%$ ), 14.17% (23 217; 95% CI 10.27-18.57;  $I^2 = 99\%$ ), 21.53% (21 324; 95% CI 16.82-26.65;  $I^2 = 99\%$ ), 24.48% (17 474; 95% CI 18.56-30.93;  $I^2 = 99\%$ ) and 27.47% (23 924; 95% CI 21.07-34.36;  $I^2 = 99\%$ ) respectively (Figure 4; Figures S7 and S8). The positive rate is slightly higher in male (129 569; 13.39%, 95% CI 11.34-15.59;  $I^2 = 99\%$ ) compared to female (120 264; 12.25%,

**FIGURE 2** (A) The global seroprevalence of anti-HEV IgG antibody (B) The global seroprevalence of anti-HEV IgM antibody (C) The global prevalence of HEV RNA positivity



95% CI 10.05-14.63;  $I^2 = 99\%$ ; Figure 4; Figure S9). To clarify the HEV prevalence among regions with different levels of economic development, we firstly calculated the anti-HEV IgG prevalence in high-income countries, upper middle income countries, lower middle income countries and low income countries. We estimated the anti-HEV IgG positivity of 8.84% (424 905; 95% CI 6.79-11.14;  $I^2 = 100\%$ ) in high-income countries, 12.79% (618 638; 95% CI 10.81-14.92;  $I^2 = 100\%$ ) in upper middle income countries, 19.04% (40 593;

95% CI 13.25%-25.60%;  $I^2 = 100\%$ ) in lower middle income countries and 30.44% (5781; 95% CI 16.60-46.39;  $I^2 = 99\%$ ) in low income countries (Figure 4; Figure S10). The pooled estimate of anti-HEV IgG seroprevalence was 14.83% (689 452; 95% CI 12.98-16.77;  $I^2 = 100\%$ ) in developing countries compared to 8.59% (401 513; 95% CI 6.46-10.99;  $I^2 = 100\%$ ) in developed countries (Figure 4; Figure S11). Of the global HEV prevalence during the period of 1993-2019, we estimated anti-HEV IgG-positive rate of 9.43% (79 998; 95% CI

TABLE 1 Hepatitis E virus (HEV) prevalence in general population

Continent	Country	Anti-HEV IgG						Anti-HEV IgM Anti-HEV IgM						HEV RNA					
		No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	No. of studies	Events	Tested (n)	Prevalence (%)	95% CI			
Asia	China	42	200 221	579 696	22.68	19.67; 25.83	18	1186	88 587	1.76	1.29; 2.31	5	106	56 319	0.41	0.07; 1.06			
	Thailand	4	121	2632	4.82	0.30; 14.31	2	5	2057	0.12	0.00; 1.05	1	26	30 115	0.09	0.06; 0.12			
	Israel	3	133	7115	3.36	0.19; 10.22	..	..	..	..	..	..	..	..	..	..	..		
	Saudi Arabia	3	428	2911	15.41	10.77; 20.70	1	39	900	4.33	3.10; 5.76	..	..	..	..	..	..		
	Japan	8	1901	40 936	4.26	3.27; 5.37	5	32	27 478	0.30	0.06; 0.73	2	41	623 325	0.05	0.00; 0.29			
	Jordan	1	139	450	30.89	26.71; 35.23	..	..	..	..	..	..	..	..	..	..	..		
	Kazakhstan	1	11	199	5.53	2.79; 9.12	..	..	..	..	..	..	..	..	..	..	..		
	Korea	3	328	3969	9.30	4.87; 14.96	1	9	1030	0.87	0.40; 1.53	..	..	..	..	..	..		
	Vietnam	2	44	833	7.25	0.39; 21.55	1	1	187	0.53	0.00; 2.08	..	..	..	..	..	..		
	Yemen	1	38	356	10.67	7.68; 14.09	..	..	..	..	..	..	..	..	..	..	..		
	Bangladesh	3	509	1707	36.87	17.16; 59.20	1	20	273	7.33	4.54; 10.71	..	..	..	..	..	..		
	India	7	3160	14 136	27.15	19.31; 35.78	5	348	5354	10.18	2.08; 23.38	4	510	8462	6.59	0.22; 20.71			
	Indonesia	2	50	858	5.83	4.36; 7.49	..	..	..	..	..	..	..	..	..	..	..		
	Mongolia	3	36	1486	2.70	0.06; 9.08	2	0	1237	0	0.00; 0.08	..	..	..	..	..	..		
	Iran	20	1208	12 547	8.98	5.74; 12.86	5	38	4148	0.80	0.19; 1.82	2	7	2031	0.24	0.00; 2.16			
	Cambodia	3	1074	3173	28.93	14.46; 46.04	2	55	2305	1.31	0.01; 4.74	2	3	1169	0.25	0.05; 0.63			
	Singapore	1	30	219	13.70	9.48; 18.56	..	..	..	..	..	..	..	..	..	..	..		
	Laos	1	38	210	18.10	13.20; 23.58	..	..	..	..	..	..	..	..	..	..	..		
	Nepal	2	1344	2602	59.19	26.14; 88.13	1	54	1686	3.20	2.42; 4.10	..	..	..	..	..	..		
Qatar	1	1198	5854	20.46	19.44; 21.51	1	34	5854	0.58	0.40; 0.79	1	4	5854	0.07	0.02; 0.15				
United Arab Emirates	..	..	..	..	..	1	93	469	19.83	16.35; 23.56	1	28	469	5.97	4.01; 8.29				
Total	111	212 011	681 373	15.80	13.29; 18.49	46	1920	141 565	1.86	1.34; 2.46	18	725	727 744	0.93	0.48; 1.52				
Africa	Nigeria	2	56	588	10.17	5.32; 16.35	2	17	927	2.83	0.00; 17.56	..	..	..	..	..	..		
	Burkina Faso	1	56	178	31.46	24.86; 38.46	..	..	..	..	..	..	..	..	..	..	..		
	Burundi	1	18	129	13.95	8.54; 20.44	..	..	..	..	..	..	..	..	..	..	..		
	Djibouti	1	14	112	12.50	7.05; 19.23	..	..	..	..	..	..	..	..	..	..	..		
	Egypt	6	6670	14 052	42.43	20.40; 66.19	1	6	100	6.00	2.22; 11.47	..	..	..	..	..	..		
	Ethiopia	2	481	1232	37.09	26.85; 47.95	2	10	1232	0.80	0.38; 1.37	..	..	..	..	..	..		
	Gabon	2	135	1083	10.23	4.02; 18.87	..	..	..	..	..	..	..	..	..	..	..		
	Ghana	3	75	789	8.74	4.39; 14.38	3	44	789	5.87	0.06; 20.10	..	..	..	..	..	..		
	Benin	1	62	278	22.30	17.61; 27.38	1	7	278	2.52	1.01; 4.68	1	0	278	0	0.00; 1.32			
	Malawi	1	80	397	20.15	16.36; 24.23	..	..	..	..	..	..	..	..	..	..	..		

(Continues)

TABLE 1 (Continued)

Continent	Country	Anti-HEV IgG					Anti-HEV IgM Anti-HEV IgM					HEV RNA					
		No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	
		South Africa	3	418	2243	16.00	1.55; 41.31	1	16	782	2.05	1.17; 3.16	..	..	..	..	..
South Sudan	1	154	206	74.76	68.61; 80.44	1	9	206	4.37	2.01; 7.58	..	..	..	..	..	..	
Tanzania	1	1	403	0.25	0.00; 0.97	..	..	..	..	..	..	..	..	..	..	..	
Tunisia	1	37	687	5.39	3.82; 7.20	1	20	687	2.91	1.79; 4.30	..	..	..	..	..	..	
Total	26	8257	22 377	21.76	13.05; 31.98	12	129	5001	3.09	1.49; 5.24	1	0	278	0	0.00; 0.95	..	
Europe	Finland	1	37	385	9.61	6.87; 12.75	1	1	385	0.26	0.00; 1.02	..	..	..	..	..	..
	France	8	3978	17 778	14.51	6.80; 24.49	5	230	15 027	1.16	0.30; 2.57	3	32	917	6.75	0.14; 22.04	..
	Germany	11	2146	11 045	14.35	6.69; 24.30	2	60	2183	2.86	0.00; 23.32	2	16	17 144	0.14	0.02; 0.37	..
	Moldova	1	63	255	24.71	19.62; 30.18	..	..	..	..	..	..	..	..	..	..	..
	Greece	1	25	265	9.43	6.22; 13.24	..	..	..	..	..	..	..	..	..	..	..
	Turkey	10	225	4656	4.93	2.68; 7.81	..	..	..	..	..	..	..	..	..	..	..
	UK	7	706	9672	5.60	3.11; 8.76	3	25	96 341	0.36	0.00; 1.66	4	600	2 201 609	0.02	0.01; 0.04	..
	Iceland	1	6	291	2.06	0.75; 4.01	..	..	..	..	..	..	..	..	..	..	..
	Ireland	2	73	1274	6.17	3.92; 8.90	1	2	1076	0.19	0.02; 0.53	1	10	24 985	0.04	0.02; 0.07	..
	Italy	14	1690	19 488	7.28	4.54; 10.60	4	49	10 559	0.44	0.19; 0.78	2	12	10 363	0.83	0.00; 7.27	..
	Czech	1	13	230	5.65	3.05; 9.00	..	..	..	..	..	..	..	..	..	..	..
	Denmark	2	296	1459	11.09	0.00; 57.44	..	..	..	..	..	..	..	..	..	..	..
	Austria	2	306	2200	13.91	12.49; 15.38	..	..	..	..	..	..	..	..	..	..	..
	Bulgaria	1	67	741	9.04	7.08; 11.21	1	0	741	0	0.00; 0.13	..	..	..	..	..	..
	Montenegro	1	24	400	6.00	3.89; 8.53	00B7.	..	..	..	..	..	..	..	..	..	..
	Netherlands	8	3521	25 786	16.07	6.09; 29.62	3	491	13 503	2.00	0.05; 6.63	4	96	67 041	0.40	0.03; 1.18	..
	Norway	1	177	1263	14.01	12.16; 15.98	..	..	..	..	..	..	..	..	..	..	..
	Poland	4	1561	4497	14.17	2.07; 34.48	3	41	3470	1.02	0.52; 1.67	1	10	12 664	0.08	0.04; 0.14	..
	Portugal	3	380	2812	8.72	2.20; 18.99	1	8	1656	0.48	0.21; 0.87	..	..	..	..	..	..
	Romania	1	22	148	14.86	9.61; 21.03	..	..	..	..	..	..	..	..	..	..	..
	Russia	1	62	341	18.18	14.27; 22.45	1	2	341	0.59	0.06; 1.67	..	..	..	..	..	..
	San Marino	1	33	2233	1.48	1.02; 2.02	..	..	..	..	..	..	..	..	..	..	..
	Serbia	2	50	726	8.46	0.95; 22.35	..	..	..	..	..	..	..	..	..	..	..
	Spain	7	2282	18 534	5.08	0.85; 12.58	1	7	1040	0.67	0.27; 2.16	3	7	22 351	0.03	0.01; 0.06	..
	Sweden	2	14	205	6.51	2.40; 12.45	..	..	..	..	..	..	..	..	..	..	..
	Switzerland	4	803	5228	7.25	0.97; 18.62	..	..	..	..	..	..	..	..	..	..	..
Total	97	18 560	132 419	9.31	7.35; 11.48	26	916	146 322	0.79	0.30; 1.51	23	807	2 441 744	0.08	0.05; 0.11	..	..


(Continues)



TABLE 1 (Continued)

Continent	Country	Anti-HEV IgG				Anti-HEV IgM Anti-HEV IgM				HEV RNA						
		No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	No. of studies	Events	Tested (n)	Prevalence (%)	95% CI	No. of studies	Events	Tested (n)	Prevalence (%)	95% CI
Oceania	New Zealand	1	98	1013	9.67	7.93; 11.57	..	..	..	..	..	..	..	..	..	..
	Australia	1	17	550	3.09	1.81; 4.70	..	..	..	..	..	1	1	74 131	0	0.00; 0.01
	Total	2	115	1563	5.99	1.22; 14.03	..	..	..	..	..	1	1	74 131	0	0.00; 0.01
North America	Cuba	3	74	1827	4.89	0.70; 12.55	1	5	1149	0.44	0.14; 0.90	..	..	..	..	..
	Mexico	6	535	4977	8.97	3.75; 16.14	..	..	..	..	..	..	..	..	..	..
	USA	12	7747	60 291	9.65	5.68; 14.53	3	12	11 048	0.16	0.00; 0.78	2	2	20 768	0.01	0.00; 0.03
	Canada	2	252	4495	4.35	1.82; 7.91	..	..	..	..	..	1	0	13 993	0	0.00; 0.01
	Nicaragua	1	17	399	4.26	2.50; 6.46	..	..	..	..	..	..	..	..	..	..
	Total	24	8625	71 989	8.05	5.47; 11.09	4	17	12 197	0.22	0.00; 0.74	3	2	34 761	0	0.00; 0.02
South America	Chile	1	82	469	17.48	14.18; 21.05	..	..	..	..	..	..	..	..	..	..
	Argentina	3	145	5831	5.92	1.29; 13.60	..	..	..	..	..	..	..	..	..	..
	Venezuela	1	23	611	3.76	2.40; 5.42	1	3	611	0.49	0.09; 1.20	..	..	..	..	..
	Bolivia	5	209	1940	9.50	3.19; 18.70	1	10	574	1.74	0.83; 2.97	1	3	123	2.44	0.47; 5.89
	Brazil	10	315	4739	6.39	3.25; 10.49	4	50	1495	3.46	0.12; 11.09	2	0	931	0	0.00; 0.10
	Guyana	1	64	996	6.43	4.99; 8.03	..	..	..	..	..	..	..	..	..	..
Total	21	838	14 586	7.28	4.83; 10.19	6	63	2680	2.43	0.43; 6.00	3	3	1054	0.18	0.00; 1.36	
Multiple		6	3158	175 410	14.64	0.40; 43.67	4	18	171 236	0.15	0.00; 0.56	1	16	165 010	0.01	0.01; 0.02
	Total	287	251 564	1 099 717	12.47	10.42; 14.67	98	3063	479 001	1.47	1.14; 1.85	50	1554	3 444 752	0.20	0.15; 0.25

Multiple: studies contained more than one countries.



**FIGURE 3** Hepatitis E virus genotype distribution in our study

6.11-13.37;  $I^2 = 100\%$ ) during 1993-2006 and 13.65% (1 019 719; 95% CI 11.15-16.35;  $I^2 = 100\%$ ) during 2007-2019 (Figure 4, Figure S12).

### 3.2 | Prevalence of HEV infection in occupational population and special population


Anti-HEV IgG seroprevalence data from veterinarians, swine workers, slaughters and pork sellers were collected to estimate the overall anti-HEV seroprevalence in occupational population. Based on 43 studies with 8776 occupational individuals, the overall seropositivity of anti-HEV IgG is 24.04% (95% CI 18.55-29.99;  $I^2 = 97\%$ ; Figure S13). In total, data from 126 studies were extracted to analyse the prevalence in special populations. The overall anti-HEV IgG, anti-HEV IgM and viral RNA-positive rates are 15.43% (95% CI 12.82-18.24;  $I^2 = 98\%$ ), 3.21% (95% CI 1.77-5.06;  $I^2 = 98\%$ ) and 1.10% (95% CI 0.53-1.87;  $I^2 = 97\%$ ) respectively (Figures S14-S16). Among these special populations, patients with acute hepatitis have the highest positive rate of anti-HEV IgG (21.49%, 95% CI 12.65-31.92;  $I^2 = 99\%$ ), anti-HEV IgM (8.62%, 95% CI 4.16-14.51;  $I^2 = 99\%$ ) and viral RNA (5.57%, 95% CI 2.26-10.21;  $I^2 = 99\%$ ; Figures S17-S19). The anti-HEV IgG-positive rates in two special groups are higher than that in general population, with 16.91% (95% CI 12.64-21.67;  $I^2 = 98\%$ ) in the HIV population and 13.10% (95% CI 9.34-17.39;  $I^2 = 96\%$ ) in haemodialysis population, while it is slightly lower in organ transplant recipients with 11.68% (95% CI 7.91-16.06;  $I^2 = 97\%$ ) seropositivity (Figures S20-S27; Table S5).

### 3.3 | Risk factors of HEV

We investigated the potential risk factors for HEV in the general population (Figure S28). Significant rising effects on anti-HEV IgG seropositivity were observed in consumption of raw meat (OR 1.45, 95% CI 1.20-1.76,  $P = .0001$ ), exposure to soil (OR 1.52, 95% CI 1.24-1.86,  $P < .0001$ ), blood transfusion (OR 1.61, 95% CI 1.10-2.36,  $P = .0138$ ), travelling to endemic areas (OR 1.39, 95% CI 1.04-1.84,  $P = .0244$ ), contacting with dogs (OR 1.45, 95% CI 1.01-2.07,  $P = .0416$ ), living in rural areas (OR 0.80, 95% CI 0.65-0.98,  $P = .0349$ ) and receiving education less than elementary school (OR 1.71, 95% CI 1.41-2.07,  $P < .0001$ ). No statistically significant differences were identified for anti-HEV IgG positivity in respect to different water source ( $P = .0909$ ), IDU experience ( $P = .4321$ ), MSM experience ( $P = .5576$ ) and contacting with cats ( $P = .4791$ ; Figure S29-S39). Sensitivity analysis detected no study having an obvious effect influence to the pooled estimates of HEV prevalence in the general population (Table S3).

### 3.4 | Anti-HEV IgG detection rate of different ELISA kits

We finally analysed the detection rates of the ELISA kits from six manufacturers. The detection rates of anti-HEV IgG seropositivity vary dramatically, with the highest of Wantai assay (20.72%, 95% CI 2.07-9.84;  $I^2 = 100\%$ ) followed by MP Diagnostics (10.75%, 95% CI 4.55-19.15;  $I^2 = 100\%$ ), Dia.Pro (8.89%, 95% CI 6.68-11.38;  $I^2 = 97\%$ ), Mikrogen (8.60%, 95% CI 4.65-13.62;  $I^2 = 98\%$ ),



**FIGURE 4** Anti-hepatitis E virus (HEV) IgG seroprevalence among six subgroups

Genelabs Diagnostics (6.22%; 95% CI 3.42-9.77;  $I^2 = 98\%$ ) and Abbott Laboratories (5.27%, 95% CI 2.07-9.84;  $I^2 = 100\%$ ) (Figure S40; Table S4).

## 4 | DISCUSSION

It has been estimated that one-third of the global population, representing over two billion people, live in HEV endemic areas at risk of infection.<sup>12</sup> This has been widely misinterpreted as that 2.3 billion of the population have been infected with HEV.<sup>25</sup> In fact, the true burden of hepatitis E remains largely unknown.<sup>26</sup> In this study, we have systematically and comprehensively assessed the global HEV prevalence by retrieving data from 75 countries of the six continents. We estimated that 12.47% of the global population, corresponding to approximately 939 million individuals, have experienced past infection of HEV based on their seropositivity of anti-HEV IgG antibody. Africa and Asia have been previously recognized for the high prevalence of HEV.<sup>27,28</sup> Our estimates confirm the high seroprevalence rates of 21.76% and 15.80% in Africa and Asia respectively. For Europe, we estimated a prevalence rate of 9.31%,

which is substantially lower than a previous estimation of 16.90% from a meta-analysis performed in 2016.<sup>14</sup> A possible explanation for the disparity could be that they collected fewer studies and included small size populations, and thus is prone to cause more bias. In Americas, we observed a slightly higher seroprevalence rate in North (8.05%) compared to South (7.28%) America, which is consistent with the results from a recent meta-analysis.<sup>13</sup>

Of a technical note, it has been well-realized that there are substantial differences in sensitivity and specificity of the anti-HEV IgG ELISA kits from different manufacturers.<sup>29,30</sup> Our results largely agree with the literature that the Wantai assay has the highest sensitivity and has been most widely used.<sup>31</sup> Thus, the use of different anti-HEV IgG ELISA kits may partially explain the disparities in estimates among different studies, and caution should be taken when interpreting the seroprevalence rate in this respect.

The bona fide disease burden of HEV lies in the actively infected patients. The global burden caused by GT 1 and GT 2 HEV in Africa and Asia has been mathematically modelled for 2005. Among the 4.7 billion people in these regions corresponding to 72.8% of the global population in 2005, it has been estimated as 20 million incident HEV infections, 3.4 million symptomatic cases, 70 000 deaths and 3000

stillbirths.<sup>32</sup> In 2011, WHO reported 14 million symptomatic cases annually worldwide with 300 000 deaths and 5200 stillbirths.<sup>33</sup> Hypothetically, if both estimates are accurate, there would be about 10 million symptomatic cases annually from developed countries, which are mainly caused by the zoonotic GT 3 strains. This clearly disagrees with the vast majority of the current literature that we do not expect the burden in respect to symptomatic infection would be three times in developed compared to developing countries. In this study, we have estimated approximately 110 million individuals with recent/current infection based on anti-HEV IgM antibody positivity and 15 million with ongoing infection based on HEV RNA positivity. As viral RNA persists for a few weeks and anti-HEV IgM antibody for a few months,<sup>34</sup> the annually global infections are probably at a range of hundred(s) millions. However, the available data regarding anti-HEV IgM antibody or viral RNA positivity are very limited. Thus, our estimates may have biases, and we were not able to further sub-analyse regional prevalence, GT-specific burden or clinical outcome, which require future studies in these aspects.

Accumulating knowledge on HEV biology and transmission routes has facilitated the identification of risk factors for the infection. A wide range of domestic or wild animals have been recognized as reservoirs for the zoonotic strains. Consumption of uncooked meat or meat product from swine, wild boar or deer has been widely reported to cause GT 3 HEV infection in European countries.<sup>35,36</sup> As expected, consumption of raw meat is an important risk factor revealed by our meta-analysis. This is in line with previous reports that humans with occupational exposure to pigs are at a high risk of HEV infection.<sup>37,38</sup> In this study, we observed twofold higher anti-HEV seropositivity in occupational population who had frequent contact with pig or pig products compared to the general population.

The host range for HEV is ever expanding and cross-species infections commonly occur.<sup>7</sup> Intriguingly, recent evidence has indicated that companion animals including dogs, cats, rabbits and horses might be accidental hosts for HEV and might constitute a source for HEV transmission to human.<sup>10,11,39</sup> Transmission of rat HEV to human has been recently reported in Hong Kong.<sup>40</sup> Dogs and cats are the most common household pets. Previous studies have reported that seroprevalence of HEV antibodies in dogs ranges from 0.8% in the UK,<sup>41</sup> 6.79% in Brazil,<sup>42</sup> 17.8% to 36.55% in different regions of China,<sup>10,11</sup> 22.7% in India<sup>43</sup> and 56.6% in Germany.<sup>44</sup> Interestingly, when comparing with the general population, veterinarians and dog farm staff who are frequently exposed to dogs have significantly higher anti-HEV antibody positivity.<sup>10</sup> The anti-HEV seroprevalence rates in cats have been reported to be 6.28% in China,<sup>11</sup> 8.1% in Korea<sup>45</sup> and 33% in Japan.<sup>46</sup> In this study, we found that people who frequently contact with dogs have higher anti-HEV IgG seropositivity. This was not found in people who contact with cats, but the number of studies is very limited. These results call more attention to address the potential role of pets in HEV zoonotic transmission, although currently it remains unconfirmed whether pets are reservoirs, requiring further investigation.

Previous studies have indicated the differences of HEV seroprevalence between rural and urban areas.<sup>47-49</sup> We found that rural

compared to urban residents have higher risk of HEV infection. This largely agrees with our findings that high exposure to soil is also a risk factor. In addition, we observed the high risk of HEV infection in individuals with lower education levels, consistent with previous studies.<sup>50,51</sup> Conceivably, this population are more likely living in rural areas with compromised sanitation conditions and more frequent exposure to animals or soil. Although we did not find differences of HEV prevalence with respect to different water source, this does not contradict to the fact that contaminated water is the main source of GT 1 infection, especially during outbreak. In our study, we have excluded studies related to outbreak, and the number of included studies reporting water source is also very limited, which may cause bias.

Of note, there are some limitations of our study. Firstly, the number of available studies, in particular on anti-HEV IgM antibody and viral RNA positivity, is limited. We were also not able to further analyse detailed regional prevalence, GT-specific burden or clinical outcome. Secondly, we have focused on HEV prevalence, but did not estimate the incidence, which is also very relevant for assessing the disease burden. Thirdly, the assays used for HEV detection are heterogeneous in sensitivity and specificity, which may affect the estimates. Fourthly, publication bias existed in our study which was reflected in Funnel and Egger test (Figures S41-S42).

In summary, we found that 1/8 of the global population, corresponding to over 900 million individuals, have ever encountered HEV infection. Importantly, 15-110 million individuals are experiencing recent or ongoing infection. Consuming raw meat, exposing to soil, blood transfusion, travelling to endemic areas, contacting with dogs, living in rural areas and receiving lower level of education were identified as risk factors for HEV infection. Thus, our results bear important implications for assessing the global burden and devising preventive measures for controlling HEV infection.

## CONFLICT OF INTEREST

The authors do not have any disclosures to report.

## AUTHOR CONTRIBUTIONS

P. Li and J. Liu performed the study, acquisition and analysis of data. W. M. Bramer performed literature database searching. Y. Li, W. Cao, M. P. Peppelenbosch and Q. Pan discussed the data. J. Su and Z. Ma provided technical assistance. P. Li drafted the manuscript. R. A. de Man, M. P. Peppelenbosch and Q. Pan contributed to the writing of the manuscript. P. Li and J. Liu contributed equally and share co-first authorship. Q. Pan and J. Liu share co-correspondence authorship.

## ORCID

Qiuwei Pan  <https://orcid.org/0000-0001-9982-6184>

## REFERENCES

1. Khuroo MS, Kamili S. Aetiology, clinical course and outcome of sporadic acute viral hepatitis in pregnancy. *J Viral Hepat.* 2003;10(1):61-69.



2. Kamar N, Selves J, Mansuy J-M, et al. Hepatitis E virus and chronic hepatitis in organ-transplant recipients. *N Engl J Med*. 2008;358(8):811-817.
3. Zhou X, de Man RA, de Knecht RJ, Metselaar HJ, Peppelenbosch MP, Pan Q. Epidemiology and management of chronic hepatitis E infection in solid organ transplantation: a comprehensive literature review. *Rev Med Virol*. 2013;23(5):295-304.
4. Purdy MA, Harrison TJ, Jameel S, et al. ICTV virus taxonomy profile: hepeviridae. *J Gen Virol*. 2017;98(11):2645-2646.
5. Zhou J-H, Li X-R, Lan XI, et al. The genetic divergences of codon usage shed new lights on transmission of hepatitis E virus from swine to human. *Infect Genet Evol*. 2019;68:23-29.
6. Wang Y, Chen G, Pan Q, Zhao J. Chronic hepatitis E in a renal transplant recipient: the first report of genotype 4 hepatitis e virus caused chronic infection in organ recipient. *Gastroenterology*. 2018;154(4):1199-1201.
7. Meng XJ. Expanding host range and cross-species infection of hepatitis E virus. *PLoS Pathog*. 2016;12(8):e1005695.
8. Westhölter D, Hiller J, Denzer U, et al. HEV-positive blood donations represent a relevant infection risk for immunosuppressed recipients. *J Hepatol*. 2018;69(1):36-42.
9. Teshale E, Grytdal S, Howard C, et al. Evidence of person-to-person transmission of hepatitis E virus during a large outbreak in Northern Uganda. *Clin Infect Dis*. 2010;50(7):1006-1010.
10. Zeng MY, Gao H, Yan XX, et al. High hepatitis E virus antibody positive rates in dogs and humans exposed to dogs in the south-west of China. *Zoonoses Public Health*. 2017;64(8):684-688.
11. Liang H, Chen J, Xie J, et al. Hepatitis E virus serosurvey among pet dogs and cats in several developed cities in China. *PLoS ONE*. 2014;9(6):e98068.
12. Mirazo S, Ramos N, Mainardi V, Gerona S, Arbiza J. Transmission, diagnosis, and management of hepatitis E: an update. *Hepat Med*. 2014;6:45-59.
13. Horvatits T, Ozga A-K, Westhölter D, et al. Hepatitis E seroprevalence in the Americas: a systematic review and meta-analysis. *Liver Int*. 2018;38(11):1951-1964.
14. Hartl J, Otto B, Madden R, et al. Hepatitis E seroprevalence in Europe: a meta-analysis. *Viruses*. 2016;8(8):211.
15. Capai L, Falchi A, Charrel R. Meta-analysis of human IgG anti-HEV seroprevalence in industrialized countries and a review of literature. *Viruses*. 2019;11(1):84.
16. Behzadifar M, Lankarani KB, Abdi S, et al. Seroprevalence of hepatitis E virus in Iran: a systematic review and meta-analysis. *Middle East J Dig Dis*. 2016;8(3):189-200.
17. Hassan-Kadle MA, Osman MS, Ogurtsov PP. Epidemiology of viral hepatitis in Somalia: systematic review and meta-analysis study. *World J Gastroenterol*. 2018;24(34):3927-3957.
18. Zhang L, Jiao S, Yang Z, et al. Prevalence of hepatitis E virus infection among blood donors in mainland China: a meta-analysis. *Transfusion*. 2017;57(2):248-257.
19. Huang X, Huang Y, Wagner AL, Chen X, Lu Y. Hepatitis E virus infection in swine workers: a meta-analysis. *Zoonoses Public Health*. 2019;66(1):155-163.
20. Hakim MS, Wang W, Bramer WM, et al. The global burden of hepatitis E outbreaks: a systematic review. *Liver Int*. 2017;37(1):19-31.
21. Aggarwal R. Diagnosis of hepatitis E. *Nat Rev Gastroenterol Hepatol*. 2013;10(1):24-33.
22. Takahashi M, Kusakai S, Mizuo H, et al. Simultaneous detection of immunoglobulin a (IgA) and IgM antibodies against hepatitis E virus (HEV) is highly specific for diagnosis of acute HEV infection. *J Clin Microbiol*. 2005;43(1):49-56.
23. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol*. 2009;62(10):1006-1012.
24. THE Joanna Briggs Institute. The Joanna Briggs Institute Critical Appraisal tools for use in JBI systematic reviews. Checklist for Systematic Reviews and Research Syntheses. [https://joannabriggs.org/sites/default/files/2019-05/JBI\\_Critical\\_Appraisal-Checklist\\_for\\_Systematic\\_Reviews2017\\_0.pdf](https://joannabriggs.org/sites/default/files/2019-05/JBI_Critical_Appraisal-Checklist_for_Systematic_Reviews2017_0.pdf). Accessed March 03, 2020.
25. Melgaco JG, Gardinali NR, de Mello VDM, Leal M, Lewis-Ximenez LL, Pinto MA. Hepatitis E: update on prevention and control. *Biomed Res Int*. 2018;2018:5769201.
26. World Health Organization. Sixty-second world health assembly A62/22. [https://apps.who.int/gb/ebwha/pdf\\_files/WHA62-REC1/WHA62\\_REC1-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA62-REC1/WHA62_REC1-en.pdf). Accessed March 03, 2020.
27. Yazbek S, Kreidieh K, Ramia S. Hepatitis E virus in the countries of the Middle East and North Africa region: an awareness of an infectious threat to blood safety. *Infection*. 2016;44(1):11-22.
28. Kmush B, Wierzba T, Krain L, Nelson K, Labrique AB. Epidemiology of hepatitis E in low- and middle-income countries of Asia and Africa. *Semin Liver Dis*. 2013;33(1):15-29.
29. Rossi-Tamisier M, Moal V, Gerolami R, Colson P. Discrepancy between anti-hepatitis E virus immunoglobulin G prevalence assessed by two assays in kidney and liver transplant recipients. *J Clin Virol*. 2013;56(1):62-64.
30. Wenzel JJ, Preiss J, Schemmerer M, Huber B, Jilg W. Test performance characteristics of Anti-HEV IgG assays strongly influence hepatitis E seroprevalence estimates. *J Infect Dis*. 2013;207(3):497-500.
31. Sommerkorn FM, Schauer B, Schreiner T, Fickenscher H, Krumbholz A. Performance of hepatitis E virus (HEV)-antibody tests: a comparative analysis based on samples from individuals with direct contact to domestic pigs or wild boar in Germany. *Med Microbiol Immunol*. 2017;206(3):277-286.
32. Rein DB, Stevens GA, Theaker J, Wittenborn JS, Wiersma ST. The global burden of hepatitis E virus genotypes 1 and 2 in 2005. *Hepatology*. 2012;55(4):988-997.
33. WHO. *Viral hepatitis in the WHO countries of south-east Asia region*. [http://hepcasia.com/wp-content/uploads/2015/03/who\\_searo\\_viral-hepatitis-report.pdf](http://hepcasia.com/wp-content/uploads/2015/03/who_searo_viral-hepatitis-report.pdf). Accessed June 30, 2019.
34. Kamar N, Bendall R, Legrand-Abbravanel F, et al. Hepatitis E. *Lancet*. 2012;379(9835):2477-2488.
35. Tei S, Kitajima N, Ohara S, et al. Consumption of uncooked deer meat as a risk factor for hepatitis E virus infection: an age- and sex-matched case-control study. *J Med Virol*. 2004;74(1):67-70.
36. Colson P, Romanet P, Moal V, et al. Autochthonous infections with hepatitis E virus genotype 4, France. *Emerg Infect Dis*. 2012;18(8):1361-1364.
37. Ivanova A, Tefanova V, Reshetnjak I, et al. Hepatitis E virus in domestic pigs, wild boars, pig farm workers, and hunters in Estonia. *Food Environ Virol*. 2015;7(4):403-412.
38. Teixeira J, Mesquita JR, Pereira SS, et al. Prevalence of hepatitis E virus antibodies in workers occupationally exposed to swine in Portugal. *Med Microbiol Immunol*. 2017;206(1):77-81.
39. Sahli R, Fraga M, Semela D, Moradpour D, Gouttenoire J. Rabbit HEV in immunosuppressed patients with hepatitis E acquired in Switzerland. *J Hepatol*. 2019;70(5):1023-1025.
40. Sridhar S, Yip C-Y, Wu S, et al. Transmission of rat hepatitis E virus infection to humans in Hong Kong: a clinical and epidemiological analysis. *Hepatology*. 2020. <https://doi.org/10.1002/hep.31138>
41. McElroy A, Hiraide R, Bexfield N, et al. Detection of hepatitis E virus antibodies in dogs in the United Kingdom. *PLoS ONE*. 2015;10(6):e0128703.
42. Vitral CL, Pinto MA, Lewis-Ximenez LL, Khudyakov YE, dos Santos DR, Gaspar AM. Serological evidence of hepatitis E virus infection in different animal species from the Southeast of Brazil. *Mem Inst Oswaldo Cruz*. 2005;100(2):117-122.

43. Arankalle VA, Joshi MV, Kulkarni AM, et al. Prevalence of anti-hepatitis E virus antibodies in different Indian animal species. *J Viral Hepat*. 2001;8(3):223-227.
44. Dahnert L, Conraths FJ, Reimer N, Groschup MH, Eiden M. Molecular and serological surveillance of hepatitis E virus in wild and domestic carnivores in Brandenburg, Germany. *Transbound Emerg Dis*. 2018;65(5):1377-1380.
45. Song Y-J, Jeong H-J, Kim Y-J, et al. Analysis of complete genome sequences of swine hepatitis E virus and possible risk factors for transmission of HEV to humans in Korea. *J Med Virol*. 2010;82(4):583-591.
46. Okamoto H, Takahashi M, Nishizawa T, Usui R, Kobayashi E. Presence of antibodies to hepatitis E virus in Japanese pet cats. *Infection*. 2004;32(1):57-58.
47. Echevarria JM. Light and darkness: prevalence of hepatitis E virus infection among the general population. *Scientifica (Cairo)*. 2014;2014:481016.
48. Alvarado-Esquivel C, Sanchez-Anguiano LF, Hernandez-Tinoco J. Seroepidemiology of hepatitis e virus infection in mennonites in Mexico. *J Clin Med Res*. 2015;7(2):103-108.
49. Caron M, Kazanji M. Hepatitis E virus is highly prevalent among pregnant women in Gabon, central Africa, with different patterns between rural and urban areas. *Viral J*. 2008;5:158.
50. Farshadpour F, Taherkhani R, Ravanbod MR, Eghbali SS, Taherkhani S, Mahdavi E. Prevalence, risk factors and molecular evaluation of hepatitis E virus infection among pregnant women resident in the northern shores of Persian Gulf, Iran. *PLoS ONE*. 2018;13(1):e0191090.
51. Oncu S, Oncu S, Okyay P, Ertug S, Sakarya S. Prevalence and risk factors for HEV infection in pregnant women. *Med Sci Monit*. 2006;12(1):CR36-39.

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Li P, Liu J, Li Y, et al. The global epidemiology of hepatitis E virus infection: A systematic review and meta-analysis. *Liver Int*. 2020;00:1-13. <https://doi.org/10.1111/liv.14468>