



Systematic review of outcome parameters following treatment of chronic exertional compartment syndrome in the lower leg

Sanne Vogels^{1,2} | Ewan D. Ritchie^{1,2} | Thijs T. C. F. van Dongen^{1,3} |
 Marc R. M. Scheltinga⁴ | Wes O. Zimmermann^{5,6} | Rigo Hoencamp^{1,2,3,7}

¹Department of Surgery, Alrijne Hospital, Leiderdorp, The Netherlands

²Trauma Research Unit, Department of Trauma Surgery, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

³Defense Healthcare Organization, Ministry of Defense, Utrecht, The Netherlands

⁴Department of Surgery, Máxima MC, Veldhoven, The Netherlands

⁵Department of Sports Medicine, Royal Netherlands Army, Utrecht, The Netherlands

⁶Uniformed Services University of the Health Sciences, Bethesda, MD, USA

⁷Department of Surgery, Leiden University Medical Center, Leiden, The Netherlands

Correspondence

Sanne Vogels, Simon Smitweg 1, 2353 GA Leiderdorp, The Netherlands.
 Email: svogels@alrijne.nl

Objective: Surgery is the gold standard in the management of chronic exertional compartment syndrome (CECS) of the lower extremity, although recent studies also reported success following gait retraining. Outcome parameters are diverse, and reporting is not standardized. The aim of this systematic review was to analyze the current evidence regarding treatment outcome of CECS in the lower leg.

Material and Methods: A literature search and systematic analysis were performed according to the PRISMA criteria. Studies reporting on outcome following treatment of lower leg CECS were included.

Results: A total of 68 reports fulfilled study criteria ($n = 3783$; age range 12–70 year; 7:4 male-to-female ratio). Conservative interventions such as gait retraining ($n = 2$) and botulinum injection ($n = 1$) decreased ICP ($-x = 68$ mm Hg to $-x = 32$ mm Hg) and resulted in a 47% ($\pm 42\%$) rate of satisfaction and a 50% ($\pm 45\%$) rate of return to physical activity. Fasciotomy significantly decreased ICP ($-x = 76$ mm Hg to $-x = 24$ mm Hg) and was associated with an 85% ($\pm 13\%$) rate of satisfaction and an 80% ($\pm 17\%$) rate of return to activity. Return to activity was significantly more often achieved ($P < .01$) in surgically treated patients, except in one study favoring gait retraining in army personnel.

Conclusion: Surgical treatment of CECS in the lower leg results in higher rates of satisfaction and return to activity, compared to conservative treatment. However, the number of studies is limited and the level of evidence is low. Randomized controlled trials with multiple treatment arms and standardized outcome parameters are needed.

KEYWORDS

chronic exertional compartment syndrome, conservative treatment, fasciotomy, lower extremity, systematic review

1 | INTRODUCTION

Chronic exertional compartment syndrome (CECS) may affect muscle compartments mostly of the lower limb and is characterized by a sensation of tightness and pain during or

after performing repetitive physical activity. Symptoms are likely the result of a mismatch between swelling of muscular tissue within a relatively noncompliant fascia, leading to supranormal intracompartmental pressures (ICP). However, strong evidence supporting this hypothesized pathogenetic mechanism of CECS is currently lacking.¹

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Scandinavian Journal of Medicine & Science In Sports* published by John Wiley & Sons Ltd.

The diagnosis of CECS is often delayed as familiarity with the disorder among physicians is limited. Moreover, clues in patient history or physical examination in patients possibly suffering from CECS are not universally accepted. The diagnostic gold standard is invasive needle or catheter manometry that can provide values of ICP before, during, and after provocative exercise. However, the validity of these ICP measurements is seriously doubted and cutoff criteria (Box 1) are questioned.²⁻¹⁰ As a consequence of all these uncertainties, diagnostic delay in CECS can be unacceptably long.

Once CECS is diagnosed, intervention is advised as its natural course is not beneficial.⁷ Conservative therapy may entail cessation of provocative physical activity, therapeutic massage, taping, stretching, or strengthening. In addition, gait retraining and shoe modifications may be tried.^{11,12} Surgical intervention entails opening of the enveloping fascia via a fasciotomy using an open, a minimally invasive, or an endoscopic technique.¹²⁻¹⁴

Traditionally, management of CECS starts with conservative measures, followed by surgical intervention in case of treatment failure or severely disabling symptoms.¹¹ This sequence is merely based on clinical experience; a surgery first approach, or a combination of surgery and conservative measures, might also be beneficial.

Apart from clinical therapeutic considerations, presentation of treatment outcome in scientific literature is far from standardized. Commonly used outcome measures are return to physical activity, improvement of symptoms or patient-reported satisfaction, though applied methodologies often vary. At present, clinical outcome seems largely dependent on population characteristics, in particular military versus civilian athletes.^{12,15-17} In addition, outcome measures may even differ between military and civilian athletes; for example, the Single Assessment Numeric Evaluation (SANE) score¹⁸ is a validated single question instrument increasingly applied in military populations, yet rarely used with civilian patients. Conversely, these factors do influence whether a conservative or surgical approach is preferred.

A systematic review focusing on outcome following various treatments for CECS in the lower leg is currently not available. The aim of this systematic review is to analyze the current evidence regarding the most commonly reported treatment outcomes of CECS in the lower leg. Results of this review may aid in proposing a standardized report for treatment outcome regarding CECS in future research.

2 | MATERIALS AND METHODS

2.1 | Search strategy

The search strategy and systematic analysis were performed according to the PRISMA statement methodology. A search was conducted in PubMed, EMBASE, Web of Science, Cochrane,

BOX Cutoff criteria of intracompartmental pressure (ICP) for the diagnosis of chronic exertional compartment syndrome (CECS)

In studies with civilian patients, usually one or a combination of the three Pedowitz⁸³ criteria is used to define CECS of the leg:

1. pre-exercise pressure ≥ 15 mm Hg
2. one minute post-exercise pressure of ≥ 30 mm Hg
3. five minute post-exercise pressure ≥ 20 mm Hg

Yet, in service members the value most commonly referred to is the one minute after exercise measurement, with a cutoff value ≥ 35 mm Hg.¹⁰

CENTRAL, and Emcare. Key words used included “chronic exertional compartment syndrome,” “anterior compartment,” “posterior compartment,” “peroneal compartment,” “exertional leg pain,” “medial tibial pain,” “overuse injuries,” “therapy,” “surgical treatment,” and “conservative treatment.” All related MeSH terms, synonyms, and plurals were entered. Language was restricted to English and Dutch. Studies published between January 1, 1970, and May 1, 2019, were selected. In addition, relevant publications that were found outside this strategy were manually added, based on opinions of experts in the field.

2.2 | Inclusion criteria

Clinical studies with fully available text including at least five subjects diagnosed with CECS of the lower leg were considered. The diagnosis was based on a suggestive history and physical examination in the presence of elevated ICP values. Outcome following a conservative and/or surgical intervention was reported as drop in ICP values, complication rate, or recurrence rates. Moreover, studies using patient-reported outcome measures such as return to activity, satisfaction, Lower Leg Outcome Survey (LLOS),¹⁹ or the SANE,¹⁸ which numerically scores functioning of affected joints or other sections of the leg, were also included. The commonly encountered, yet heterogeneous outcome variable patient satisfaction was summarized dichotomously, using the categories “satisfied and/or improvement of symptoms” or “very satisfied and/or free of symptoms.”

2.3 | Exclusion criteria

Studies concerning acute compartment syndrome, compartment syndrome secondary to a condition other than repetitive physical activity, or a compartment syndrome in body parts other than the lower leg were excluded. Moreover, papers on combinations of CECS with medial tibial stress syndrome (MTSS) or popliteal artery entrapment syndrome were not considered, as were




FIGURE 1 Flow chart of selected studies.

TABLE 1 CECS study characteristics (n =; 68)

Author	Design	Level of Evidence	patients (n)	Patient population?	Male/Female	μ age in year (min-max)	Affected compartments	μ duration symptoms in months (min-max)	n conservative patients/n surgical patients
Akermark et al ⁴²	R	4	30	C	19/11	23 (15-36)	DP	-	-/30
Ali et al ²⁴	P	4	20	-	-	-	A	-	4/16
Allen & Barnes ⁴³	P	4	110	C	86/24	- (12-44)	A, DP	-	-/110
Balius et al ⁴⁴	P	4	7	C	-	26 (18-34)	A	-	-/7
Beck et al ⁴⁵	R	4	135	C	-	-	A, L, DP, SP	-	-/135
Biedert & Marti ²⁹	R	4	15	C	14/1	29 (-)	DP	54 (12-180)	-/15
Blackman et al ⁴⁶	P	4	7	C	6/1	25 (21-29)	A	-	7/-
Breen et al ²³	P	4	10	C	9/1	31 (-)	-	-	10/-
Cook & Bruce ⁴⁷	R	4	14	M	10/4	27 (22-38)	A, L, DP, SP	63 (6-120)	-/14
de Bruijn et al ⁴⁸	P	4	14	C	5/9	26 (18-48)	A	- (6-240)	-/14
de Fijter et al ³³	R	4	72	C + M	65/7	21 (18-37)	A	-	-/72
Detmer et al ⁴⁹	R	4	100	C	51/49	26 (-)	A, L, DP, SP	22 (-)	-/100
Diebal et al ¹⁹	P	4	10	M	8/2	20 (-)	A, L	-	10/-
Drexler et al ⁵⁰	R	4	53	C	49/4	24 (16-43)	A, L	22 (1-120)	-/53
Edmundsson et al ⁵¹	P	4	18	C	8/10	36 (16-65)	-	31 (6-180)	-/18
Finestone et al ²	R	4	36	C + M	-	24 (16-54)	A	-	-/36
Fronek et al ²⁵	R	4	18	C	8/10	24 (12-43)	A, L	-	5/13
Garcia-Mata et al ³	R	4	23	C	10/13	16 (14-18)	A, L, DP, SP	24 (7-72)	-/23
Gatenby et al ⁵²	R	4	20	C	8/12	28 (16-50)	A, L	32 (1-131)	-/20
Helmhout et al ²¹	P	3	19	M	18/1	25 (19-53)	-	-	19/-
Helmhout et al ⁵³	P	4	6	M	6/0	21 (18-27)	-	- (6-36)	6/-
Howard et al ⁵⁴	R	4	39	C	14/25	32 (-)	A, L, DP, SP	-	-/39
Irion et al ⁵⁵	R	4	13	C	6/7	20 (17-24)	A, L, DP, SP	- (0-4)	-/13
Islam & Robbs ³⁹	P	3	120	C	86/34	28 (18-53)	A, L, DP, SP	42 (12-72)	-/120
Isner-Horobeti et al ²²	R	4	16	C + M	13/3	23 (18-36)	A, L	40 (4-240)	-/16
Jarvinnen et al ⁵⁶	R	4	34	C	26/8	24 (15-41)	DP	18 (3-60)	-/34
Lohrer & Nauck ⁵⁷	R	4	17	C	8/9	24 (14-43)	A, L, DP	38 (6-360)	-/17
Maffulli et al ⁵⁸	P	4	18	C	12/6	27 (18-35)	A, L	17 (5-31)	-/18
Maher et al ⁵⁹	R	4	21	C	5/16	25 (-)	-	15 (-)	-/21
McCallum et al ⁶⁰	R	4	46	M	38/8	30 (19-50)	A, L, DP, SP	-	-/46
Micheli et al ¹⁶	R	4	47	C	-	17 (14-21)	A, L, DP, SP	15 (-)	-/47
Moeyersoons et al ⁶¹	R	4	100	C	81/19	14 (-)	-	24 (-)	-/100

Conservative Intervention(s)	Surgical approach	μ Follow-up in months (min-max)	Outcome measurements used							
			ICP	Patient satisfaction	Return to activity	SANE	LLOS	Complications (in %)	Recurrence (in %)	Reoperations (in %)
-	Open	34 (6-85)	N	Y	Y	N	N	-	-	-
PT	ES	6 (-)	N	Y	N	N	N	0	-	-
-	MI	-	Y	N	Y	N	N	0	1	-
-	MI	25 (12-38)	N	N	Y	N	N	-	-	-
-	Open, MI & ES	11 (6-28)	N	N	Y	N	N	11.2	-	19
-	Open	27 (8-72)	Y	N	N	N	N	-	-	-
M	-	1 (-)	Y	N	N	N	N	-	-	-
GR	-	12 (-)	N	Y	N	N	N	-	-	-
-	Open	37 (11-90)	N	Y	N	N	N	11.1	3.7	3.7
-	MI	21 (16-25)	N	Y	Y	N	N	3.6	-	-
-	MI	62 (-)	N	N	Y	N	N	18	2	2
-	Open & MI	5 (0-47)	N	Y	Y	N	N	7.7	3.4	3.4
GR	-	12 (-)	Y	N	Y	Y	Y	-	-	-
-	MI	50 (5-98)	N	Y	N	N	N	16.8	8.4	-
-	Open	12 (-)	N	Y	N	N	N	10.5	-	-
-	-	116 (-)	N	N	N	N	N	4.9	-	1.6
AM	Open	50 (-)	Y	Y	Y	N	N	10	-	5
-	Open	58 (12-84)	Y	Y	Y	N	N	2.3	0	2.3
-	Open	-	N	N	Y	N	N	5.6	5.6	2.8
GR	-	4 (-)	Y	N	N	Y	Y	-	-	-
GR	-	9 (-)	N	N	N	Y	Y	-	-	-
-	Open	185 (-)	N	Y	Y	N	N	13	-	6
-	Open	11 (2-60)	N	N	Y	N	N	7.7	31	7.7
-	Open	12 (-)	N	Y	N	N	N	11	0.5	0.5
BI	-	4 (3-9)	Y	Y	Y	N	N	-	-	-
-	Open	- (12-120)	N	Y	N	N	N	9	6	6
-	ES	47 (5-84)	N	Y	Y	N	N	10.5	0	5.3
-	MI	8 (5-12)	N	Y	Y	N	N	14.8	0	-
-	Open	213(32-329)	N	N	Y	N	N	-	-	-
-	-	26 (8-51)	N	Y	Y	Y	N	20	-	1.4
-	MI	50 (3-162)	N	Y	Y	N	N	-	-	-
-	Open	-	N	Y	Y	N	N	-	-	-

(Continues)

TABLE 1 (Continued)

Author	Design	Level of Evidence	patients (n)	Patient population?	Male/Female	μ age in year (min-max)	Affected compartments	μ duration symptoms in months (min-max)	n conservative patients/n surgical patients
Mouhsine et al ⁶²	R	4	18	C	10/8	25 (19-38)	A, L	-	-/18
Orlin et al ⁶³	R	4	37	C	17/20	37 (-)	A, L, DP, SP	-	-/37
Packer et al ²⁶	R	3	100	C	32/68	26 (-)	-	-	27/73
Pandya & Ganley. ⁶⁴	R	4	6	C	-	- (15-17)	A, L	-	-/6
Pasic et al ⁶⁵	R	4	46	C	23/23	30 (16-57)	A, L, DP, SP	48 (0-252)	-/46
Puranen & Alavaikko. ⁶⁶	R	4	24	C	11/13	29 (16-63)	A, DP	-	-/24
Qvarfordt et al ³⁴	R	4	15	C	8/7	29 (17-50)	A, L	36 (5-108)	-/15
Raikin et al ³⁵	R	4	16	C	6/10	25 (14-50)	A, L, DP	30 (7-72)	-/16
Reneman. ³⁶	R	4	61	C + M	58/3	21 (18-57)	A, L	-	-/61
Rettig et al ⁶⁷	R	4	12	C	1/11	21 (15-30)	A, L, DP	17 (1-36)	-/12
Roberts et al ⁶⁸	R	4	98	M	88/10	28 (-)	A	-	-/98
Rorabeck et al ⁶⁹	R	4	12	C	9/3	21 (18-26)	A, L, DP, SP	11 (5-18)	-/12
Rorabeck et al ⁷⁰	R	4	25	C	14/11	22 (-)	A, L, DP	- (12-84)	-/25
Schepsis et al ⁷¹	P	4	20	C	8/12	23 (16-37)	A, L	- (4-30)	-/20
Schepsis et al ³⁷	R	4	28	C	15/13	- (15-39)	A, L, DP	- (2-30)	-/28
Sebik & Dogan. ³⁸	P	4	6	C	4/2	28 (-)	A	-	-/6
Simpson et al ⁴	R	4	41	M	-	-	A	40 (9-110)	-/41
Singh et al ⁷²	R	4	15	C + M	13/2	31 (20-43)	A, L, DP, SP	-	-/15
Slimmon et al ⁷³	R	3	62	C	27/35	26 (-)	-	30 (2-300)	-/62
Styf & Korner. ⁷⁴	R	4	19	C	14/5	26 (17-51)	A	30 (10-84)	-/19
Sudmann. ⁷⁵	R	4	29	C + M	11/18	- (14-70)	A	- (1-120)	-/29
Takebayashi et al ⁷⁶	R	4	9	C	6/3	22 (18-24)	A, L, DP, SP	-	-/9
Thein et al ³¹	R	4	55	C	36/7	24 (-)	A	-	12/43
Turnipseed. ⁵	R	4	796	C	279/517	-	A, L, DP, SP	-	-/796
van den Brand et al ³⁰	P	3	10	C + M	8/2	23 (-)	A	-	-/10
van den Brand et al ⁷⁷	P	3	42	M	-	-	A	-	-/42
van der Wal et al ⁷	R	4	12	M	11/1	30 (-)	A	50 (-)	12/6
van Zantvoort et al ⁷⁸	R	4	30	C	14/16	29 (17-65)	A, L, DP, SP	-	-/30
van Zoest et al ²⁷	R	4	46	C	19/27	35 (-)	DP	-	19/27
Verleisdonk et al ⁸	P	4	53	C + M	47/6	- (18-41)	A	24 (-)	3/50
Verleisdonk et al ²⁸	R	4	81	C + M	77/4	24 (18-54)	A, L	6 (1-60)	-/81
Waterman et al ⁷⁹	R	4	611	M	561/50	28 (-)	A, L, DP, SP	-	-/611
Winkes et al ⁸⁰	P	4	52	C	23/29	33 (-)	A, L, DP	-	-/52

Conservative Intervention(s)	Surgical approach	μ Follow-up in months (min-max)	Outcome measurements used							
			ICP	Patient satisfaction	Return to activity	SANE	LLOS	Complications (in %)	Recurrence (in %)	Reoperations (in %)
-	Open	24 (-)	N	N	Y	N	N	0	0	0
-	Open	34 (24-52)	N	Y	N	N	N	2.7	-	-
AM	-	67 (-)	N	Y	Y	N	N	6.4	-	-
-	ES	-	N	N	Y	N	N	9.1	-	0
-	Open	55 (4-127)	N	Y	Y	N	N	-	-	11
-	-	- (2-8)	Y	N	N	N	N	-	-	-
-	Open	3 (-)	Y	Y	N	N	N	-	-	-
-	Open	16 (6-48)	N	Y	Y	N	N	-	-	-
-	Open	- (2-48)	Y	N	Y	N	N	-	-	-
-	-	- (6-24)	N	Y	Y	N	N	4.8	-	-
-	Open	23 (-)	N	Y	N	N	N	-	-	-
-	Open	12 (6-24)	N	Y	Y	N	N	-	-	-
-	Open	- (24-42)	N	Y	Y	N	N	4	12	8
-	Open	26 (12-42)	N	Y	N	N	N	3.3	-	-
-	Open	50 (-)	N	Y	N	N	N	8.7	-	2.2
-	ES	24 (-)	N	Y	Y	N	N	0	-	-
-	MI	-	N	N	Y	N	N	-	-	-
-	Open	3 (1-6)	N	Y	N	N	N	-	-	-
-	Open	51 (24-107)	N	Y	Y	N	N	3.4	11	11
-	Open	25 (19-46)	N	Y	Y	N	N	-	6.7	6.7
-	MI	- (8-30)	N	Y	N	N	N	-	-	-
-	-	-	N	Y	N	N	N	-	-	-
AM	Open	28 (-)	N	N	Y	N	N	7.4	-	-
-	Open & MI	-	N	Y	N	N	N	7	3.9	-
-	MI	-	Y	N	N	N	N	-	-	-
-	MI	-	Y	N	N	N	N	-	-	-
LM	MI	2 (-)	Y	Y	N	N	N	0	-	-
-	Open	- (12-108)	N	Y	Y	N	N	-	-	-
LM	Open	36 (19-44)	N	Y	N	N	N	-	-	-
AM	MI	-	Y	Y	N	N	N	-	5.7	1.4
-	MI	6 (-)	Y	Y	Y	N	N	2.6	-	-
-	Open	-	N	N	Y	N	N	14.3	45	5.9
-	Open	39 (3-89)	N	Y	N	N	N	-	-	-

(Continues)

TABLE 1 (Continued)

Author	Design	Level of Evidence	patients (n)	Patient population?	Male/Female	μ age in year (min-max)	Affected compartments	μ duration symptoms in months (min-max)	n conservative patients/n surgical patients
Winkes et al ⁸¹	P	4	42	C	23/19	- (17-52)	DP	- (3-72)	-/42
Wittstein et al ⁸²	R	4	9	C	4/5	24 (13-54)	A, L, DP, SP	-	-/9
Zimmermann et al ²⁰	R	3	37	M	32/5	23 (19-30)	A	11 (3-28)	37/-

Abbreviations: -, information not available; A, anterior compartment; AM, Activity modification; BI, botulinum injection; C, civil population; DP, deep posterior compartment; ES, endoscopic; GR, Gait retraining; L, lateral compartment; LM, Lifestyle modification; M, military population; MI, minimally invasive; N, no; P, prospective; PT, Physical therapy; R, retrospective; SP, superficial posterior compartment; Y, yes.

reviews, case reports, letters, expert opinions, and narrative articles. Finally, if two selected articles were reporting on the same (retrospective) cohort, the smallest study was excluded.

2.4 | Data analysis

Data extracted from included studies were study design, demographics of participants, diagnostics, type of intervention,

comparator groups, and all available outcome measures. All relevant data were independently entered into an Excel spreadsheet (Microsoft, Redmond, Washington, 2010) by two researchers (SV & ER). If absolute numbers were available, rates of recurrence, reoperation, or complication were calculated by dividing by the total number of legs. Discrepancies between reviewers were resolved by discussion.

For quantitative data, results from comparable groups of studies were pooled and means with corresponding standard

TABLE 2 Treatment outcome following conservative intervention in CECS

	n	Conservative Intervention	ICP values			Change (P-value)	SANE
			Measurement	Before intervention μ in mm Hg (\pm SD)	After intervention μ in mm Hg (\pm SD)		Before intervention (\pm SD)
Ali et al ²⁴	4	PT	-	-	-	-	-
Blackman et al ⁴⁶	7	M	3-min PE	63 (\pm 21)	68 (\pm 24)	0.156	-
Breen et al ²³	10	GR	-	-	-	-	-
Diebal et al ¹⁹	10	GR	Resting 1-min PE	40 (\pm 11) 78 (\pm 32)	36 (\pm 12) 38 (\pm 12)	- 0.002	50 (\pm 21)
Fronek et al ²⁵	5	AM	-	-	-	-	-
Helmhout et al ²¹	19	GR	1-min PE	73 (-)	47 (-)	<0.05	56 (\pm 15)
Isner-Horobeti et al ²²	16	BI	anterior 1-min PE 5-min PE lateral 1-min PE 5-min PE	65 (-) 40 (-) 60 (-) 39 (-)	22 (-) 12 (-) 19 (-) 10 (-)	<0.0001 <0.0001 <0.001 <0.01	
Packer et al ²⁶	27	AM	-	-	-	-	-
Thein et al ³¹	12	AM	-	-	-	-	-
van der Wal et al ⁷	12	LM	PE	58 (\pm 15)	51 (\pm 15)	NS	-
Van Zoest et al ²⁷	19	LM	-	-	-	-	-
Verleisdonk et al ⁸	3	AM	-	-	-	-	-
Zimmermann et al ²⁰	37	GR	-	-	-	-	51 (\pm 15)

Abbreviations: -, information not available; AM, Activity modification; BI, botulinum injection; GR, Gait retraining; ICP, intracompartmental pressure; LLOS, Lower Leg Outcome Survey (0-60, with 60 being normal); LM, Lifestyle modification; M, Massage; PE, post-exercise; PT, Physical therapy; SANE, Single Assessment Numeric Evaluation (0-100 scale, with 100 being normal).

Conservative Intervention(s)	Surgical approach	μ Follow-up in months (min-max)	Outcome measurements used							
			ICP	Patient satisfaction	Return to activity	SANE	LLOS	Complications (in %)	Recurrence (in %)	Reoperations (in %)
-	Open	26 (12-42)	N	Y	Y	N	N	6.3	6.2	1.6
-	ES	45 (5-90)	N	N	Y	N	N	14.3	0	0
GR	-	11 (3-28)	N	Y	N	Y	N	-	-	-

deviations (SD) were calculated. *P*-values < .05 were considered significant.

2.5 | Assessing the quality of evidence

The quality of studies was evaluated according to Cochrane's GRADE evidence profile. Subsequently, levels of evidence were established for all selected studies.

3 | RESULTS

A total of 7421 studies were identified (Figure 1). Following removal of duplicates and screening of title and abstract, 286 articles were reviewed for potential eligibility. Subsequently, 92 articles fitted all study criteria. After studying outcome variables, 68 studies were included (patients *n* =; 3783). The majority of the studied populations received surgical treatment (*n* =; 3612), whereas only 171 patients were treated conservatively.

After intervention (±SD)	Change (P-value)	LLOS		Change (P-value)	Satisfaction (in %)		Return to activity (in %)	
		Before intervention (±SD)	After intervention (±SD)		Satisfied or improved	Very satisfied or symptom free	Previous level	Full activity
-	-	-	-	-	0	-	-	-
-	-	-	-	-	-	-	-	-
-	-	-	-	-	17	83	-	-
90 (±10)	<0.01	67.3 (±13.7)	91.5 (±8.5)	<0.01	-	-	100	-
-	-	-	-	-	20	-	0	-
77 (±22)	0.00	72.0 (±11.3)	84.6 (±15.5)	0.00	-	-	-	-
						94		94
-	-	-	-	-	56	-	-	30
-	-	-	-	-	-	-	25	-
-	-	-	-	-	0	-	-	-
-	-	-	-	-	84	-	-	-
-	-	-	-	-	0	-	-	-
73 (±22)	<0.01	-	-	-	70	19	-	-

TABLE 3 Treatment outcome following surgical intervention for CECS

	Legs (n)	Compartments (n)	Type of Surgery	ICP in mm Hg			
				Measurement	Before intervention μ (\pm SD)	After intervention μ (\pm SD)	Change (P-value)
Akermark et al ⁴²	60	60	Open	-	-	-	-
Ali et al ²⁴	24	24	ES	-	-	-	-
Allen & Barnes ⁴³	73	84	MI	-	-	-	-
Balius et al ⁴⁴	9	9	MI	-	-	-	-
Beck et al ⁴⁵	250	741	Open, MI & ES	-	-	-	-
Biedert & Marti. ²⁹	15	-	Open	Rest PE	6 (-) 19 (-)	2 (-) 2 (-)	<0.005 <0.0001
Cook & Bruce ⁴⁷	27	56	Open	-	-	-	-
de Bruijn et al ⁴⁸	28	28	MI	-	-	-	-
de Fijter et al ³³	118	118	MI	-	-	-	-
Detmer et al ⁴⁹	-	233	Open & MI	-	-	-	-
Drexler et al ⁵⁰	95	95	MI	-	-	-	-
Edmundsson et al ⁵¹	57	121	Open	-	-	-	-
Fronek et al ²⁵	20	40	Open	-	-	-	-
Garcia-Mata et al ³	43	-	Open	-	-	-	-
Gatenby et al ⁵²	36	72	Open	-	-	-	-
Howard et al ⁵⁴	39	78	Open	-	-	-	-
Irion et al ⁵⁵	20	48	Open	-	-	-	-
Islam & Robbs. ³⁹	216	376	Open	-	-	-	-
Jarvinnen et al ⁵⁶	34	48	Open	-	-	-	-
Lohrer & Nauck ⁵⁷	38	38	ES	-	-	-	-
Maffulli et al ⁵⁸	27	38	MI	-	-	-	-
Maher et al ⁵⁹	36	-	Open	-	-	-	-
McCallum et al ⁶⁰	70	114	-	-	-	-	-
Micheli et al ¹⁶	72	103	MI	-	-	-	-
Moeyersoons & Martens ⁶¹	85	-	Open	-	-	-	-
Mouhsine et al ⁶²	29	36	Open	-	-	-	-
Orlin et al ⁶³	74	296	Open	-	-	-	-
Packer et al ²⁶	125	-	-	-	-	-	-
Pandya & Ganley. ⁶⁴	11	22	ES	-	-	-	-
Pasic et al ⁶⁵	84	244	Open	-	-	-	-
Qvarfordt et al ³⁴	30	60	Open	-	-	-	-
Raikin et al ³⁵	-	-	Open	-	-	-	-
Reneman et al ³⁶	119	-	Open	-	-	-	-
Rettig et al ⁶⁷	20	21	-	-	-	-	-
Roberts et al ⁶⁸	189	189	Open	-	-	-	-
Rorabeck et al ⁶⁹	24	56	Open	-	-	-	-
Rorabeck et al ⁷⁰	-	-	Open	-	-	-	-
Schepsis et al ⁷¹	30	45	Open	-	-	-	-

Likert Scale (in %)					Satisfaction (in %)		Return to activity (in %)	
Excellent	Good	Fair	Poor	Bad	Satisfied/ improved	Very satisfied/ symptom free	Previous level	Full activity
-	-	-	-	-	30	57	-	67
-	-	-	-	-	-	100	-	-
-	-	-	-	-	-	-	-	96
-	-	-	-	-	-	-	86	-
-	-	-	-	-	-	-	80	-
-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	78.5	-	-
31	31	23	8	8	-	-	100	-
-	-	-	-	-	-	-	-	94
-	-	-	-	-	9	73	16	75
-	-	-	-	-	75.5	-	-	-
11	61	26	2	-	-	-	-	-
-	-	-	-	-	-	94	94	-
-	-	-	-	-	-	100	-	100
-	-	-	-	-	-	-	50	40
-	-	-	-	-	79	-	78	-
-	-	-	-	-	-	-	85	-
-	-	-	-	-	6	90	-	-
41	37	15	7	-	-	-	-	-
53	6	24	18	-	-	59	82	-
-	94	-	-	-	-	-	83	11
-	-	-	-	-	-	-	75	-
-	-	-	-	-	71.4	-	37	41
47	28	15	9	-	-	-	75	-
75	6	19	-	-	-	83.5	84	-
-	-	-	-	-	-	-	100	-
-	-	-	-	-	30	63	-	-
-	-	-	-	-	81	-	-	79
-	-	-	-	-	-	-	100	-
-	-	-	-	-	30	48	63	-
-	-	-	-	-	-	93	-	-
-	-	-	-	-	20	80	87	-
-	-	-	-	-	-	-	90	-
-	-	-	-	-	83	17	25	66
-	-	-	-	-	52	-	-	-
-	-	-	-	-	-	83	-	83
-	-	-	-	-	92	-	64	-
43	47	7	3	-	90	-	-	-

(Continues)

TABLE 3 (Continued)

	Legs (n)	Compartments (n)	Type of Surgery	ICP in mm Hg			
				Measurement	Before intervention μ (\pm SD)	After intervention μ (\pm SD)	Change (P-value)
Schepesis et al ³⁷	46	64	Open	-	-	-	-
Sebik & Dogan. ³⁸	9	9	ES	-	-	-	-
Simpson et al ⁴	82	82	MI	-	-	-	-
Singh et al ⁷²	17	64	Open	-	-	-	-
Slimmon et al ⁷³	117	148	Open	-	-	-	-
Styf & Korner. ⁷⁴	30	30	Open	-	-	-	-
Sudmann. ⁷⁵	40	40	MI	-	-	-	-
Takebayashi et al ⁷⁶	12	20	-	-	-	-	-
Thein et al ³¹	54	54	Open	-	-	-	-
Turnipseed. ⁵	1396	2401	Open & MI	-	-	-	-
Van den Brand et al ³⁰	10	20	MI	PE	61 (\pm 27)	30 (\pm 8)	<0.05
Van der Wal et al ⁷	10	10	MI	PE	51 (\pm 15)	36 (\pm 5)	Sig.
van Zantvoort et al ⁷⁸	54	95	Open	-	-	-	-
Van Zoest et al ²⁷	-	-	Open	-	-	-	-
Verleisdonk et al ⁸	100	100	MI	<i>Median and range instead of mean and SD</i>			
				Rest	17 (3-23)	15 (4-29)	>0.05
				PE	62 (30-103)	22 (11-29)	<0.05
				5-min PE	37 (21-55)	16 (7-28)	<0.05
Verleisdonk et al ²⁸	151	151	MI	Rest	22 (-)	14 (-)	<0.05
				DE	58 (-)	25.4 (-)	<0.01
				5-min PE	34 (-)	25.2 (-)	<0.05
Waterman et al ⁷⁹	754	1794	Open	-	-	-	-
Winkes et al ⁸⁰	-	-	Open	-	-	-	-
Winkes et al ⁸¹	64	64	Open	-	-	-	-
Wittstein et al ⁸²	14	30	ES	-	-	-	-

Abbreviations: -, information not available; DE, during exercise; ES, endoscopic; ICP, intracompartmental pressure; MI, minimally invasive; PE, post-exercise.

An overview of study characteristics is found in Table 1. The majority (72%) was of retrospective design. An overall 7:4 male-to-female ratio was found. Study populations were dominated by adults between 20 and 30 years of age, although CECS cases up to 70 years old were identified. Additionally, more than half of the studies (56%) reported on CECS in multiple compartments, whereas 22 (32%) studies analyzed results of only one compartment. In eight (12%) articles, the affected compartments were not specified.

Inclusion of CECS patients was done by using a suggestive history of pain during exercise as a criterium in 62 articles (91%). In a total of 58 studies (85%), ICP manometry was performed, of which 24 studies applied the Pedowitz criteria. Additional imaging using radiographic images, MRI, or scintigraphy, for exclusion of stress fractures, was performed

in 23 (34%), eight (12%), and 20 (29%) articles, respectively. Ultrasonography either traditionally and/or with Doppler, for exclusion of vascular pathologies, was conducted by ten studies (15%).

3.1 | Outcome following conservative treatment

Studies reporting on ICP measurements, SANE,^{19,20} LLOS,¹⁹ patient satisfaction, or return to physical activity following conservative interventions are depicted in Table 2. Interestingly, none of the studies used similar intervention strategies (Appendix S1) or outcome measurements.

A significant drop in ICP was reported in two studies using gait retraining^{19,21} and one applying botulinum

Likert Scale (in %)					Satisfaction (in %)		Return to activity (in %)	
Excellent	Good	Fair	Poor	Bad	Satisfied/ improved	Very satisfied/ symptom free	Previous level	Full activity
49	23	14	0	-	-	-	-	-
-	-	-	-	-	-	100	100	-
-	-	-	-	-	-	-	29	46
-	-	-	-	-	-	100	-	-
31	18	19	13	-	-	-	42	-
-	-	-	-	-	74	-	63	32
-	-	-	-	-	15	70	-	-
33	33	33	-	-	-	-	-	-
-	-	-	-	-	-	-	77.4	-
-	-	-	-	-	-	91	-	-
-	-	-	-	-	100	-	-	-
13	20	23	10	-	-	-	30	-
-	-	19	-	-	52	-	-	-
-	-	-	-	-	83	-	-	-
-	-	-	-	-	76	-	-	76
-	-	-	-	-	-	-	-	72
17	31	-	-	-	-	-	-	-
23	31	9	8	-	76	-	29	-
-	-	-	-	-	-	-	89	-

injections.²² Moreover, lower ICP values were associated with an improved outcome as reflected by SANE and LLOS scores. Improvement of symptoms or satisfaction was reported by 47% ($\pm 42\%$) of the patients who completed a follow-up analysis, whereas 50% ($\pm 45\%$) returned to a form of physical activity. The well-structured gait retraining programs^{20,23} and treatment with botulinum injections²² scored highest with satisfaction rates ranging from 89% to 100%, whereas all studies with patients alone initiated modifications in activity and/or lifestyle^{7,8,24-27} scored between 0% and 84% satisfaction.

Among the 171 conservatively treated patients, six cases were reported to eventually opt for surgery.⁷ Additionally, a significant reduction of individuals requiring subsequent surgery was found in military populations (not mentioned in Table 2).^{19,20}

3.2 | Outcome following surgical treatment

Clinical outcome with respect to lowered ICP values, patient satisfaction, return to activity, rates of complication, recurrence, and reoperation after surgical intervention is depicted in Table 3.

ICP values were obtained both before and after surgical intervention in nine of thirty studies. Five^{7,8,28-30} of these found a statistically significant reduction of ICP, suggesting that surgical intervention is effective in reducing muscle compartment pressures.

Patient-reported outcome measures and rates of return to activity reveal that the majority of CECS patients were satisfied and returned to previous levels of activity. In addition, 58% ($\pm 29.6\%$) were satisfied with the treatment results and experienced reduction of symptoms, whereas 78% ($\pm 21.2\%$)

TABLE 4 Comparing conservative and surgical interventions in CECS

Study				ICP in mm Hg				
	Conservative	Surgical		Measurement	Conservative		Surgical	
		Patients (n)	Legs (n)		Compartments (n)	Before intervention μ (\pm SD)	After intervention μ (\pm SD)	Before intervention μ (\pm SD)
Ali et al ²⁴	4	24	24	-	-	-	-	-
Fronck et al ²⁵	5	20	40	Resting	17.14(\pm 9.05)	-	18.5 (\pm 7.94)	9.2 (\pm 0.98)
				PE	57.0 (\pm 22.4)	-	55.7(\pm 33.25)	12.7(\pm 2.49)
				1-min PE	42.3 (\pm 21.0)	-	37.4 (\pm 14.3)	10 (\pm 0)
				5-min PE	34.3 (\pm 22.3)	-	27.8 (\pm 9.61)	9.8 (\pm 2.23)
Packer et al ²⁶	27	125	-	Rest	6.30 (\pm 2.92)	-	6.60 (\pm 3.23)	-
				PE	26.67(\pm 11.26)	-	40.44(\pm 9.60)	-
Thein et al ³¹	12	54	54	-	-	-	-	-
Van der Wal et al ⁷	12	10	10	PE	58 (\pm 15)	51 (\pm 15)	51 (\pm 15)	36 (\pm 5)
Van Zoest et al ²⁷	19	-	-	Rest	14.5 (-)	-	22 (-)	-
				PE	15.5 (-)	-	34.5 (-)	-
				1-min PE	13.5 (-)	-	31.5 (-)	-
				5-min PE	12 (-)	-	29.5 (-)	-
Verleisdonk et al ⁸	3	100	100	<i>Median and range instead of mean and SD</i>				
				Rest	-	-	17 (3-23)	15 (4-29)
				PE	-	-	62 (30-103)	22 (11-29)
				5-min PE	-	-	37 (21-55)	16 (7-28)

Abbreviations: -, information not available; ICP, intracompartmental pressure; NS, non-significant; PE, post-exercise.

were very satisfied and/or free of symptoms. Combining these results allows for calculating an 85% (\pm 13%) overall satisfaction rate. Moreover, the average proportion that returned to some form of physical activity after surgery was 80% (\pm 17.3). However, return to previous level and/or full activity was on average 69% (\pm 25.5%) and 65% (\pm 25.0%), respectively.

Surgical complications and rates of recurrence and reoperations (Table 1) indicate that approximately 8% (\pm 5.3%) of the studied CECS patients experienced surgical complications, mainly wound problems or nerve damage. Irrespective of surgical technique or operated compartment, recurrence, and reoperation rates were 7% (\pm 10.8%) and 5% (\pm 4.3%), respectively. Comparing studies that focused on civilian (n =; 32) or military patients (n =; 3) revealed a significantly higher complication rate among patients that serve in the armed forces (civilian 7.1% \pm 4.6% versus military 15.1% \pm 4.5%, $P =$; .01). Similar results are found with respect to recurrence (civilian 5.6% \pm 7.7% versus military 24.4% \pm 29.2%, $P =$; .03) and reoperations rates (civilian 5.2% \pm 4.8% versus military 21.4% \pm 32.6%, $P =$; .03).

A list of different postoperative treatment protocols after surgical intervention is found in Appendix S2. Days of rest, weight bearing, use of compressive bandages, and sport limitations varied widely among studies.

3.3 | Comparison of conservative and surgical interventions

Table 4 lists studies comparing conservative and surgical interventions. Packer et al²⁶ and Thein et al³¹ compared rates of return to activity and found significant differences favoring surgical intervention. However, Packer et al²⁶ found similar satisfaction rates. Interestingly, Zimmermann et al²⁰ reported in a military population a higher percentage that returned to active duty following conservative treatment compared to surgical intervention.

4 | DISCUSSION & CONCLUSION

This systematic review is the first to analyze studies reporting on outcome following conservative and surgical treatment in patients with CECS in any compartment of the lower leg, not just the posterior compartment.³² No randomized controlled trials were found.

Most CECS studies report on beneficial effects of surgical therapy, with an overall 85% satisfaction rate and an 80% rate of return to physical activity. In contrast, conservative interventions were seemingly associated with lower rates of satisfaction and return to activity (47% and 50%, respectively). Only two studies compared both modalities in one

Overall Difference (p-value)	Satisfaction (in %)					Return to activity (in%)					
	Conservative		Surgical			Conservative		Surgical			Overall Difference (p-value)
	Satisfied/improved	Very satisfied/symptom free	Satisfied/improved	Very satisfied/symptom free	Difference (p-value)	Previous level	Full activity	Previous level	Full activity		
-	0	-	-	100	-	-	-	-	-	-	
-	20	-	-	94	-	0	-	94	-	-	
NS <0.001	56	-	81	-	0.011	-	30	-	79	<0.001	
-	-	-	-	-	-	25	-	77.4	-	0.001	
-	0	-	100	-	-	-	-	-	-	-	
NS <0.05 <0.05 <0.05	84	-	52	-	-	-	-	-	-	-	
-	0	-	83	-	-	-	-	-	-	-	
-											
-											
-											

model, reporting statistically superior results following a fasciotomy.^{26,31} However, caution regarding an interpretation is required due to the limited number of studies on conservative treatment with substantial smaller study populations.

This review demonstrates that ICP measurements are infrequently used as a treatment outcome parameter, even though they are considered the gold standard in diagnosing CECS. Only sixteen of the included studies measured ICPs before and after intervention, with only nine studies reporting on corresponding P-values. The use of ICP measurements as outcome measure cannot be confirmed, nor discarded with current literature.

Another interesting finding is that this overview consistently found a potential difference between surgically treated civilian and military study populations with significant higher rates of postoperative complications, recurrence, and reoperations in the military, as was already suggested by previous literature.^{12,15,17} Even though these observations were made on the basis of different population sizes (civilian n =; 1975, military n =; 671), these findings may suggest conservative treatment in military patients may be preferred compared to surgery.

This review was subject to a number of limitations, the most prominent being the lack of uniformity among outcome measures. Moreover, follow-up data were often obtained in substantially smaller number of patients than

initially treated, potentially introducing selection bias. This principle also applies to the exact determination of overall recurrence rates and complications, especially when information on unilaterality or bilaterality of symptoms was missing.

This review was further hampered by the heterogeneity among study populations. Variation was found in studies with respect to the inclusion of patients with fascial herniae,^{2,3,5,6,8,28,33-38} presence of concomitant MTSS²⁰ or affected upper extremities.³⁹ An attempt to overcome this heterogeneity was made by solely including studies that allowed for extraction of data only concerning CECS in the lower extremity. Nevertheless, any conclusion based on the present review must be taken with caution.

Defining uniform and generally applicable outcome parameters will likely simplify future data comparison. This process is facilitated by initiating a consensus via the Delphi method as was already conducted for various other entities by the International Consortium for Health Outcomes Measurement.⁴⁰ Based on the content of Hip & Knee osteoarthritis set,⁴¹ we wish to propose a potential outline from which standardization can be initiated (Figure 2). The use of a 5-point Likert scale is preferred for all questions related to symptoms or performance. Currently, such a set of standardized outcome measurements will be applied by our study group to military civilian collaboration, with special




FIGURE 2 Proposed outline for a standardized Patient Reported Outcome Measurement in patients with CECS

emphasis on prevention, conservative treatment, and non-invasive diagnostics.

In conclusion, the present review found that surgical treatment for CECS resulted in a minimal 80% overall satisfaction and return to physical activity rate. In contrast, conservative interventions were associated with lower rates of satisfaction and return to activity up to 50%. As these findings are based on low-quality studies demonstrating a large heterogeneity, higher quality studies including randomized controlled trials with univocal endpoints are required for determining any superior treatment regimen in the lower leg CECS.

5 | PERSPECTIVE

Surgery is currently the gold standard in the management of CECS of the lower extremity, although recent studies also reported success following gait retraining. This review

provides an extensive overview of all published evidence regarding treatment outcome for both conservative and surgical therapy. This study therefore serves educational purposes for healthcare professionals working with CECS patients, who can be found among all areas of sport in both civil and military populations. The presented overview aids evidence-based and shared decision making in the discussion between healthcare provider and patients; it offers clear implications and guidelines for future treatment and research.

CONFLICT OF INTEREST

There is no conflict of interest to declare.


AUTHOR CONTRIBUTION

SV and ER contributed equally to this manuscript. SV, ER, and RH conceptualized the study. SV wrote the study protocol. SV and ER conducted the literature searches, the study selection, the data extraction, and the study quality

assessment. SV and ER performed all statistical analyses. SV and ER drafted all sections of the manuscript. All authors critically revised the draft manuscript and contributed to the subsequent revisions and the final version of the manuscript.

ORCID

Sanne Vogels  <https://orcid.org/0000-0002-3686-325X>

Wes O. Zimmermann  <https://orcid.org/0000-0002-5814-9794>

REFERENCES

- Wuellner JC, Nathe CD, Kreulen CD, Burnham KJ, Giza E. Chronic exertional compartment syndrome: the Athlete's claudication. *Oper Tech Sports Med.* 2017;25(2):52-58.
- Finestone AS, Noff M, Nassar Y, Moshe S, Agar G, Tamir E. Management of chronic exertional compartment syndrome and fascial hernias in the anterior lower leg with the forefoot rise test and limited fasciotomy. *Foot Ankle Int.* 2014;35(3):285-292.
- Garcia-Mata S, Hidalgo-Ovejero A, Martinez-Grande M. Chronic exertional compartment syndrome of the legs in adolescents. *J Pediatr Orthop.* 2001;21(3):328-334.
- Simpson C, Roscoe D, Hughes S, Hulse D, Guthrie H. Surgical outcomes for chronic exertional compartment syndrome following improved diagnostic criteria. *J R Army Med Corps.* 2019. <https://militaryhealth.bmj.com/content/early/2019/04/15/jramc-2019-001171>.
- Turnipseed WD. Clinical review of patients treated for atypical claudication: a 28-year experience. *J Vasc Surg.* 2004;40(1):79-85.
- Turnipseed WD. Diagnosis and management of chronic compartment syndrome. *Surgery.* 2002;132(4):613-617.
- Van der Wal WA, Heesterbeek PJ, Van den Brand JG, Verleisdonk EJ. The natural course of chronic exertional compartment syndrome of the lower leg. *Knee Surg Sports Traumatol Arthrosc.* 2015;23(7):2136-2141.
- Verleisdonk EJ, Schmitz RF, van der Werken C. Long-term results of fasciotomy of the anterior compartment in patients with exercise-induced pain in the lower leg. *Int J Sports Med.* 2004;25(3):224-229.
- Roberts A, Franklyn-Miller A. The validity of the diagnostic criteria used in chronic exertional compartment syndrome: a systematic review. *Scand J Med Sci Sports.* 2012;22(5):585-595.
- Zimmermann WO, Ligthert E, Helmhout PH, et al. Intracompartmental pressure measurements in 501 service members with exercise-related leg pain. *Transl J Am Coll Sports Med.* 2018;3(14):107-112.
- Rajasekaran S, Hall MM. Nonoperative management of chronic exertional compartment syndrome: a systematic review. *Curr Sports Med Rep.* 2016;15(3):191-198.
- Zimmermann WO, Helmhout PH, Beutler A. Prevention and treatment of exercise related leg pain in young soldiers; a review of the literature and current practice in the Dutch Armed Forces. *J R Army Med Corps.* 2017;163(2):94-103.
- Lohrer H, Malliaropoulos N, Korakakis V, Padhiar N. Exercise-induced leg pain in athletes: diagnostic, assessment, and management strategies. *Phys Sportsmed.* 2019;47(1):47-59.
- Vajapey S, Miller TL. Evaluation, diagnosis, and treatment of chronic exertional compartment syndrome: a review of current literature. *Phys Sportsmed.* 2017;45(4):391-398.
- Dunn JC, Waterman BR. Chronic exertional compartment syndrome of the leg in the military. *Clin Sports Med.* 2014;33(4):693-705.
- Micheli LJ, Solomon R, Solomon J, Plasschaert VF, Mitchell R. Surgical treatment for chronic lower-leg compartment syndrome in young female athletes. *Am J Sports Med.* 1999;27(2):197-201.
- Meulekamp MZ, van der Wurff P, van der Meer A, Lucas C. Identifying prognostic factors for conservative treatment outcomes in servicemen with chronic exertional compartment syndrome treated at a rehabilitation center. *Mil Med Res.* 2017;4(1):36.
- Williams GN, Gangel TJ, Arciero RA, Uhorchak JM, Taylor DC. Comparison of the single assessment numeric evaluation method and two shoulder rating scales. *Am J Sports Med.* 1999;27(2):214-221.
- Diebal AR, Gregory R, Alitz C, Gerber JP. Forefoot running improves pain and disability associated with chronic exertional compartment syndrome. *Am J Sports Med.* 2012;40(5):1060-1067.
- Zimmermann WO, Hutchinson MR, Van den Berg R, Hoencamp R, Backx FJG, Bakker EWP. Conservative treatment of anterior chronic exertional compartment syndrome in the military, with a mid-term follow-up. *BMJ Open Sport Exerc Med.* 2019;5(1):e000532.
- Helmhout PH, Diebal AR, van der Kaaden L, Harts CC, Beutler A, Zimmermann WO. The effectiveness of a 6-week intervention program aimed at modifying running style in patients with chronic exertional compartment syndrome: results from a series of case studies. *Orthop J Sports Med.* 2015;3(3):2325967115575691.
- Isner-Horobeti ME, Dufour SP, Blaes C, Lecocq J. Intramuscular pressure before and after botulinum toxin in chronic exertional compartment syndrome of the leg: a preliminary study. *Am J Sports Med.* 2013;41(11):2558-2566.
- Breen DT, Foster J, Falvey E, Franklyn-Miller A. Gait re-training to alleviate the symptoms of anterior exertional lower leg pain: a case series. *Int J Sports Phys Ther.* 2015;10(1):85-94.
- Ali T, Mohammed F, Mencia M, Maharaj D, Hoford R. Surgical management of exertional anterior compartment syndrome of the leg. *West Indian Med J.* 2013;62(6):529-532.
- Fronek J, Mubarak SJ, Hargens AR, et al. Management of chronic exertional anterior compartment syndrome of the lower extremity. *Clin Orthop Relat Res.* 1987;220:217-227.
- Packer JD, Day MS, Nguyen JT, Hobart SJ, Hannafin JA, Metz J. Functional outcomes and patient satisfaction after fasciotomy for chronic exertional compartment syndrome. *Am J Sports Med.* 2013;41(2):430-436.
- van Zoest WJ, Hoogveen AR, Scheltinga MR, Sala HA, van Mourik JB, Brink PR. Chronic deep posterior compartment syndrome of the leg in athletes: postoperative results of fasciotomy. *Int J Sports Med.* 2008;29(5):419-423.
- Verleisdonk EJ, van den Helder CJ, Hoogendoorn HA, van der Werken C. Good results of fasciotomy in chronic compartment syndrome of the lower leg. *Ned Tijdschr Geneesk.* 1996;140(50):2513-2517.
- Biedert RM, Marti B. Intracompartmental pressure before and after fasciotomy in runners with chronic deep posterior compartment syndrome. *Int J Sports Med.* 1997;18(5):381-386.
- Van Den Brand JGH, Verleisdonk EJMM, van der Werken C. Near infrared spectroscopy in the diagnosis of chronic exertional compartment syndrome. *Am J Sports Med.* 2004;32(2):452-456.
- Thein R, Tilbor I, Rom E, et al. Return to sports after chronic anterior exertional compartment syndrome of the leg:

- conservative treatment versus surgery. *J Orthop Surg (Hong Kong)*. 2019;27(2):2309499019835651.
32. Winkes MB, Hoogeveen AR, Scheltinga MR. Is surgery effective for deep posterior compartment syndrome of the leg? A systematic review. *Br J Sports Med*. 2014;48(22):1592-1598.
 33. de Fijter WM, Scheltinga MR, Luiting MG. Minimally invasive fasciotomy in chronic exertional compartment syndrome and fascial hernias of the anterior lower leg: short- and long-term results. *Mil Med*. 2006;171(5):399-403.
 34. Qvarfordt P, Christenson JT, Eklof B, Ohlin P, Saltin B. Intramuscular pressure, muscle blood flow, and skeletal muscle metabolism in chronic anterior tibial compartment syndrome. *Clin Orthop Relat Res*. 1983;179:284-290.
 35. Raikin SM, Rapuri VR, Vitanzo P. Bilateral simultaneous fasciotomy for chronic exertional compartment syndrome. *Foot Ankle Int*. 2005;26(12):1007-1011.
 36. Reneman RS. The anterior and the lateral compartmental syndrome of the leg due to intensive use of muscles. *Clin Orthop Relat Res*. 1975;113:69-80.
 37. Schepesis AA, Martini D, Corbett M. Surgical management of exertional compartment syndrome of the lower leg. Long-term follow-up. *Am J Sports Med*. 1993;21(6):811-817.
 38. Sebik A, Dogan A. A technique for arthroscopic fasciotomy for the chronic exertional tibialis anterior compartment syndrome. *Knee Surg Sports Traumatol Arthrosc*. 2008;16(5):531-534.
 39. Islam J, Robbs JV. Management of chronic exertional compartment syndrome. *S Afr J Surg*. 2015;53(2):59-61.
 40. Porter ME, Larsson S, Lee TH. Standardizing patient outcomes measurement. *N Engl J Med*. 2016;374(6):504-506.
 41. Ackerman IN, Cavka B, Lippa J, Bucknill A. The feasibility of implementing the ICHOM Standard Set for Hip and Knee Osteoarthritis: a mixed-methods evaluation in public and private hospital settings. *J Patient Rep Outcomes*. 2018;2:32.
 42. Akermark C, Ljungdahl M, Johansson C. Long-term results of fasciotomy caused by medial tibial syndromes in athletes. *Scand J Med Sci Sports*. 1991;1(1):59-61.
 43. Allen MJ, Barnes MR. Exercise pain in the lower leg. Chronic compartment syndrome and medial tibial syndrome. *J Bone Joint Surg Br*. 1986;68(5):818-823.
 44. Balius R, Bong DA, Ardevol J, Pedret C, Codina D, Dalmau A. Ultrasound-guided fasciotomy for anterior chronic exertional compartment syndrome of the leg. *J Ultrasound Med*. 2016;35(4):823-829.
 45. Beck JJ, Tepolt FA, Miller PE, Micheli LJ, Kocher MS. Surgical treatment of chronic exertional compartment syndrome in pediatric patients. *Am J Sports Med*. 2016;44(10):2644-2650.
 46. Blackman PG, Simmons LR, Crossley KM. Treatment of chronic exertional anterior compartment syndrome with massage: a pilot study. *Clin J Sport Med*. 1998;8(1):14-17.
 47. Cook S, Bruce G. Fasciotomy for chronic compartment syndrome in the lower limb. *ANZ J Surg*. 2002;72(10):720-723.
 48. de Bruijn JA, van Zantvoort APM, Winkes MB, et al. Feasibility and safety of an operative tool for anterior chronic exertional compartment syndrome treatment. *Foot Ankle Int*. 2015;36(12):1475-1482.
 49. Detmer DE, Sharpe K, Sufit RL, Girdley FM. Chronic compartment syndrome: diagnosis, management, and outcomes. *Am J Sports Med*. 1985;13(3):162-170.
 50. Drexler M, Rutenberg TF, Rozen N, et al. Single minimal incision fasciotomy for the treatment of chronic exertional compartment syndrome: outcomes and complications. *Arch Orthop Trauma Surg*. 2017;137(1):73-79.
 51. Edmundsson D, Toolanen G, Sojka P. Chronic compartment syndrome also affects nonathletic subjects: a prospective study of 63 cases with exercise-induced lower leg pain. *Acta Orthop*. 2007;78(1):136-142.
 52. Gatenby G, Haysom S, Twaddle B, Walsh S. Functional outcomes after the surgical management of isolated anterolateral leg chronic exertional compartment syndrome. *Orthop J Sports Med*. 2017;5(11):2325967117737020.
 53. Helmhout PH, Diebal-Lee MA, Poelsma LR, Harts CC, Zimmermann LW. Modifying marching technique in military service members with chronic exertional compartment syndrome: a case series. *Int J Sports Phys Ther*. 2016;11(7):1106-1124.
 54. Howard JL, Mohtadi NG, Wiley JP. Evaluation of outcomes in patients following surgical treatment of chronic exertional compartment syndrome in the leg. *Clin J Sport Med*. 2000;10(3):176-184.
 55. Irion V, Magnussen RA, Miller TL, Kaeding CC. Return to activity following fasciotomy for chronic exertional compartment syndrome. *Eur J Orthop Surg Traumatol*. 2014;24(7):1223-1228.
 56. Jarvinnen M, Aho H, Niittymaki S. Results of the surgical treatment of the medial tibial syndrome in athletes. *Int J Sports Med*. 1989;10(1):55-57.
 57. Lohrer H, Nauck T. Endoscopically assisted release for exertional compartment syndromes of the lower leg. *Arch Orthop Trauma Surg*. 2007;127(9):827-834.
 58. Maffulli N, Loppini M, Spiezia F, D'Addona A, Maffulli GD. Single minimal incision fasciotomy for chronic exertional compartment syndrome of the lower leg. *J Orthop Surg Res*. 2016;11(1):61.
 59. Maher JM, Brook EM, Chiodo C, Smith J, Bluman EM, Matzkin EG. Patient-reported outcomes following fasciotomy for chronic exertional compartment syndrome. *Foot Ankle Spec*. 2018;11(5):471-477.
 60. McCallum JR, Cook JB, Hines AC, Shaha JS, Jex JW, Orchowski JR. Return to duty after elective fasciotomy for chronic exertional compartment syndrome. *Foot Ankle Int*. 2014;35(9):871-875.
 61. Moeyersoons JP, Martens M. Chronic compartment syndrome: diagnosis and management. *Acta Orthop*. 1992;58(1):23-27.
 62. Mouhsine E, Garofalo R, Moretti B, Gremion G, Akiki A. Two minimal incision fasciotomy for chronic exertional compartment syndrome of the lower leg. *Knee Surg Sports Traumatol Arthrosc*. 2006;14(2):193-197.
 63. Orlin JR, Oen J, Andersen JR. Changes in leg pain after bilateral fasciotomy to treat chronic compartment syndrome: a case series study. *J Orthop Surg Res*. 2013;8:6.
 64. Pandya NK, Ganley TJ. Single-incision endoscopically-assisted compartment releases for exertional compartment syndrome in pediatric patients. *Curr Orthop Pract*. 2010;21(2):213-217.
 65. Pasic N, Bryant D, Willits K, Whitehead D. Assessing outcomes in individuals undergoing fasciotomy for chronic exertional compartment syndrome of the leg. *Arthroscopy*. 2015;31(4):707-713.
 66. Puranen J, Alavaikko A. Intracompartmental pressure increase on exertion in patients with chronic compartment syndrome in the leg. *J Bone Joint Surg Am*. 1981;63(8):1304-1309.
 67. Rettig AC, McCarroll JR, Hahn RG. Chronic compartment syndrome: surgical intervention in 12 cases. *Phys Sports Med*. 1991;19(4):63-70.

68. Roberts AJ, Krishnasamy P, Quayle JM, Houghton JM. Outcomes of surgery for chronic exertional compartment syndrome in a military population. *J R Army Med Corps*. 2015;161(1):42-45.
69. Rorabeck CH, Bourne RB, Fowler PJ. The surgical treatment of exertional compartment syndrome in athletes. *J Bone Joint Surg Am*. 1983;65(9):1245-1251.
70. Rorabeck CH, Fowler PJ, Nott L. The results of fasciotomy in the management of chronic exertional compartment syndrome. *Am J Sports Med*. 1988;16(3):224-227.
71. Schepsis AA, Gill SS, Foster TA. Fasciotomy for exertional anterior compartment syndrome: is lateral compartment release necessary? *Am J Sports Med*. 1999;27(4):430-435.
72. Singh N, Sidawy AN, Bottoni CR, et al. Physiological changes in venous hemodynamics associated with elective fasciotomy. *Ann Vasc Surg*. 2006;20(3):301-305.
73. Slimmon D, Bennell K, Brukner P, Crossley K, Bell SN. Long-term outcome of fasciotomy with partial fasciectomy for chronic exertional compartment syndrome of the lower leg. *Am J Sports Med*. 2002;30(4):581-588.
74. Styf JR, Korner LM. Chronic anterior-compartment syndrome of the leg. Results of treatment by fasciotomy. *J Bone Joint Surg Am*. 1986;68(9):1338-1347.
75. Sudmann E. The painful chronic anterior lower leg syndrome. A prospective clinical and experimental study. *Acta Orthop Scand*. 1979;50(5):573-581.
76. Takebayashi S, Takazawa H, Sasaki R, Miki H, Soh R, Nishimura J. Chronic exertional compartment syndrome in lower legs: localization and follow-up with thallium-201 SPECT imaging. *J Nucl Med*. 1997;38(6):972-976.
77. Van den Brand JG, Nelson T, Verleisdonk EJ, van der Werken C. The diagnostic value of intracompartmental pressure measurement, magnetic resonance imaging, and near-infrared spectroscopy in chronic exertional compartment syndrome: a prospective study in 50 patients. *Am J Sports Med*. 2005;33(5):699-704.
78. van Zantvoort APM, de Bruijn JA, Hundscheid HPH, Crujjsen-Raaijmakers M, Teijink JAW, Scheltinga MR. Fasciotomy for lateral lower-leg chronic exertional compartment syndrome. *Int J Sports Med*. 2018;39(14):1081-1087.
79. Waterman BR, Laughlin M, Kilcoyne K, Cameron KL, Owens BD. Surgical treatment of chronic exertional compartment syndrome of the leg: failure rates and postoperative disability in an active patient population. *J Bone Joint Surg Am*. 2013;95(7):592-596.
80. Winkes MB, Hoogveen AR, Houterman S, Giesberts A, Wijn PF, Scheltinga MR. Compartment pressure curves predict surgical outcome in chronic deep posterior compartment syndrome. *Am J Sports Med*. 2012;40(8):1899-1905.
81. Winkes MB, van Zantvoort APM, de Bruijn JA, et al. Fasciotomy for deep posterior compartment syndrome in the lower leg: a prospective study. *Am J Sports Med*. 2016;44(5):1309-1316.
82. Wittstein J, Moorman CT III, Levin LS. Endoscopic compartment release for chronic exertional compartment syndrome: surgical technique and results. *Am J Sports Med*. 2010;38(8):1661-1666.
83. Pedowitz RA, Hargens AR, Mubarak SJ, Gershuni DH. Modified criteria for the objective diagnosis of chronic compartment syndrome of the leg. *Am J Sports Med*. 1990;18(1):35-40.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Vogels S, Ritchie ED, van Dongen TTCF, Scheltinga MRM, Zimmermann WO, Hoencamp R. Systematic review of outcome parameters following treatment of chronic exertional compartment syndrome in the lower leg. *Scand J Med Sci Sports*. 2020;00:1–19. <https://doi.org/10.1111/sms.13747>