

EUR Research Information Portal

No evidence of aberrant amyloid β and phosphorylated tau expression in herpes simplex virus-infected neurons of the trigeminal ganglia and brain

Published in:
Brain Pathology

Publication status and date:
Published: 01/07/2022

DOI (link to publisher):
[10.1111/bpa.13044](https://doi.org/10.1111/bpa.13044)

Document Version
Publisher's PDF, also known as Version of record

Document License/Available under:
CC BY-NC-ND

Citation for the published version (APA):
Tran, D. N., Bakx, A. T. C. M., van Dis, V., Aronica, E., Verdijk, R. M., & Ouwendijk, W. J. D. (2022). No evidence of aberrant amyloid β and phosphorylated tau expression in herpes simplex virus-infected neurons of the trigeminal ganglia and brain. *Brain Pathology*, 32(4), Article e13044. <https://doi.org/10.1111/bpa.13044>
[Link to publication on the EUR Research Information Portal](#)

Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.


In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: openaccess.library@eur.nl. Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

RESEARCH ARTICLE

No evidence of aberrant amyloid β and phosphorylated tau expression in herpes simplex virus-infected neurons of the trigeminal ganglia and brain

Diana N. Tran¹ | Amy T. C. M. Bakx¹ | Vera van Dis² | Eleonora Aronica³ | Robert M. Verdijk² | Werner J. D. Ouwendijk¹ 

¹Department of Viroscience, Erasmus MC, University Medical Center Rotterdam, the Netherlands

²Department of Pathology, Erasmus MC, University Medical Center Rotterdam, the Netherlands

³Department of (Neuro)Pathology, Amsterdam Neuroscience, Amsterdam University Medical Centers, University of Amsterdam, Amsterdam, the Netherlands

Correspondence

Werner J. D. Ouwendijk, Department of Viroscience, Erasmus MC, University Medical Center Rotterdam, PO Box 2040, Rotterdam 3000 CA, the Netherlands. Email: w.ouwendijk@erasmusmc.nl

Funding information

Research reported in this publication was in part supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award Number R01AI151290 (W.J.D.O.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. This study was funded in part by a Human Disease Model Award 2020 (Erasmus MC)

Abstract

Increasing evidence supports the role of neurotropic herpes simplex virus 1 (HSV-1) in the pathogenesis of Alzheimer's disease (AD). However, it is unclear whether previously reported findings in HSV-1 cell culture and animal models can be translated to humans. Here, we analyzed clinical specimens from latently HSV-1 infected individuals and individuals with lytic HSV infection of the brain (herpes simplex encephalitis; HSE). Latent HSV-1 DNA load and latency-associated transcript (LAT) expression were identical between trigeminal ganglia (TG) of AD patients and controls. Amyloid β (A β) and hyperphosphorylated tau (pTau) were not detected in latently HSV-infected TG neurons. Aging-related intraneuronal A β accumulations, neurofibrillary tangles (NFT), and/or extracellular A β plaques were observed in the brain of some HSE patients, but these were neither restricted to HSV-infected neurons nor brain regions containing virus-infected cells. Analysis of unique brain material from an AD patient with concurrent HSE showed that HSV-infected cells frequently localized close to A β plaques and NFT, but were not associated with exacerbated AD-related pathology. HSE-associated neuroinflammation was not associated with specific A β or pTau phenotypes. Collectively, we observed that neither latent nor lytic HSV infection of human neurons is directly associated with aberrant A β or pTau protein expression in ganglia and brain.

KEYWORDS

Alzheimer's disease, amyloid β , herpes simplex encephalitis, herpes simplex virus, neurofibrillary tangles, varicella-zoster virus

1 | BACKGROUND

Alzheimer's disease (AD) is a progressive neurodegenerative disease that accounts for about 80% of dementia cases [1]. Pathological hallmarks of AD include the formation of

extracellular amyloid β (A β) protein plaques and intracellular neurofibrillary tangles (NFT) composed of hyperphosphorylated tau (pTau) protein [2]. The deposition of A β plaques is an early event in AD pathogenesis, which induces subsequent inflammation, accumulation of NFTs, and

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Brain Pathology* published by John Wiley & Sons Ltd on behalf of International Society of NeuroPathology.

ultimately neuronal dysfunction and cell death [3]. Genetic risk factors for AD development are principally involved in A β processing and clearance – most notably apolipoprotein ϵ 4 allele (*APOE4*) – and are expressed by CNS-resident immune cells [4]. Additionally, an estimated 21%–42% of lifetime risk on AD is attributed to environmental factors [5]. Recent studies demonstrated that A β can function as an antimicrobial peptide that protects neurons from bacterial and viral infections [6, 7], supporting the hypothesis that microbial infections – especially herpes simplex virus 1 (HSV-1) – could play a role in the pathogenesis of AD.

Most adults worldwide are infected with the human neurotropic alphaherpesvirus herpes simplex virus 1 (HSV-1). Latent HSV-1 predominantly resides in the somata of pseudounipolar sensory trigeminal ganglion (TG) neurons, which innervate both the orofacial epithelia and the brainstem [8]. Periodic HSV-1 reactivation results in virus spread to the oral mucosa and the central nervous system (CNS). Whereas oral HSV-1 shedding can be both asymptomatic and symptomatic (cold sores), symptomatic HSV-1 infection of the CNS is extremely rare and associated with severe morbidity and mortality (herpes simplex encephalitis; HSE) [9, 10]. Multiple studies have detected HSV-1 DNA in the brain of elderly individuals with and without AD [11–15]. In the brain of AD patients, viral DNA was found to preferentially colocalize with A β plaques [16]. The presence of HSV-1 in brain could contribute to the pathogenesis of AD by inducing the accumulation of intracellular A β , extracellular A β aggregates, and nuclear accumulation and hyperphosphorylation of tau protein, as observed in cultures of mouse primary cortical neurons, human neuroblastoma cells, and human iPSC-derived neuron models [17–24]. In mice, these AD-related pathological changes were found to progressively accumulate with repeated HSV-1 infection and to correlate with cognitive impairment [25]. However, humans are the only natural host of HSV-1. Unlike mice, lytic HSV-1 infection of the human CNS is extremely rare, in part, because human cortical neurons are intrinsically resistant to HSV-1 infection [26]. It remains to be determined whether lytic or latent HSV-1 infection of human neurons in vivo is directly associated with aberrant A β and pTau expression. Therefore, the aim of this study was to investigate the relationship between AD-related neuropathological changes and HSV infection in the TG of latently HSV-1 infected individuals and the brain of individuals with lytic HSV infection of the brain (HSE).

2 | METHODS

2.1 | Human clinical specimens

Temporal cortex samples from AD patients and paired human TG and plasma samples from healthy controls, AD patients, and patients with other neurological diseases were

obtained from the Netherlands Brain Bank (Netherlands Institute for Neuroscience; Amsterdam, the Netherlands). All donors had provided written informed consent for brain autopsy and the use of material and clinical information for research purposes. All study procedures were performed in compliance with relevant laws in the Netherlands and in accordance with the ethical standards of the Declaration of Helsinki. Institutional guidelines were approved by the local ethical committee (VU University Medical Center, Amsterdam, project number 2009/148). AD brain tissues and some of the TG samples were fixed in neutral buffered formalin and embedded in paraffin (FFPE) for histological analysis. Additional TG samples were used to generate single-cell suspensions or snap-frozen for RNA extraction [27]. FFPE brain samples from patients with acute traumatic brain injury (TBI), HSE patients, and a VZV encephalitis patient were obtained for diagnostic purposes and provided by the BioBanks of the Erasmus MC (TBI patients) and Amsterdam University Medical Center (HSE and VZV encephalitis). According to the institutional “Opt-Out” system, which is defined by the National “Code of Good Conduct” [Dutch: Code Goed Gebruik, May 2011], these surplus human brain tissues were available for the current study.

2.2 | Nucleic acid extraction, quantitative TaqMan real-time PCR (qPCR), and *APOE* genotyping

One-tenth of a TG single-cell suspension was used for DNA extraction using the QIAamp DNA Kit (Qiagen). RNA was extracted from homogenized TG using TRIzol (Thermo Fisher Scientific) and the RNeasy Mini Kit including on-column DNase I treatment (Qiagen), as described [28]. cDNA synthesis was performed using 5 μ g total RNA and superscript IV with oligo(dT)12–18 primers (Thermo Fisher Scientific). TaqMan qPCR was performed in duplicate on the 7500 Real-Time PCR system using 2X PCR Universal Master Mix (Applied Biosystems) and primer/probe pairs specific for HSV-1 *US4*, HSV-1 *LAT*, VZV *ORF38*, and the human single-copy gene hydroxymethylbilane synthase [28, 29] (Table S1). *APOE* genotyping was performed using allele ϵ 2-, ϵ 3-, and ϵ 4-specific primer/probe combinations and TaqMan qPCR, as described [30] (Table S1). Results were confirmed by PCR amplification of the *APOE* gene region containing allele-differentiating SNPs rs429358 and rs7412 using PfuUltra II Fusion High-Fidelity DNA Polymerase (Agilent), followed by Sanger sequencing using the BigDye 3.1 Cycle Sequencing Kit on the 3130XL Genetic Analyzer (Applied Biosystems) [30].

2.3 | Immunohistochemical and immunofluorescent staining

FFPE tissue sections were subjected to heat-induced antigen retrieval using citrate buffer or Trilogy™ (Cell Marque).

Immunohistochemical (IHC) staining was performed using the following primary antibodies: monoclonal mouse anti- β -Amyloid (clone 4G8, BioLegend), anti-phosphorylated-Tau (Ser²⁰²/Thr²⁰⁵) (AT8, Invitrogen), and anti-HSV-1 infected cell protein 8 (ICP8) (10A3, Cell Marque). Staining was visualized using biotinylated polyclonal rabbit anti-mouse Ig (Dako) and goat anti-rabbit Ig (Dako), followed by horseradish peroxidase-conjugated streptavidin (Dako) and 3-amino-9-ethylcarbazole substrate. Images were obtained with an Olympus BX51 microscope or by scanning the slides using the Hamamatsu NanoZoomer 2.0 HT.

Immunofluorescent (IF) staining was performed using the following primary antibodies: polyclonal rabbit anti-HSV-1 (cross-reactive with HSV-2; Agilent), anti-Ibal (Wako), or anti-GFAP (Dako), polyclonal chicken anti-GFAP (Abcam), monoclonal mouse anti-NeuN IgG1 (A60, Sigma-Aldrich), anti-MBP (1.B.645, Santa Cruz Biotechnology), anti-Ibal (GT10312, Invitrogen), anti- β -Amyloid (4G8, BioLegend), anti-phosphorylated-Tau (Ser²⁰²/Thr²⁰⁵) (AT8, Invitrogen), anti-CD45 (2B11 + PD7/26, Dako), and monoclonal rat anti-CD3 (CD3-12, Abcam). Alexa Fluor® 488 (AF488)-, AF594-, and AF647-conjugated polyclonal goat anti-rabbit IgG, anti-mouse IgG1, and IgG2b, goat anti-chicken IgY and goat anti-rat IgG secondary antibodies (Invitrogen) were used. Nuclei were stained with Hoechst 33342 Solution and mounted with ProLong™ Diamond Antifade Mountant (Thermo Fisher Scientific). Images were obtained using a Zeiss LSM700 confocal microscope.

2.4 | In situ hybridization

Tissue sections were stained for HSV-1 and VZV RNA by in situ hybridization (ISH), using the RNAscope® 2.5 HD Kit-RED and probes HSV-1-LAT (Cat No. 315651) and VZV-Pool (Cat No. 400701) (Advanced Cell Diagnostics). Slides were counterstained with hematoxylin (Sigma) and mounted with EcoMount (Biocare Medical).

2.5 | Statistical analyses

All statistical analyses were performed using GraphPad Prism 8.0.2 (GraphPad Software Inc).

3 | RESULTS

3.1 | HSV-1 DNA load and LAT expression in the TG of AD patients and controls

Previous studies suggest that HSV-1 infection or reactivation, as measured by plasma HSV-1 IgG and IgM levels, could be a risk factor for AD development [31].

Clinical HSV-1 reactivation frequency correlates with both *APOE4* carriage (humans) and latent viral DNA load (mice) [32–35]. To investigate whether latent HSV-1 DNA load was associated with AD development or *APOE* genotype, we performed qPCR on human TG that were infected with HSV-1 and obtained from AD patients and controls. Additionally, we performed qPCR to detect the closely related varicella-zoster virus (VZV), because most human TG is co-infected by HSV-1 and VZV [10]. HSV-1 and VZV DNA loads were similar between the TG obtained from AD patients and controls (Figure 1A). Further, *APOE* genotyping of all analyzed TG specimens demonstrated comparable HSV-1 and VZV DNA levels in *APOE4* carriers and non-carriers (Figure 1B and Figure S1), despite the reported effect of *APOE4* on promoting HSV neurovirulence [36, 37]. We performed RT-qPCR and ISH to quantify expression of the HSV-1 latency-associated transcripts (LAT) in human TG. Similar to HSV-1 DNA loads, the prevalence of LAT-positive neurons and abundance of LAT RNA were comparable between AD patients and controls (Figure 1C).

3.2 | No expression of A β and pTau in latently HSV-infected TG neurons of AD patients and controls

Most individuals acquire HSV-1 infection in their childhood, followed by a lifelong latent infection and frequent (asymptomatic) virus reactivation [38, 39]. Given that HSV-1 replication leads to the accumulation of A β and increased pTau expression in cultured neurons and in the brains of HSV-infected mice [17, 20, 21, 25, 40], we analyzed whether latent HSV-1 infection was associated with A β or pTau expression in TG neurons of AD patients and controls (Table S2). Analysis of consecutive TG sections from HSV IgG seropositive ($n = 3$ AD; $n = 5$ controls) and HSV IgG seronegative ($n = 2$ AD) donors did not demonstrate A β , A β plaques, or pTau staining in latently HSV-1-infected TG neurons of either AD patients or controls (Figure 1D). Similarly, we did not observe aberrant A β or pTau staining in neurons not infected by HSV-1.

3.3 | Expression of A β and pTau in brain of HSE patients

As we did not observe aberrant A β or pTau expression in HSV-infected peripheral TG neurons, we hypothesized that HSV-induced AD-related pathological changes could be (1) restricted to CNS neurons, or (2) reversible and therefore limited to lytic (productive) virus infection. To test these hypotheses, we acquired rare post-mortem brain specimens from five HSE patients (Table 1), four cases of HSV-1 encephalitis and




FIGURE 1 HSV-1 infection is not associated with aberrant A β or pTau expression in latently infected human trigeminal ganglia (TG) neurons. (A, B) HSV-1- and VZV-specific qPCR was performed on DNA extracted from the TG of Alzheimer's disease (AD) patients and control subjects, stratified on disease status (A; 20 controls and 10 AD patients) or *APOE4* allele carrier status (B; HSV-1: 22 *APOE4*-negative and 8 *APOE4*-positive individuals; VZV: 24 *APOE4*-negative and 13 *APOE4*-positive individuals). Horizontal line: median. (C) Detection of LAT RNA by RT-qPCR (7 AD patients and 7 controls) and ISH (11 controls and 5 AD patients; 3 sections per donor analyzed) in human TG from AD patients and control subjects. (D) Sequential TG sections from AD ($n = 4$) and control ($n = 6$) subjects were stained for amyloid β protein, HSV-1 latency-associated transcript (LAT) RNA and phosphorylated Tau protein (pTau; Ser²⁰²/Thr²⁰⁵) by immunohistochemistry (IHC) and RNA in situ hybridization (ISH). Arrowheads indicate LAT-positive neurons and asterisks indicate lipofuscin granules. Scale bar: 50 μ m

one case of neonatal HSV-2 encephalitis. Brain sections from all patients contained lytically HSV-infected cells, as evidenced by the detection of intranuclear eosinophilic (Cowdry type A) inclusion bodies, HSV-1 LAT RNA (in all HSV-1 HSE patients) and HSV ICP8 protein (Figure 2A). In all patients, the majority of HSV-infected ICP8^{POS} cells were identified as NeuN^{POS} neurons, with lower frequencies of infected non-neuronal cells, mainly Iba1^{POS} microglia (Figure 2B,C). Thus, brain sections from HSE patients provide a snapshot of the appearance of lytically HSV-infected CNS neurons in vivo.

We then investigated whether lytic HSV infection was associated with increased intracellular production of A β or pTau, or the deposition of extracellular A β plaques in brain sections from HSE patients (Table 1). As a control, we included brain sections from patients with acute traumatic brain injury (TBI, $n = 3$; Table S3) and AD patients ($n = 3$; Table S4). TBI-induced neuronal damage is associated with increased production of A β , resulting in its accumulation in axonal spheroids as well as plaques in about 30% of patients [41]. Indeed, we observed axonal spheroids in all three TBI patients and increased expression of intracellular A β protein and

TABLE 1 HSE patient characteristics and Alzheimer's disease-related pathology

Patient	Age ^a	Gender ^b	Causative virus	Brain regions analyzed	A β intraneuronal ^c	A β plaques	NFT
1	–	–	HSV-1	Cortex	No	No	No
2	79	F	HSV-1	Hippocampus/cortex	Yes	Yes	Yes
				Amygdala/entorhinal Cortex	Yes	Yes	Yes
				Cortex	Yes	Yes	Yes
				Insula	Yes	Yes	Yes
3	0 ^d	F	HSV-2	Cortex	No	No	No
4	81	F	HSV-1	Putamen	No	Yes	No
5	76	M	HSV-1	Hippocampus/entorhinal cortex	No	Yes	Yes

Abbreviation: NFT, neurofibrillary tangle.

^aAge in years.

^bF, female; M, male.

^cIntraneuronal A β accumulation.

^dNeonatal HSE patient, 2 wk old.




FIGURE 2 Lytic HSV infection of neurons in the brain of herpes simplex virus encephalitis (HSE) patients. (A) Brain sections from HSE patients (Table 1) were stained with hematoxylin and eosin, or stained for HSV-1 LAT RNA by ISH and HSV-1 ICP8 protein by IHC. Boxes indicate the area shown at higher magnification in the inset. Scale bar: 50 μ m. (B, C) Brain sections of HSE patients were immunofluorescently stained for (B) HSV-1 protein (green), neurons (NeuN; white) and oligodendrocytes (MBP; red), as well as (C) HSV-1 protein (green), microglia (Iba1; white), and astrocytes (GFAP; red). Nuclei were stained with Hoechst-33342 (blue). Representative images are shown for patient #2 (amygdala/entorhinal cortex; Table 1). Scale bar: 20 μ m

(diffuse) extracellular A β plaques in two TBI patients (Figure 3A). NFTs and pTau staining was not observed in TBI patients (Figure 3A). Abundant intracellular A β , extracellular A β plaques, and NFT were observed

in all three AD patients (Figure 3A). By contrast, brain sections from two of five HSE patients showed neither A β protein/plaques nor pTau protein/NFTs (Table 1). A β plaques were observed in one of five HSE patients




FIGURE 3 Lytic HSV infection is not consistently associated with aberrant Aβ or pTau expression in the brain of herpes simplex virus encephalitis (HSE) patients. (A) Brain sections from Alzheimer's disease (AD; patient 1, Table S4), trauma brain injury (TBI; patient 2, Table S3), and HSE patients (Table 1) were stained for Aβ and pTau (Ser²⁰²/Thr²⁰⁵) by IHC. Filled and open arrowheads indicate mature senile and diffuse Aβ plaques, respectively. Filled black arrows indicate intracellular Aβ accumulations. Open arrows indicate neurofibrillary tangles. Boxes indicate the area shown at higher magnification in the inset. Scale bar: 50 μm. (B) Consecutive brain sections from HSE patients were stained for Aβ and pTau (Ser²⁰²/Thr²⁰⁵) protein by IHC and HSV-1 RNA by ISH. Boxes indicate the area shown at a higher magnification. Scale bars indicate 500 μm (panels 1), 100 μm (panels 2), and 50 μm (panels 3)

(patient #2), whereas intracellular accumulations of Aβ protein were detected in three of five HSE patients. NFTs were detected in two of five HSE patients (patients #2 and #5) (Figure 3A and Table 1).

Subsequently, we determined the spatial relationship between HSV-infected cells and AD-related pathological changes. Although HSV-infected cells were occasionally observed in close proximity to NFT and/or Aβ plaques,

majority of these AD-related pathological changes were not associated with sites of HSV replication in the brains of HSE patients #2 and #5 (Figure 3B). Similarly, intracellular accumulations of Aβ protein were widely spread through brain sections of HSV-1 HSE patients and not restricted to areas with HSV-1 infection (Figure 3B and Figure S2). IF staining was performed to investigate potential colocalization between HSV-1 infected cells and

plaques/NFT and to assess the intracellular expression of A β and pTau in more detail. Again, occasional HSV-infected cells were observed adjacent to A β plaques and NFT, but HSV antigen never colocalized with A β plaques nor NFT (Figure 4A). In HSE patients #2, #4, and #5, abundant intracellular A β protein staining was detected not only in most HSV-infected neurons but also in non-infected neurons (Figure 4B). Nuclear pTau staining was observed in some HSV-infected cells in HSE patients containing NFTs (Figure 4C). However, we did not observe increased A β or pTau expression in HSV-infected neurons in HSE patients #1 and #3 (Figure 4D).

Thereafter, we investigated whether the widespread increased expression of intracellular A β protein without diffuse A β plaques in the brain of some HSE patients was specific to HSV infection. For this, we studied A β and

pTau expression in brain sections from a VZV encephalitis patient. The patient was a 72-year-old male who had no known history of dementia and died 15 days after the onset of disease. Abundant VZV RNA-positive cells were detected (Figure S3A), as well as VZV antigen-positive cells, indicative of lytic VZV infection. A β plaques and NFT, as well as intracellular A β protein, were observed in hippocampal sections, whereas only widespread prominent intracellular A β protein depositions were observed in medulla oblongata sections (Figure S3B). Similar to the HSE brains, we did not observe an association between the spatial relationship of virus-infected cells and AD-related pathological changes (Figure S3C). Overall, these results suggest that lytic α -herpesvirus, especially HSV, infection of human CNS neurons is not directly associated with aberrant A β or pTau protein expression.




FIGURE 4 HSV-infected cells do not consistently express aberrant A β or pTau in the brain of herpes simplex encephalitis (HSE) patients. Brain sections from HSE patients were immunofluorescently stained for HSV protein (green), A β (red), and pTau (Ser²⁰²/Thr²⁰⁵; white). Nuclei were stained with Hoechst-33342 (blue). Images are shown for HSE patients #2 (A left panel), #5 (A right panel, and C), #4 (B), and #3 (D) (Table 1). White boxes indicate the areas shown at higher magnification. Scale bars indicate 10 μ m (A: Amyloid β plaques; and C) and 20 μ m (A right panel, B and D)

3.4 | Expression of A β and pTau in brain of an AD patient with concurrent HSE

A previous study reported that HSV-1 DNA colocalizes with A β plaques in brain of AD patients [16]. To investigate whether pre-existing AD pathology impacts the relationship between HSV-infected cells and A β and pTau expression, we obtained rare brain tissue specimens from an AD patient presenting with HSE. The patient was a 63-year-old female diagnosed with AD who developed

HSE and died from septic shock resulting from HSE and aspiration pneumonia 1.5 wk after hospitalization. Analysis of six cortical brain regions revealed intermediate AD-related pathological changes, that is, amyloid Thal phase 5/5 and Braak NFT stage 4/6 (A3B2), consistent with dementia. HSV antigen-positive cells were abundantly detected in insular and temporal cortical tissue sections and less prominent in parietal and occipital cortex sections (Figure 5A). Importantly, HSV-1 infected cells were often found in close proximity to NFT and




FIGURE 5 Detection of HSV-infected cells in close proximity to A β plaques in the brain of an Alzheimer's disease patient with herpes simplex encephalitis. (A) Brain section stained for HSV protein (brown) by immunohistochemistry (IHC). Scale bar indicates 50 μ m. (B) Consecutive brain sections were stained for A β (4G8) and pTau (Ser²⁰², Thr²⁰⁵) by IHC or stained for HSV-1 RNA by in situ hybridization (ISH). Boxes indicate areas shown at higher magnification. Asterisks indicate the same blood vessel in consecutive sections. Scale bars indicate 500 μ m (low magnification) and 100 μ m (high magnification). (C, D) Brain sections were immunofluorescently stained for HSV protein (green), A β (red), and pTau (Ser²⁰²/Thr²⁰⁵; white) protein. Nuclei were stained with Hoechst-33342 (blue). Scale bar: 20 μ m

especially A β plaques (Figure 5B). However, AD-related pathological changes were not different between regions with and without HSV-1 infected cells. HSV-1 infected cells were situated around but did not colocalize with A β plaques and NFTs (Figure 5C). Similar to the other HSE patients, increased expression of intracellular A β protein was present in most HSV-infected cells, as well as non-infected cells (Figure 5D). Thus, AD and HSV infection may affect the same brain regions, but we did not observe enhanced AD-related pathology proximate to HSV infection.

3.5 | Neuroinflammation in HSE patients with and without AD-related pathology

Because neuroinflammation is a major contributor to neurodegeneration in AD [42], we hypothesized that inflammatory cells rather than HSV replication itself could be associated with specific AD-related pathological changes. Brain tissue sections of all HSE patients, including the AD patient with HSE, indeed demonstrated widespread inflammation and gliosis in comparison to AD patients without viral encephalitis (Figure 6A and Table S4). HSE patients and the AD patient with concurrent HSE showed prominent perivascular cuffs, mainly composed of CD45^{POS} CD3^{NEG} Iba1^{NEG} mononuclear cells – most likely macrophages or histiocytes – and lower numbers of CD45^{POS} CD3^{POS} T-cells (Figure 6B). Astrogliosis was observed in the brain parenchyma of all HSE patients, irrespective of A β or pTau expression pattern. This was also observed in the AD patient with HSE, but not as extensively in the AD patients without HSE (Figure 6C). Notably, while astrocytes colocalized with A β plaques of all AD patients and HSE patient #2, more intense GFAP staining was observed in astrocytes entangling A β plaques in the AD patient with HSE (Figure S4). Additionally, extensive microgliosis was observed in the brain parenchyma of the HSE patients and AD patient with HSE, but not AD patients without HSE (Figure 6D). Both amoeboid (Figure 6D) and ramified microglia (Figure S5) were observed throughout the tissue sections and in association with A β plaques and/or pTau in the AD patient with concurrent HSE and the HSE patients #2 and #5 (Table 1). Microglia were mostly amoeboid in HSE patients #1 and #3, whereas predominantly ramified microglia were present in HSE patient #4 (Figure 6D). Thus, although lytic HSV infection of the CNS induced significant neuroinflammation, the overall pattern of inflammation and gliosis was not associated with specific A β or pTau expression patterns.

4 | DISCUSSION

Cumulative evidence has led to the hypothesis that the endemic human neurotropic virus, HSV-1, could play

a role in the pathogenesis of AD [17–19, 21, 22, 25, 43]. Here, we used clinical specimens from HSV-infected individuals to investigate the relationship between HSV infection and A β and pTau expression within human neurons in TG and brain. We report that latent HSV infection is not associated with A β or pTau expression in TG neurons located in the PNS. Similarly, lytic HSV infection in the CNS was not consistently associated with increased expression of intracellular A β or pTau proteins nor with the deposition of A β plaques and NFT. Analysis of unique material from an AD patient with concurrent HSE showed that the same brain regions are often affected by HSV and AD-related pathology, but did not reveal exacerbated A β or pTau production proximate to regions containing virus-infected cells.

Previous studies showed that increased anti-HSV IgM plasma levels and possibly anti-HSV IgG avidity can be a measurement of HSV reactivation that correlates with AD development [31, 44]. One proposed explanation is that genetic risk factors associated with AD development, especially *APOE4* and paired immunoglobulin-like type 2 receptor alpha (*PILRA*), can directly impact lytic HSV-1 infection. The *APOE4* allele is associated with increased frequency of HSV-1 DNA detection in the brain of AD patients and symptomatic HSV-1 reactivation (cold sores) [45, 46]. By contrast, asymptomatic oral HSV-1 shedding was not affected by *APOE* genotype [41], suggesting that *APOE4* does not directly influence virus reactivation but may play a role in peripheral control of HSV infection. In this study, we did not observe differences in latent HSV-1 DNA load nor LAT expression in the TG of AD patients compared to control subjects without AD, or in *APOE4* carriers compared to non-carriers. These data support the hypothesis that latent HSV-1 infection is not directly associated with increased risk of AD development.

Detailed analysis of rare human HSE brain specimens demonstrated that lytic HSV infection was not associated with increased levels of intraneuronal A β or A β plaque deposition in human neurons in vivo. Although we observed accumulation of intraneuronal A β protein in three HSE and one VZV encephalitis patient, A β depositions were widely present in both virus-infected and non-infected neurons and tissue sections. These findings are consistent with the ubiquitous and progressive accumulation of intraneuronal A β in the absence of extracellular A β plaques and NFT that is observed with aging [47]. Similarly, we only observed A β plaques in one HSE patient and locally (hippocampus, but not medulla oblongata) in one VZV encephalitis patient. The elderly age of these two patients, lack of an association between sites of virus replication and A β plaques, and the presence of both senile plaques and NFT suggests that these pathological changes were most likely part of ongoing development of AD, rather than induced by HSV infection.




FIGURE 6 Neuroinflammation is similar in herpes simplex encephalitis (HSE) patients with and without A β plaques and/or NFT. Brain sections from Alzheimer's disease (AD) patients ($n = 3$, Table S4), HSE patients ($n = 4$, Table 1), and an AD with HSE patient ($n = 1$) were stained with (A) hematoxylin and eosin staining or IF stained for: (B) CD3 (green) and GFAP (white) (left panel) or Iba1 (green) and CD45 (white) (right panel); (C) A β (red) and GFAP (white); or (D) A β (red), Iba1 (green), and CD45 (white). Nuclei were stained with Hoechst-33342 (blue). Images are shown for the AD patient with HSE (B–D), AD patient #1 (C, D; Table S4) and indicated HSE patients. Boxes indicate the areas shown at higher magnification. Scale bar: 50 μ m

HSV particles bind to A β protein and catalyze A β 42 oligomerization, leading to A β aggregates that physically entrap virus particles [7, 48]. Consistent with the idea of HSV initiating the deposition of A β plaques in the CNS, a previous study used in situ PCR to detect HSV-1 DNA in the brain of AD patients and controls, and reported that HSV-1 preferentially colocalizes with A β plaques in AD patients [12]. Here, we did not detect viral antigen nor RNA within A β plaques in the brain of elderly

individuals with HSV-1 or VZV encephalitis. Although HSV-infected cells were often found in close proximity to A β plaques and/or NFT in the brain of the AD patient with HSE, AD-related pathology was not restricted to regions with HSV-infected cells. We also did not observe HSV antigen or RNA in A β plaques in this patient. HSV-1 infection rapidly induces A β plaque deposition in the brain of genetically susceptible AD mouse models, but repeated HSV-1 reactivation and lytic virus replication in



brain results in the progressive accumulation of both senile plaques and NFT [25]. While HSE is extremely rare in humans, asymptomatic HSV reactivation and viral spread to the CNS, as measured by the presence of viral DNA, occur frequently [9, 10, 16, 45]. Our data suggest that lytic HSV infection does not directly induce the formation of A β plaques in the human brain. This is possible because of the extensive neuronal cell death observed in HSE patients [49], which may limit the time available for neurons to produce substantial A β aggregates. Instead, repeated exposure to abortive HSV infections may be required to induce persistent A β plaques or could enhance A β oligomerization in existing plaques.

HSV infection of diverse human neuron models in vitro leads to the hyperphosphorylation of tau protein, including Ser²⁰²/Thr²⁰⁵, by cellular cyclin-dependent kinases and relocation of pTau to the nucleus [17–19]. We report that HSV-infected neurons in the brain of human HSE patients occasionally expressed nuclear pTau but only in individuals presenting with NFT. However, NFTs were not restricted to areas containing HSV-infected cells in HSE patients and the AD patient with HSE. These findings are in agreement with prior studies in 3xTg-AD mice (prone to develop NFT [50]), in which the degree and residues involved in tau phosphorylation in response to HSV infection varied between brain regions, and progressively increased with repeated viral reactivation events [25]. The heterogeneity of neurons in vivo – including the expression of specific cyclin-dependent kinases – could determine their differential susceptibility to HSV-induced aberrant tau phosphorylation and cellular localization.

Specific subsets of microglia and astrocytes colocalize with A β plaques in the brain of AD patients [51–53] and are thought to be involved in both early and advanced stages of AD pathogenesis [54, 55]. Lytic HSV infection induces robust innate immune responses in the brain, involving both microglia and astrocytes, which not only control ongoing virus replication but also causes immunopathology [56]. Prominent microgliosis and astrogliosis were present in all HSE patients, but we did not observe overall differences in glia cell morphology or abundance in patients with or without A β plaques and/or NFT. Interestingly, GFAP staining tended to be more abundant in A β plaque-associated astrocytes in brain sections from the patient with combined AD and HSE, compared to patients with either AD or HSE (Figure S4). Although these observations suggest that HSV infection may influence reactive astrocyte function in the brain of AD patients, more detailed analyses in more patients or experimental animal models are warranted.

In conclusion, we demonstrate that latent and lytic HSV infection of human neurons in TG and CNS is not consistently associated with aberrant A β or pTau expression. Human in vitro neuron cultures and murine AD models highlight potential mechanisms by which HSV infection could contribute to the initiation

or perpetuation of AD. However, our data suggest that the human CNS could be more resilient to HSV-induced AD-related neuropathology than previously anticipated. Future studies comparing the effects of HSV infection on the healthy aging brain and AD brain may provide valuable insight into mechanisms by which HSV may affect AD pathogenesis.

ACKNOWLEDGMENTS

We thank Dr. Georges Verjans for the critical discussion of the data and Tamana Khemai-Mehraban for technical assistance. Research reported in this publication was in part supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award Number R01AI151290 (W.J.D.O.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. This study was funded in part by a Human Disease Model Award 2020 (Erasmus MC).

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

AUTHOR CONTRIBUTIONS

WJDO conceived and designed the study. DNT and ATCMB conducted the experiments. DNT, ATCMB, VvD, RMV and WJDO analyzed and interpreted the data. EA and RMV contributed patient samples. DNT and WJDO wrote the manuscript.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Werner J. D. Ouwendijk  <https://orcid.org/0000-0001-8393-296X>

REFERENCES

- 2020 Alzheimer's disease facts and figures. *Alzheimers Dement.* 2020;16(3):391–460.
- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science.* 2002;297(5580):353–6.
- Hardy JA, Higgins GA. Alzheimer's disease: the amyloid Alzheimer's disease. *Science.* 1992;256:184–5.
- Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C, et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet.* 2013;45(12):1452–8.
- Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet Neurol.* 2011;10(9):819–28.
- Kumar DKV, Choi SH, Washicosky KJ, Eimer WA, Tucker S, Ghofrani J, et al. Amyloid- β peptide protects against microbial infection in mouse and worm models of Alzheimer's disease. *Sci Transl Med.* 2016;340(8):340ra72.
- Eimer WA, Vijaya Kumar DK, Navalpur Shanmugam NK, Rodriguez AS, Mitchell T, Washicosky KJ, et al. Alzheimer's

- disease-associated β -amyloid is rapidly seeded by herpesviridae to protect against brain infection. *Neuron*. 2018;99(1):56–63.
8. Smith G. Herpesvirus transport to the nervous system and back again. *Annu Rev Microbiol*. 2012;66(1):153–76.
 9. Roizman B, Whitley RJ. An inquiry into the molecular basis of HSV latency and reactivation. *Annu Rev Microbiol*. 2013;67:355–74.
 10. Arvin A, Campadelli-Fiume G, Mocarski E, Moore PS, Roizman B, Whitley R, et al. Human herpesviruses: biology, therapy, and immunoprophylaxis. Cambridge University Press; 2007.
 11. Baringer JR, Pisani P. Herpes simplex virus genomes in human nervous system tissue analyzed by polymerase chain reaction. *Ann Neurol*. 1994;36(6):823–9.
 12. Itzhaki RF. Herpes and Alzheimer's Disease: Subversion in the Central Nervous System and How It Might Be Halted. *J Alzheimer's Dis*. 2016;54(4):1273–81. <https://doi.org/10.3233/JAD-160607>
 13. Gordon L, McQuaid S, Cosby SL. Detection of herpes simplex virus (types 1 and 2) and human herpesvirus 6 DNA in human brain tissue by polymerase chain reaction. *Clin Diagn Virol*. 1996;6(1):33–40.
 14. Readhead B, Haure-Mirande JV, Funk CC, Richards MA, Shannon P, Haroutunian V, et al. Multiscale analysis of independent Alzheimer's cohorts finds disruption of molecular, genetic, and clinical networks by human herpesvirus. *Neuron*. 2018;99(1):64–82.e7. <https://doi.org/10.1016/j.neuron.2018.05.023>
 15. Hemling N, Røytta M, Rinne J, Pöllänen P, Broberg E, Tapio V, et al. Herpesviruses in brains in Alzheimer's and Parkinson's diseases. *Ann Neurol*. 2003;54(2):267–71.
 16. Wozniak MA, Mee AP, Itzhaki RF. Herpes simplex virus type 1 DNA is located within Alzheimer's disease amyloid plaques. *J Pathol*. 2009;217(1):131–8.
 17. Zambrano A, Solis L, Salvadores N, Cortes M, Lerchundia R, Otthb C. Neuronal cytoskeletal dynamic modification and neurodegeneration induced by infection with herpes simplex virus type 1. *J Alzheimers Dis*. 2008;14:259–69.
 18. Wozniak MA, Frost AL, Itzhaki RF. Alzheimer's disease-specific tau phosphorylation is induced by herpes simplex virus type 1. *J Alzheimers Dis*. 2009;16(2):341–50.
 19. Álvarez G, Aldudo J, Alonso M, Santana S, Valdivieso F. Herpes simplex virus type 1 induces nuclear accumulation of hyperphosphorylated tau in neuronal cells. *J Neurosci Res*. 2012;90(5):1020–9.
 20. Ill-Raga G, Palomer E, Wozniak MA, Ramos-ferna E, Antu C, Tajés M, et al. Activation of PKR causes amyloid β -peptide accumulation via de-repression of BACE1 expression. *PLoS One*. 2011;6(6):1–10.
 21. Wozniak MA, Itzhaki RF, Shipley SJ, Dobson CB. Herpes simplex virus infection causes cellular β -amyloid accumulation and secretase upregulation. *Neurosci Lett*. 2007;429:95–100.
 22. Shipley SJ, Parkin ET, Itzhaki RF, Dobson CB. Herpes simplex virus interferes with amyloid precursor protein processing. *BMC Microbiol*. 2005;5:1–8.
 23. Cairns DM, Rouleau N, Parker RN, Walsh KG, Gehrke L, Kaplan DL. A 3D human brain-like tissue model of herpes-induced Alzheimer's disease. *Sci Adv*. 2020;6(19):1–14.
 24. Abrahamson EE, Zheng W, Muralidaran V, Ikonovic MD, Bloom DC, Nimgaonkar VL, et al. Modeling A β 42 accumulation in response to herpes simplex virus 1 infection: two dimensional or three dimensional? *J Virol*. 2020;95(5):1–9.
 25. de Chiara G, Piacentini R, Fabiani M, Mastrodonato A, Marcocci ME, Limongi D, et al. Recurrent herpes simplex virus-1 infection induces hallmarks of neurodegeneration and cognitive deficits in mice. *PLoS Pathog*. 2019;15(3):1–30.
 26. Lafaille FG, Harschnitz O, Lee YS, Zhang P, Hasek ML, Kerner G, et al. Human SNORA31 variations impair cortical neuron-intrinsic immunity to HSV-1 and underlie herpes simplex encephalitis. *Nat Med*. 2019;25(12):1873–84.
 27. Verjans GMGM, Hintzen RQ, van Dun JM, Poot A, Milikan JC, Laman JD, et al. Selective retention of herpes simplex virus-specific T cells in latently infected human trigeminal ganglia. *Proc Natl Acad Sci U S A*. 2007;104(9):3496–501.
 28. Depledge DP, Ouwendijk WJD, Sadaoka T, Braspenning SE, Mori Y, Cohrs RJ, et al. A spliced latency-associated VZV transcript maps antisense to the viral transactivator gene. *Nat Commun*. 2018;9(1):1–12. <https://doi.org/10.1038/s41467-018-03569-2>
 29. van Doornum GJJ, Guldemeester J, Osterhaus ADME, Niesters HGM. Diagnosing herpesvirus infections by real-time amplification and rapid culture. *J Clin Microbiol*. 2003;41(2):576–80.
 30. Zhong L, Xie Y-Z, Cao T-T, Wang Z, Wang T, Li X, et al. A rapid and cost-effective method for genotyping apolipoprotein E gene polymorphism. *Mol Neurodegener*. 2016;11:2.
 31. Lövheim H, Gilthorpe J, Adolffson R, Nilsson LG, Elgh F. Reactivated herpes simplex infection increases the risk of Alzheimer's disease. *Alzheimers Dement*. 2015;11(6):593–9.
 32. Hoshino Y, Pesnicak L, Cohen JI, Straus SE. Rates of reactivation of latent herpes simplex virus from mouse trigeminal ganglia ex vivo correlate directly with viral load and inversely with number of infiltrating CD8+ T cells. *J Virol*. 2007;81(15):8157–64.
 33. Linard M, Letenneur L, Garrigue I, Doize A, Dartigues JF, Helmer C. Interaction between APOE4 and herpes simplex virus type 1 in Alzheimer's disease. *Alzheimers Dement*. 2020;16(1):200–8.
 34. Letenneur L, Pères K, Fleury H, Garrigue I, Barberger-Gateau P, Helmer C, et al. Seropositivity to herpes simplex virus antibodies and risk of Alzheimer's disease: a population-based cohort study. *PLoS One*. 2008;3(11):1–5.
 35. Lopatko Lindman K, Weidung B, Olsson J, Josefsson M, Kok E, Johansson A, et al. A genetic signature including apolipoprotein E ϵ 4 potentiates the risk of herpes simplex-associated Alzheimer's disease. *Alzheimers Dement*. 2019;5:697–704.
 36. Burgos JS, Ramirez C, Sastre I, Valdivieso F. Effect of apolipoprotein e on the cerebral load of latent herpes simplex virus type 1 DNA. *J Virol*. 2006;80(11):5383–7.
 37. Burgos JS, Ramirez C, Sastre I, Bullido MJ, Valdivieso F. ApoE4 is more efficient than E3 in brain access by herpes simplex virus type 1. *NeuroReport*. 2003;14(14):1825–7.
 38. Mark KE, Wald A, Magaret AS, Selke S, Olin L, Huang M-L, et al. Rapidly cleared episodes of herpes simplex virus reactivation in immunocompetent adults. *J Infect Dis*. 2008;198(8):1141–9.
 39. Smith JS, Robinson NJ. Age-specific prevalence of infection with herpes simplex virus types 2 and 1: a global review. *J Infect Dis*. 2002;186:S3–28.
 40. Martin C, Aguila B, Araya P, Vio K, Valdivia S, Zambrano A, et al. Inflammatory and neurodegeneration markers during asymptomatic HSV-1 reactivation. *J Alzheimers Dis*. 2014;39(4):849–59.
 41. Johnson VE, Stewart W, Smith DH. Traumatic brain injury and amyloid- β pathology: a link to Alzheimer's disease? *Nat Rev Neurosci*. 2010;11(5):361–70. Available from: <http://www.nature.com/articles/nrn2808>
 42. Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT. Inflammation as a central mechanism in Alzheimer's disease. *Alzheimers Dement*. 2018;4:575–90. <https://doi.org/10.1016/j.trci.2018.06.014>
 43. Ashraf GM, Tarasov VV, Makhmutova A, Chubarev VN, Avila-Rodriguez M, Bachurin SO, et al. The possibility of an infectious etiology of Alzheimer disease. *Mol Neurobiol*. 2019;56(6):4479–91.
 44. Agostini S, Mancuso R, Baglio F, Cabinio M, Hernis A, Costa AS, et al. High avidity HSV-1 antibodies correlate with absence of amnesic Mild Cognitive Impairment conversion to Alzheimer's disease. *Brain Behav Immun*. 2016;58:254–60. Available from: <https://www.sciencedirect.com/science/article/pii/S0889159116303476>

45. Itzhaki RF, Lin W, Shang D, Wilcock GK, Faragher B, Jamieson GA. Early reports Herpes simplex virus type 1 in brain and risk of Alzheimer's disease. *FASEB J*. 1997;349:241–4.
46. Koelle DM, Magaret A, Warren T, Schellenberg GD, Wald A. APOE genotype is associated with oral herpetic lesions but not genital or oral herpes simplex virus shedding. *BoneSex Transm Infect*. 2010;86(3):202–6.
47. Welikovich LA, Do Carmo S, Maglóczy Z, Szocsics P, Löke J, Freund T, et al. Evidence of intraneuronal A β accumulation preceding tau pathology in the entorhinal cortex. *Acta Neuropathol*. 2018;136(6):901–17. <https://doi.org/10.1007/s00401-018-1922-z>
48. Ezzat K, Pernemalm M, Pålsson S, Roberts TC, Järver P, Dondalska A, et al. The viral protein corona directs viral pathogenesis and amyloid aggregation. *Nat Commun*. 2019;10(1):1–16. <https://doi.org/10.1038/s41467-019-10192-2>
49. DeBiasi RL, Kleinschmidt-DeMasters BK, Richardson-Burns S, Tyler KL. Central nervous system apoptosis in human herpes simplex virus and cytomegalovirus encephalitis. *J Infect Dis*. 2002;186(11):1547–57.
50. Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, Kaye R, et al. Triple-transgenic model of Alzheimer's Disease with plaques and tangles: Intracellular A β and synaptic dysfunction. *Neuron*. 2003;39(3):409–21.
51. Habib N, McCabe C, Medina S, Varshavsky M, Kitsberg D, Dvir-Szternfeld R, et al. Disease-associated astrocytes in Alzheimer's disease and aging. *Nat Neurosci*. 2020;23(6):701–6. <https://doi.org/10.1038/s41593-020-0624-8>
52. Olah M, Menon V, Habib N, Taga MF, Ma Y, Yung CJ, et al. Single cell RNA sequencing of human microglia uncovers a subset associated with Alzheimer's disease. *Nat Commun*. 2020;11(1):6129. <https://doi.org/10.1038/s41467-020-19737-2>
53. Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R, Ulland TK, et al. A unique microglia type associated with restricting development of Alzheimer's disease. *Cell*. 2017;169(7):1276–1290.e17. <https://doi.org/10.1016/j.cell.2017.05.018>
54. Rodríguez-Arellano JJ, Parpura V, Zorec R, Verkhratsky A. Astrocytes in physiological aging and Alzheimer's disease. *Neuroscience*. 2016;323:170–82. <https://doi.org/10.1016/j.neuroscience.2015.01.007>
55. Bates KA, Fonte J, Robertson TA, Martins RN, Harvey AR. Chronic gliosis triggers Alzheimer's disease-like processing of amyloid precursor protein. *Neuroscience*. 2002;113(4):785–96.
56. Mancini M, Vidal SM. Insights into the pathogenesis of herpes simplex encephalitis from mouse models. *Mamm Genome*. 2018;29(7–8):425–45. <https://doi.org/10.1007/s00335-018-9772-5>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

TABLE S1 Primers and probes used in this study

TABLE S2 FFPE TG samples used in this study

TABLE S3 TBI patients used in this study

TABLE S4 FFPE AD samples used in this study

FIGURE S1 Quantification of latent HSV-1 and VZV DNA load in human TG stratified on *APOE* genotype. HSV-1- and VZV-specific qPCR and *APOE* genotyping was performed on DNA extracted from the trigeminal ganglia (TG) of AD patients and controls

FIGURE S2 Intraneuronal accumulation of A β protein is not restricted to areas of HSV-1 infection in brains of HSE patients. Consecutive slides were stained for HSV-1 RNA by ISH and A β by IHC. A β can be seen in areas with HSV-1 RNA (A) and in areas without HSV-1 RNA (B). Data shown for HSE donor 5. Scale bar: 500 μ m (A) and 250 μ m (B)

FIGURE S3 Lytic VZV infection is not associated with A β plaques or NFT in brain of a VZV encephalitis patient. (A) Brain section stained for VZV RNA by ISH. Box indicates area shown at higher magnification. Open arrowhead indicates examples of VZV RNA-expressing cells. Scale bars indicate 250 μ m (top) and 50 μ m (bottom). (B) Brain sections IHC stained for A β and pTau (Ser202/Thr205). Scale bars indicate 50 μ m. (C) Consecutive brain sections were IHC stained for A β and pTau (Ser202/Thr205) or stained for HSV-1 RNA by ISH. Boxes indicate areas shown at higher magnification. Filled arrow indicates intracellular A β protein, open arrow indicates NFT and open arrowhead indicates VZV RNA-expressing cells. Scale bars indicate 50 μ m (A β , high magnification), 100 μ m (A β , low magnification; pTau high magnification) and 500 μ m (pTau, low magnification)

FIGURE S4 Prominent GFAP staining of astrocytes interacting with A β plaques brain of an AD patient with concurrent HSE. IF staining for A β (red), GFAP (white) and nuclei (Hoechst-33342; blue). Scale bar: 50 μ m

FIGURE S5 Microglia morphology and density in the brain of HSE patients and AD patient with HSE. Brain tissue sections were IF stained for A β (red), Iba1 (green) and nuclei (Hoechst-33342; blue). Boxes indicate areas shown at higher magnification. Scale bar: 50 μ m

How to cite this article: Tran DN, Bakx ATCM, van Dis V, Aronica E, Verdijk RM, Ouwendijk WJD. No evidence of aberrant amyloid β and phosphorylated tau expression in herpes simplex virus-infected neurons of the trigeminal ganglia and brain. *Brain Pathol*. 2021;00:e13044. <https://doi.org/10.1111/bpa.13044>