

EUR Research Information Portal

European Association of Urology Guidelines on the Management of Female Non-neurogenic Lower Urinary Tract Symptoms. Part 2

Published in:
European Urology

Publication status and date:
Published: 01/07/2022

DOI (link to publisher):
[10.1016/j.eururo.2022.01.044](https://doi.org/10.1016/j.eururo.2022.01.044)

Document Version
Publisher's PDF, also known as Version of record

Document License/Available under:
CC BY

Citation for the published version (APA):
Arlandis, S., Bø, K., Cobussen-Boekhorst, H., Costantini, E., de Heide, M., Farag, F., Groen, J., Karavitakis, M., Lapitan, M. C., Manso, M., Arteaga, S. M., Nambiar, A. K., Riogh, A. N. A., O'Connor, E., Omar, M. I., Peyronnet, B., Phé, V., Sakalis, V. I., Sihra, N., ... Harding, C. K. (2022). European Association of Urology Guidelines on the Management of Female Non-neurogenic Lower Urinary Tract Symptoms. Part 2: Underactive Bladder, Bladder Outlet Obstruction, and Nocturia. *European Urology*, 82(1), 60-70. <https://doi.org/10.1016/j.eururo.2022.01.044>

[Link to publication on the EUR Research Information Portal](#)

Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.

In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: openaccess.library@eur.nl. Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

Platinum Priority – Review – Female Urology – Incontinence
Editorial by Jean-Nicolas Cornu on pp. 71–72 of this issue

European Association of Urology Guidelines on the Management of Female Non-neurogenic Lower Urinary Tract Symptoms. Part 2: Underactive Bladder, Bladder Outlet Obstruction, and Nocturia

Salvador Arlandis^{a,*}, Kari Bø^b, Hanny Cobussen-Boekhorst^c, Elisabetta Costantini^d,
Monica de Heide^e, Fawzy Farag^{f,g}, Jan Groen^h, Markos Karavitakisⁱ, Marie Carmela Lapitan^j,
Margarida Manso^k, Serenella Monagas Arteaga^l, Arjun K. Nambiar^m, Aisling Nic An Ríoghⁿ,
Eabhann O'Connor^o, Muhammad Imran Omar^p, Benoit Peyronnet^{q,r}, Veronique Phé^s,
Vasileios I. Sakalis^t, Néha Sihra^u, Lazaros Tzelves^v, Mary-Lynne van Poelgeest-Pomfret^w,
Tine W.L. van den Bos^e, Huub van der Vaart^x, Christopher K. Harding^{m,y}

^a Urology Department, La Fe University and Polytechnic Hospital, Valencia, Spain; ^b Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway; ^c Department of Urology, Radboud UMC, Nijmegen, The Netherlands; ^d Department of Medicine and Surgery, University of Perugia, Perugia, Italy; ^e Bekkenbodem4All, Tiel, The Netherlands; ^f Department of Urology, Sohag University Hospital, Sohag, Egypt; ^g Department of Urology, East Suffolk and North Essex NHS Foundation Trust, Ipswich, UK; ^h Department of Urology, Erasmus Medical Center, Rotterdam, The Netherlands; ⁱ Department of Urology, Charing Cross Hospital, London, UK; ^j College of Medicine, Philippine General Hospital, National Institutes of Health, University of the Philippines Manila, Manila, Philippines; ^k Department of Urology, Centro Hospitalar Universitário São João, Porto, Portugal; ^l Department of Urology, University Hospital of San Agustín, Avilés, Spain; ^m Department of Urology, Freeman Hospital, Newcastle-upon-Tyne, UK; ⁿ Department of Urology, Cork University Hospital, Cork, Ireland; ^o Department of Urology, Beaumont Hospital, Dublin, Ireland; ^p University of Aberdeen, Aberdeen, UK; ^q European Association of Urology, Arnhem, The Netherlands; ^r Department of Urology, University of Rennes, Rennes, France; ^s Department of Urology, AP-HP, Pitié-Salpêtrière Academic Hospital, Sorbonne University, Paris, France; ^t Department of Urology, Agios Pavlos General Hospital of Thessaloniki, Thessaloniki, Greece; ^u Department of Urology, Guy's and St. Thomas' NHS Foundation Trust, London, UK; ^v Second Department of Urology, National and Kapodistrian University of Athens, Sismanogleio General Hospital, Athens, Greece; ^w World Federation of Incontinence and Pelvic Problems, Leiden, The Netherlands; ^x Department of Obstetrics and Gynecology, University Medical Center, Utrecht, The Netherlands; ^y Translational and Clinical Research Institute, Newcastle University, Newcastle-upon-Tyne, UK

Article info

Article history:

Accepted January 26, 2022

Associate Editor:

James Catto

Keywords:

Female lower urinary tracts symptoms
Underactive bladder
Bladder outlet obstruction

Abstract

Context: Female lower urinary tract symptoms (LUTS) are a common presentation in urological practice. Thus far, only a limited number of female LUTS conditions have been included in the European Association of Urology (EAU) guidelines compendium. The new non-neurogenic female LUTS guidelines expand the remit to include these symptoms and conditions.

Objective: To summarise the management of underactive bladder (UAB), bladder outlet obstruction (BOO), and nocturia in females.

Evidence acquisition: The literature search was updated in September 2021 and evidence synthesis was conducted using modified GRADE approach as outlined for all EAU guidelines. A new systematic review on BOO was carried out by the panel for purposes of this guideline.

* Corresponding author. Department of Urology, La Fe University and Polytechnic Hospital, Avenida Fernando Abril Martorell 106, 46026 Valencia, Spain.
E-mail address: salvador.arlandis@gmail.com (S. Arlandis).

Nocturia
 Conservative management
 European Association of Urology
 guidelines

Evidence synthesis: The important considerations for informing guideline recommendations are presented, along with a summary of all the guideline recommendations.

Conclusions: Non-neurogenic female LUTS are an important presentation of urological dysfunction. Initial evaluation, diagnosis, and management should be carried out in a structured and logical fashion on the basis of the best available evidence. This guideline serves to present this evidence to practising urologists and other health care providers in an easily accessible and digestible format.

Patient summary: This report summarises the main recommendations from the European Association of Urology guideline on symptoms and diseases of the female lower urinary tract (bladder and urethra) not associated with neurological disease. We cover recommendations related to the treatment of underactive bladder, obstruction of the bladder outlet, and nighttime urination.

© 2022 European Association of Urology. Published by Elsevier B.V. All rights reserved.

1. Introduction

Part 2 of the European Association of Urology (EAU) guideline summary on non-neurogenic female lower urinary tract symptoms (LUTS) presented here focuses on the sections relating to underactive bladder (UAB), bladder outlet obstruction (BOO), and nocturia. This summary relates primarily to the patient pathway from presentation through diagnostics and to management of the specific conditions. The best available evidence is summarised and the main recommendations from the full version of the guidelines are presented in a concise and easily digestible format.

2. Evidence acquisition

The scope of the EAU guidelines on non-neurogenic female LUTS was expanded, so a new literature search was carried out, with expansion of the terminology and criteria. The full details of the search strategy are available on the EAU website (<https://uroweb.org/wp-content/uploads/2021-EAU-Non-neurogenic-Female-LUTS-Guidelines-Search-Strategy.pdf>).

The EAU Guidelines Office uses a modified GRADE approach for evaluating the relevant literature on each topic area. High-quality systematic reviews (SRs) are referenced when available, and lower-quality evidence is evaluated if SRs are not available. For this edition of the guideline, new SRs on overactive bladder (OAB) and female bladder outlet obstruction (BOO) were conducted by the panel.

Evidence summary statements and assessments of the quality of the evidence available are reinforced by certainty ratings (ranging from very low to high). Recommendations are then produced in accordance with these certainty ratings, the benefit/harm balance, and consideration of patient values and preferences, where feasible, to give an overall recommendation with a strength rating of “strong” or “weak”. It should be noted that the balance between “strong” and “weak” recommendations is related to these three factors rather than just the evidence base for the intervention. Our panel recommendations are reinforced by the inclusion of patient representatives in the panel to provide a valuable input into discussions regarding patient values and preferences.

3. Evidence synthesis

3.1. Underactive bladder

UAB is a common condition, defined by the International Continence Society (ICS) as “a symptom complex characterised by a slow urinary stream, hesitancy, and straining to void, with or without a feeling of incomplete bladder emptying sometimes with storage symptoms” [1].

Detrusor underactivity (DU) is a diagnosis based on urodynamic studies and defined by the ICS as “a detrusor contraction of reduced strength and/or duration, resulting in prolonged bladder emptying and/or failure to achieve complete bladder emptying within a normal time span” [2].

3.1.1. Diagnostic evaluation

3.1.1.1. Symptoms associated with DU. According to current data, a pivotal symptom or collection of symptoms to specifically identify DU patients has not been identified. The ICI Questionnaire-Underactive Bladder (ICIQ-UAB) is a research tool that needs further validation before use as a patient-reported outcome measure in routine clinical practice [3].

3.1.1.2. Urodynamic studies. Noninvasive studies such as uroflowmetry, postvoid residual (PVR) volume measurement, and bladder voiding efficiency determination are potentially useful in identifying women who might have DU. There is considerable symptomatic overlap with BOO, and uroflowmetry and PVR findings may also be similar. Only invasive urodynamics with pressure-flow studies can reliably distinguish DU from BOO and these urodynamic diagnoses can co-exist. In addition, diagnosis in women is particularly difficult as females can void by relaxing the pelvic floor, that is, without a detectable detrusor contraction during the pressure-flow study and without an increase in abdominal pressure [4]. The simplest methods for defining and diagnosing DU involve the use of cutoff values for the maximum flow rate (Q_{max}) and the detrusor pressure at Q_{max} ($P_{detQ_{max}}$). There is no consensus on which threshold values should be used [5] and consequently the prevalence of DU depends on the criteria used [6].

Several proposed measures of contractile strength exist. Watt’s factor estimates the power generated by the detrusor per unit area of bladder [7]. Projected isovolumetric

pressure (PIP) is a gross simplification of the bladder output relation and estimates the maximum detrusor pressure that can be generated by the bladder when the outlet is closed, the isovolumetric detrusor pressure. The bladder contractility index is simply a reduction of PIP to an index [8]. PIP also estimates the isovolumetric detrusor pressure, but was developed in an entirely female population via an experimental method [9]. These parameters do not necessarily reflect what the detrusor might potentially achieve under optimum conditions [10].

3.1.2. Disease management

Treatment of female DU includes strategies to ensure bladder drainage, increase bladder contraction, decrease urethral resistance, or a combination [11]. The management goals for UAB are to improve symptoms and quality of life (QoL) and reduce the risk of complications.

3.1.2.1. Conservative management.

3.1.2.1.1. Behavioural interventions. Regular or timed voiding should be encouraged in women with impaired bladder sensations. Assisted voiding via abdominal straining with adequate relaxation of the pelvic floor muscle (PFM) has been recommended, as well as double or triple voiding, in an attempt to improve bladder emptying. None of these manoeuvres have proven efficacious in any randomised study. There is a possible association between voiding via excessive abdominal straining and the risk of pelvic organ prolapse (POP) or rectal prolapse [12]. A small retrospective study in neurogenic patients showed that Valsalva voiding may increase the risk of rectal prolapse when compared to clean intermittent self-catheterisation (CISC) [13].

3.1.2.1.2. PFM relaxation training with biofeedback. There are no randomised controlled trials (RCTs) examining PFM relaxation training in adult women with UAB. One study found significant relaxation of the PFM after PFM contraction [14] and another study found that PFM relaxation training over time increased the speed of relaxation after a single contraction [15]. In the absence of RCT data for women, the findings of an RCT for children with non-neuropathic UAB and voiding dysfunction comparing the effect of PFM relaxation and biofeedback plus combined treatment (hydration, scheduled voiding, toilet training, and diet) versus combined treatment alone can be cautiously extrapolated to an adult population [16]. The paediatric trial showed that additional PFM relaxation led to significant increases in the mean number of voiding episodes and Q_{max} and decreases in PVR volume and voiding time [16].

3.1.2.1.3. Clean intermittent self-catheterisation. CISC has proven efficacy in patients who are unable to empty their bladder and remains a gold standard for reducing the adverse consequences of a high PVR and incomplete voiding, despite the low level of evidence supporting this approach.

3.1.2.1.4. Indwelling catheter. An indwelling urinary catheter may be an option for some women for whom all other treatments have failed and who are unable to perform CISC. Complications include urinary tract infection (UTI),

stone formation, and urethral damage. Suprapubic catheterisation may be preferable over urethral catheterisation to minimise the risk of urethral trauma and pain [17].

3.1.2.1.5. Intravesical electrical stimulation. According to a retrospective study [18], intravesical electrical stimulation may be useful in some patients after prolonged bladder overdistension. However, this must be investigated in high-quality RCTs.

3.1.2.2. Pharmacological management.

3.1.2.2.1. Parasympathomimetics. An SR on the use of parasympathomimetics in patients with UAB included ten RCTs [19]. The SR did not support the use of parasympathomimetics for treating UAB, especially when frequent and/or serious adverse effects are taken into account.

3.1.2.2.2. α -Adrenergic Blockers. There is limited evidence regarding the effectiveness of α -blockers. One prospective study showed similar improvements in uroflowmetry parameters (specifically in the percentage of patients who had a good therapeutic response) with tamsulosin in women with BOO (39.4%) or DU (32.7%) [20]. Another longitudinal study including 14 women with DU showed clinical and urodynamic improvements after tamsulosin [21]. A prospective single-blind RCT in female patients with DU compared the efficacy of α -blockers, cholinergic drugs, and combination therapy, with the latter exhibiting the best results [22].

3.1.2.2.3. Prostaglandins. Prostaglandins E2 and F2 have been used intravesically to treat urinary retention after surgery. A Cochrane SR showed a statistically significant association between intravesically administered prostaglandin and successful voiding among postoperative patients with urinary retention. However, the success rate was low (32%) compared to placebo, with very low certainty of evidence [23].

3.1.2.3. Surgical management.

3.1.2.3.1. Sacral nerve stimulation. An RCT included 37 patients in the implantation arm and 31 in the standard medical therapy arm, showing a mean decrease in PVR volume in the implantation group [24]. A meta-analysis of seven studies (one RCT and six observational studies) showed a mean difference in PVR volume reduction of 236 ml and a mean voided volume increase of 299 ml [25]. The response rate during the trial phase ranged from 33% to 90% (mean 54.2%) and the success rate for permanent implantation ranged from 55% to 100% (mean 73.9%), highlighting that patient selection is crucial [26]. A subgroup of women with idiopathic urinary retention (Fowler's syndrome) had a higher response rate of 68–77% [27]. Sacral nerve stimulation (SNS) is a valid option for female patients with DU, with proper patient selection. Patients with evidence of anatomical BOO, suspected loss of intrinsic detrusor contractility, or neurogenic bladder dysfunction showed lower response rates [28].

3.1.2.3.2. OnabotulinumtoxinA. There is low-level evidence that onabotulinumtoxinA injections to the external striated urethral sphincter may improve voiding in patients with DU by reducing outlet resistance and suppressing the guarding reflex. Retrospective case studies have shown improvements in voiding symptoms, recovery of sponta-

neous voiding, and improvements in urodynamic parameters [29,30]. The duration of symptomatic relief is typically 3 mo.

3.1.2.3.3. Transurethral incision of the bladder neck. Transurethral incision of the bladder neck has been described in short series of women with refractory DU. In a retrospective case study, 40/82 women (48.8%) achieved satisfactory outcomes (spontaneous voiding with voiding efficiency $\geq 50\%$), but five (6.1%) of the patients developed stress urinary incontinence (SUI) and two (2.4%) developed a vesicovaginal fistula (VVF) [31].

3.1.2.3.4. Other procedures. Reduction cystoplasty and myoplasty are uncommon procedures with very limited evidence regarding their effectiveness.

Recommendations for the management of UAB are provided in Table 1.

3.2. Bladder outlet obstruction

BOO is defined by the ICS as “obstruction during voiding, characterised by increased detrusor pressure and reduced urine flow rate” [2].

3.2.1. Diagnostic evaluation

3.2.1.1. Clinical history. Evidence regarding the clinical utility of symptoms in the diagnosis of BOO is inconclusive. In a single-centre retrospective study including women with BOO, the authors concluded that symptom assessment alone was insufficient for diagnosis and a full urodynamic evaluation was essential [32]. Studies have found that significant proportions of women presenting with symptoms of urinary incontinence (UI) also have concomitant voiding symptoms and BOO on urodynamics [33,34].

3.2.1.2. Clinical examination. There are no studies evaluating the clinical utility of physical examination in women with suspected BOO; nevertheless, this is universally considered a key part of the medical assessment.

3.2.1.3. Uroflowmetry and PVR volume. Studies have shown reasonable correlation between low flow rates, significant PVR volume, and urodynamic BOO [34–37].

3.2.1.4. Ultrasound. The major utility of ultrasound scanning in women with BOO is in detecting possible complications such as bladder wall thickening or upper tract dilatation/hydronephrosis. One study reported that transvaginal ultrasonography was able to demonstrate a closed bladder neck during attempts at micturition and concluded that this modality was useful for the evaluation of possible causal factors in female BOO [38].

3.2.1.5. Magnetic resonance imaging. There are no reports on the clinical utility of magnetic resonance imaging (MRI) in the diagnosis of female BOO. MRI in patients with a urethral stricture can reveal the degree of periurethral fibrosis, although the prognostic and clinical significance of such a finding has not been established [39].

3.2.1.6. Electromyography. Abnormal electromyography (EMG) activity may be associated with nonrelaxation of the striated sphincter, abnormally high urethral pressure, poor bladder sensation, and reduced detrusor contractile strength [40,41]. Complex repetitive discharges and decelerating bursts are specific EMG abnormalities (using periurethral concentric needles) that have been described for patients with high-tone nonrelaxing sphincter, although these abnormalities also occurred in asymptomatic volunteers [42,43]. A review of voiding dysfunction in women showed that increased EMG activity of the PFM using surface electrodes during voiding or nonrelaxation, coupled with pressure-flow information from urodynamics, may be useful in differentiating between functional and anatomical obstruction [44].

3.2.1.7. Cystourethroscopy. Cystourethroscopy can be useful for visualising anatomical/mechanical obstruction and providing information regarding its nature, location, and calibre. Given that pelvic malignancy may cause anatomical BOO, cystourethroscopy is considered an essential part of the diagnostic pathway.

3.2.1.8. Urodynamics and video-urodynamics. Pressure-flow studies are the mainstay of BOO diagnosis and the characteristic abnormalities are a combination of low flow and high voiding pressure [45]. The urodynamic definition of female BOO remains controversial [46]. The Blaivas-Groutz nomogram is one of the most popular urodynamic criteria for female BOO [47] but it has been suggested that it overestimates obstruction [48]. The addition of fluoroscopic imaging introduces a video-urodynamic criterion for obstruction [49]. However, both methods lack data supporting their clinical validity, especially regarding their predictive value for treatment outcomes [50].

Table 1 – Recommendations for underactive bladder

Recommendation	Strength rating
Encourage double voiding in women who are unable to completely empty their bladder.	Weak
Warn women with UAB who use abdominal straining to improve emptying about the risk of pelvic organ prolapse.	Weak
Use CISC as a standard treatment in patients who are unable to empty their bladder.	Strong
Thoroughly instruct patients in the technique and risks of CISC.	Strong
Offer indwelling transurethral catheterisation and suprapubic cystostomy only when other modalities for urinary drainage have failed or are unsuitable.	Weak
Do not routinely recommend intravesical electrical stimulation in women with UAB.	Weak
Do not routinely recommend parasympathomimetics in the treatment of women with UAB.	Strong
Offer α -blockers before more invasive techniques.	Weak
Offer intravesical prostaglandins to women with urinary retention after surgery only in the context of well-regulated clinical trials.	Weak
Offer onabotulinumtoxinA external sphincter injections before more invasive techniques as long as the patient is informed that the evidence to support this treatment is of low quality.	Weak
Offer sacral nerve stimulation to women with UAB refractory to conservative measures.	Strong
Do not routinely offer detrusor myoplasty as a treatment for detrusor underactivity.	Weak

CISC = clean intermittent self-catheterisation; UAB = underactive bladder.

Several urodynamic cutoff values have been proposed to optimise the diagnostic accuracy of video-urodynamic studies [36]:

- $P_{\text{det}Q_{\text{max}}} \geq 30$ cm H₂O for differentiating BOO from bladder dysfunction and normal studies (area under the receiver operating characteristic curve [AUC] 0.78);
- Abrams-Griffiths number >30 for differentiating anatomical from functional BOO (AUC 0.66); and
- $P_{\text{det}Q_{\text{max}}} \geq 30$ cm H₂O for differentiating dysfunctional voiding from poor sphincter relaxation (AUC 0.93).

More recently, Solomon et al [51] devised a nomogram for calculation of the female BOO index (BOOIf) using the formula $\text{BOOIf} = P_{\text{det}Q_{\text{max}}} - 2.2Q_{\text{max}}$:

- BOOIf <0: <10% probability of obstruction;
- BOOIf 5–18: equivocal, $\geq 50\%$ likelihood of obstruction; and
- BOOIf >18: >90% likelihood of obstruction,

Voiding cystourethrography alone or in conjunction with concomitant pressure-flow studies may be useful in delineating the site of the obstruction [49].

3.2.2. Disease management

Therapeutic interventions for BOO aim to decrease outlet resistance and increase urinary flow, improve bladder emptying, and reduce LUTS [46,50,52]. Treatment choice is dictated by the nature of the underlying cause of the obstruction.

3.2.2.1. Conservative management.

3.2.2.1.1. Behavioural modification. Behavioural modification aims to improve or correct maladaptive voiding. It can include elements such as education regarding normal voiding function, self-monitoring of symptoms, changes in lifestyle factors, avoidance of constipation, and alteration of voiding technique. Ultimately, techniques aim to improve the coordination and synergistic action between the detrusor and sphincter [46,50,52]. General interventions such as those listed above may help with symptoms resulting from BOO, but no quantification of their effect is possible from existing published data.

3.2.2.1.2. PFM training \pm biofeedback. PFM relaxation training with biofeedback may result in relaxation of the PFM/urethral sphincter in women with dysfunctional voiding. A case series involving women with pelvic muscle or external urethral sphincter hyperactivity during voiding showed improved relaxation and voiding function following PFM training (PFMT) with biofeedback [53]. High-quality RCTs are needed to confirm such observations.

3.2.2.1.3. Vaginal pessaries. In a prospective study of 18 women with grade 3 or 4 cystoceles and urodynamic BOO, normal voiding was noted in 17 (94%) following placement of a vaginal pessary [54].

3.2.2.1.4. Urinary catheterisation. In a series of 20 patients with voiding dysfunction after tension-free vaginal tape surgery who adopted a CISC programme, 59% had a consistent residual volume <100 ml and 50% were voiding normally within 12 wk [55].

3.2.2.1.5. Intraurethral inserts. In a study among women with voiding dysfunction who received an intraurethral

insert, device removal within 7 d of insertion occurred in 60% of cases because of discomfort, pericatheter leakage, or technical difficulty. The 20% who continued to use the device in the long term were satisfied, with PVR volumes remaining <100 ml. Adverse events included device migration and symptomatic UTI [56,57]. There is no convincing evidence from RCTs to support the use of intraurethral inserts.

3.2.2.1.6. Extracorporeal magnetic stimulation. In a small prospective nonrandomised trial, alfuzosin was compared to electromagnetic stimulation and to the combination of both in women with functional BOO. Significant increases in Q_{max} and decreases in symptoms were observed in all groups, with greater improvements in the combination therapy group [58].

3.2.2.2. Pharmacological management.

3.2.2.2.1. α -Adrenergic blockers. In the only placebo-controlled RCT reporting subgroup analyses among women with urodynamically proven BOO, no significant difference was observed in symptoms, Q_{max} , or PVR after 8 wk of alfuzosin versus placebo [59]. A small nonrandomised trial compared the use of tamsulosin and prazosin. More patients treated with tamsulosin experienced a decrease in symptoms and treatment satisfaction. More adverse events were reported with prazosin [60].

3.2.2.2.2. Striated muscle relaxants. A randomised placebo-controlled crossover trial investigated oral baclofen in 60 women diagnosed with BOO. The results showed a lower number of voids and improvements in Q_{max} and $P_{\text{det}Q_{\text{max}}}$ with 4 wk of baclofen in comparison to placebo [61].

3.2.2.2.3. Sildenafil. A placebo-controlled, randomised crossover trial in women with BOO showed that sildenafil is not superior to placebo in improving symptoms or urodynamic parameters of female BOO [62].

3.2.2.2.4. Thyrotropin-releasing hormone. A small RCT including women with voiding problems of mixed aetiologies showed no difference in urodynamic outcomes between intravenous thyrotropin-releasing hormone and placebo [63].

3.2.2.3. Surgical management.

3.2.2.3.1. Intrasphincter botulinum toxin injection. A SR in women with dysfunctional voiding showed improvements in symptoms and reductions in residual volume as well as voiding pressure. Larger series in adults describe success rates of 86–100% [64]. In a randomised study, 100 U of onabotulinumtoxinA resulted in a significantly lower International Prostate Symptom Score (IPSS) and larger voided volume in adults with voiding dysfunction [65]. Two small case series of women with BOO who received an intrasphincter injection of onabotulinumtoxinA (100 U) showed improvements in symptoms, a significant reduction in PVR, an increase in Q_{max} , and an improvement in static urethral pressure profile [40,66]. The average symptom-free duration was 16.8 wk [66]. Adverse events included UTI and a temporary need for CISC. No SUI was reported.

3.2.2.3.2. Sacral nerve stimulation. A cohort study of women who underwent SNS for urinary retention associated with outlet obstruction showed an overall spontaneous voiding rate of 72% over mean follow-up of 4 yr [67]. In a

single-centre series of patients with idiopathic urinary retention who underwent SNS, 62.5% achieved a >50% reduction in CISC rate [68].

3.2.2.3.3. POP surgery. A multicentre prospective study involving women with grade ≥ 2 symptomatic POP who underwent surgery demonstrated a significant reduction in voiding symptoms and PVR volume at 1 yr after surgery [69]. A retrospective study of women who underwent laparoscopic sacrocolpopexy for POP showed a significant increase in mean postoperative Q_{\max} and decreases in P_{\det} - Q_{\max} and PVR volume in those aged ≥ 65 yr [70].

3.2.2.3.4. Urethral dilatation. Pooled analysis of data from an SR of retrospective studies of females with urethral stricture showed a mean success rate of 49% after urethral dilatation to 41 Fr at mean follow-up of 46 mo. The mean time to failure was 12 mo. Among treatment-naïve patients, the success rate was 58%, compared to 27.2% among patients who had undergone previous dilatation [39]. Significantly greater improvements in Q_{\max} and PVR were seen with intermittent urethral dilatation compared to on-demand dilatation for primary urethral stricture [71]. Worsening or new-onset SUI, frequency, and urgency after dilatation have been reported [72].

3.2.2.3.5. Urethrotomy. A prospective study of women with urethral strictures who underwent Otis urethrotomy to 40 Fr followed by 6-weekly dilatations demonstrated improvement in IPSS, QoL, voided volume, Q_{\max} , and PVR volume at 6 mo. Only the improvements in PVR volume and QoL were maintained on long-term follow-up [73].

3.2.2.3.6. Bladder neck incision or resection. A review of case studies on bladder neck incision for the treatment of bladder neck obstruction in women reported success rates of 76–100% [45]. Several prospective case series consistently reported significant improvements in IPSS, QoL, Q_{\max} , P_{\det} - Q_{\max} , and PVR after treatment, regardless of the site of the incision, type of energy used, or length of follow-up [74–77]. Complications reported included VVF (3.6%), SUI (4.7%), and urethral stricture (3.6%). Complications of VVF and SUI were noted in the cohort of patients who had their incisions at the 5- and 7-o'clock positions, and not in those who had their incisions at the 2- and 10-o'clock positions [77].

Bladder neck incision and V-Y reconstruction using Nesbit's technique in women with BOO showed similar rates of improvement in symptoms and postoperative PVR volume. V-Y plasty had longer operating and catheter times, a lower improvement rate, a higher transfusion rate, and a higher adverse event rate [78].

3.2.2.3.7. Urethroplasty or urethral reconstruction. Retrospective studies reporting outcomes for urethroplasty detail success rates of 57–100% [39,79]. Pooled analysis from studies using vaginal or labial flaps showed a mean success rate of 91% over mean follow-up of 32 mo. Vaginal or labial graft urethroplasty had a success rate of 80% at mean follow-up of 22 mo. Oral mucosal grafts had a mean success of 94% after mean follow-up of 15 mo [39]. A later review of retrospective studies on dorsal buccal mucosal grafts reported success rates of 62–100%, with a pooled success rate of 86% [80]. A long-term study with mean follow-up of 32 mo showed a stricture recurrence rate of 23.1% [79].

A retrospective study comparing women who underwent urethral dilatation or urethroplasty with a dorsal-onlay pedicled labium flap reported significant improvements in both groups. The urethroplasty group had significantly better QoL scores and Q_{\max} at follow-up in comparison to the dilatation group [81]. Adverse events associated with urethroplasty include new-onset SUI and urgency and worsening of urge UI.

3.2.2.3.8. Urethrolisis. Case series show improved voiding and lower PVR volumes, improvement or resolution of symptoms and QoL, and improvement of urodynamic parameters after urethrolisis treatment [82–84]. De novo SUI was reported in 39% of cases in one study [84]. A greater delay in performing urethrolisis was associated with persistent bladder symptoms [85].

3.2.2.3.9. Removal, excision, section, or loosening of mid-urethral slings. Several small retrospective reviews of cases using different techniques for sling revision (incision, partial excision, or excision) showed good success rates in terms of symptom reduction, resumption of voiding with a significant reduction in PVR volume, and improvement of urodynamic parameters. SUI recurs in a small proportion of patients and often to a lesser degree than before the sling procedure. Studies have shown long-term efficacy, including preservation of continence.

No significant difference in success rates was demonstrated on comparison of different techniques. There was a greater need for surgery for recurrent SUI after partial sling excision in the group without an anti-SUI procedure [86].

One study showed that patients who underwent surgical release >180 d after initial anti-SUI surgery had significantly less recurrent SUI in comparison to patients who underwent the release sooner [87].

Recommendations for the management of female BOO are provided in Table 2.

3.3. Nocturia

Nocturia was defined by the ICS in 2002 as “the complaint that the individual has to wake at night one or more times to void” and quantified in an updated document in 2019 as “the number of times an individual passes urine during their main sleep period, from the time they have fallen asleep up to the intention to rise from that period” [88].

3.3.1. Diagnostic evaluation

Evaluation of nocturia should include a thorough medical history and physical examination, with particular reference to history of sleep disorders, fluid balance, associated LUTS, cardiovascular and endocrine comorbidity, renal disease, current medications, and history of urological disease [89].

A bladder diary is a vital initial investigation in patients complaining of nocturia. A low nocturnal bladder capacity or global bladder capacity will be highlighted by lower voided volumes. Global polyuria is defined as 24-h urine production >40 ml/kg [90] and may be present in conditions such as diabetes mellitus and diabetes insipidus. The definition of nocturnal polyuria is age-dependent and the thresholds for this diagnosis range from 20% (in younger individuals) to 33% (age >65 yr) of the 24-h urine volume

Table 2 – Recommendations for female BOO

Recommendation	Strength rating
Diagnosis	
Take a full clinical history and perform a thorough clinical examination in women with suspected BOO.	Strong
Do not rely on measurements from urine flow studies alone to diagnose female BOO.	Strong
Perform cystourethroscopy in women with suspected anatomical BOO.	Strong
Perform urodynamic evaluation in women with suspected BOO.	Strong
Conservative treatment	
Offer PFMT aimed at PFM relaxation to women with functional BOO.	Weak
Prioritise research that investigates and advances understanding of the mechanisms and impact of PFMT on the coordinated relaxation of the pelvic floor during voiding.	Strong
Offer the use of a vaginal pessary to women with grade 3 or 4 cystoeces and BOO who are not eligible/inclined towards other treatment options.	Weak
Offer urinary containment devices to women with BOO to address urinary leakage as a result of BOO, but not as a treatment to correct the condition.	Weak
Offer CISC to women with urethral strictures or post-UI surgery for BOO.	Weak
Do not offer an intraurethral device to women with BOO.	Strong
Pharmacological treatment	
Offer uroselective α -blockers as an off-label option to women with functional BOO following discussion of the potential benefits and adverse events.	Weak
Offer oral baclofen to women with BOO, particularly those with increased EMG activity and sustained detrusor contraction during voiding.	Weak
Only offer sildenafil to women with BOO as part of a well-regulated clinical trial.	Strong
Do not offer thyrotropin-releasing hormone to women with BOO.	Strong
Surgical treatment	
Offer intrasphincter injection of botulinum toxin to women with functional BOO.	Weak
Offer sacral neuromodulation to women with functional BOO.	Weak
Advise women with voiding symptoms associated with POP that symptoms may improve after surgery.	Weak
Offer urethral dilatation to women with urethral stenosis causing BOO, but advise on the likely need for repeated intervention.	Weak
Offer internal urethrotomy with postoperative urethral self-dilatation to women with BOO due to urethral stricture disease but advise on its limited long-term improvement and the risk of postoperative UI.	Weak
Do not offer urethral dilatation or urethrotomy as a treatment for BOO to women who have previously undergone mid-urethral synthetic tape insertion owing to the theoretical risk of causing urethral mesh extrusion.	Weak
Inform women of the limited long-term improvement (only in terms of PVR volume and QoL) after internal urethrotomy.	Weak
Offer bladder-neck incision to women with BOO secondary to primary bladder-neck obstruction.	Weak
Advise women who undergo bladder-neck incision on the small risk of developing SUI, VVF, or urethral stricture postoperatively.	Strong
Offer urethroplasty to women with BOO due to recurrent urethral stricture after failed primary treatment.	Weak
Caution women on the possible recurrence of strictures on long-term follow-up after urethroplasty.	Weak
Offer urethrolysis to women who have voiding difficulties after anti-UI surgery.	Weak
Offer sling revision (release, incision, partial excision, or excision) to women who develop urinary retention or significant voiding difficulty after tape surgery for UI.	Strong
Caution women about the risk of recurrent SUI and the need for repeat/concurrent anti-UI surgery after sling revision.	Strong
BOO = bladder outlet obstruction; CISC = clean intermittent self-catheterisation; EMG = electromyography; PFM = pelvic floor muscle; PFMT = PFM training; POP = pelvic organ prolapse; PVR = postvoid residual; QoL = quality of life; SUI = stress urinary incontinence; VVF= vesicovaginal fistula; UI = urinary incontinence.	

produced during sleep. A large study conducted across European and American centres involving ~2000 patients identified nocturnal polyuria as a contributory cause of nocturia in 89% of patients who were being treated for LUT abnormalities.

3.3.2. Disease management

3.3.2.1. *Conservative management.* Owing to the lack of high-quality evidence, most recommendations are derived from consensus methodology. Interventions that may help with nocturia include:

- Reduction of fluid intake at specific times;
- Avoidance/moderation of intake of caffeine or alcohol;
- Distraction techniques;
- Bladder retraining;
- PFMT;
- Review of medication; and
- Treatment of constipation.

In the EAU systematic review [91], three studies [92–94] were favourable for conservative treatment with PFMT, with another failing to confirm a benefit [95].

Individual and group PFMT approaches appear to be equally effective in reducing nocturia episodes [95]. Most studies evaluating PFMT for nocturia in women with additional urinary symptoms have shown positive results compared with placebo, transcutaneous electrical nerve stimulation (TENS), or anticholinergic drugs [92,93,95].

In patients with obstructive sleep apnoea who complain of nocturia, continuous positive airway pressure was shown to be effective in an SR and meta-analysis of five RCTs involving both sexes [96].

3.3.2.2. Pharmacological management.

3.3.2.2.1. *Desmopressin.* A recent SR [91] identified three trials of desmopressin specifically conducted in women. A dose-response relationship was observed. Significant changes in nocturnal urine volumes were reported in favour of higher desmopressin doses. Differences in the nocturnal polyuria index also tended to favour desmopressin over placebo. The level of certainty of the evidence from these trials is low. Desmopressin treatment for nocturia led to significant reductions in nocturnal urine output, nocturnal urinary frequency, and the nocturnal polyuria index [97–99]. Most

nocturia patients tolerate desmopressin treatment without clinically significant hyponatraemia; however, the risk of hyponatraemia increases with increasing age and lower baseline serum sodium concentration [91].

Desmopressin treatment in elderly patients should include careful monitoring of serum sodium concentrations and should be avoided in patients with a baseline serum sodium concentration below the normal range [100].

Desmopressin can be safely combined with anticholinergics with significant additional benefit in women with OAB and nocturnal polyuria, as shown by a multicentre RCT of 97 patients [101].

3.3.2.2.2. Anticholinergics. A SR [91] identified three RCTs involving anticholinergics such as oxybutynin [94] and tolterodine [95,101]. Treatment of nocturia in OAB patients with anticholinergic drugs led to a reduction in nocturia episodes.

3.3.2.2.3. Oestrogens. In a recent SR [91] only a single RCT investigating the efficacy of oestrogen for nocturia was identified [102]. Vaginal oestrogen may be beneficial in the treatment of nocturia in approximately 50% of women, but the certainty of evidence for this outcome was low.

3.3.2.2.4. Diuretic treatment. A randomised placebo-controlled study investigating afternoon (timed) diuretic treatment with furosemide showed a reduction in nocturia

episodes and nocturnal voided volume in men, but no similar studies have been conducted in women [103].

Recommendations for the management of nocturia are provided in Table 3.

4. Conclusions

Non-neurogenic female LUTS comprise a broad range of symptomatology and conditions, and diagnostic uncertainty is common. A thorough history and a stepwise logical approach to investigation are required to arrive at an accurate diagnosis. Management should be guided by individual patient factors and a collaborative approach with patients to guide treatment decisions.

Author contributions: Salvador Arlandis had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Arlandis, Nambiar, Bø, Cobussen-Boekhorst, Costantini, de Heide, Groen, Lapitan, Phé, van Poelgeest-Pomfret, van den Bos, van der Vaart, Harding.

Acquisition of data: Farag, Karavitakis, Manso, Monagas Arteaga, Nic An Riogh, O'Connor, Peyronnet, Sakalis, Sihra, Tselves.

Analysis and interpretation of data: Omar, Arlandis.

Drafting of the manuscript: Arlandis, Nambiar, Harding, Lapitan.

Critical revision of the manuscript for important intellectual content: Arlandis, Nambiar, Bø, Cobussen-Boekhorst, Costantini, de Heide, Groen, Lapitan, Phé, van Poelgeest-Pomfret, van den Bos, van der Vaart, Harding.

Statistical analysis: None.

Obtaining funding: None.

Administrative, technical, or material support: None.

Supervision: Arlandis, Harding, Lapitan.

Other: None.

Financial disclosures: Salvador Arlandis certifies that all conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript (eg, employment/affiliation, grants or funding, consultancies, honoraria, stock ownership or options, expert testimony, royalties, or patents filed, received, or pending), are the following: Salvador Arlandis has received company speaker honoraria from Astellas Pharma S.A., Bayer Hispania S.L., and Lacer-Holliste; has received honoraria for consultation from Presurgy S.L., Boston Scientific Europe, and Coloplast Productos Medicos; has participated in clinical trials by GlaxoSmithKline, Astellas, Ipsen, and Urovant; has received travel grants from Astellas Pharma S. A., WellSpect, Lacer, and Rovi; and has been a company consultant for Lacer Laboratories. Christopher K. Harding is a company consultant for Teleflex Medical; has received speaker honoraria from Astellas, Allergan, and Medtronic; has participated in clinical trials by Medtronic; has received fellowship and travel grants from Medtronic; and has received research support from the National Institute for Health Research and The Urological Foundation. Marie Carmela Lapitan has received speaker honoraria from Astellas, AstraZeneca, and GSK. Elisabetta Costantini has received research support from S&R Farmaceutici. Jan Groen has participated in clinical trials by Neuspera Medical Inc. Veronique Phé has received speaker honoraria from Medtronic and Viatrix and is a company consultant for Allergan and Coloplast. Huub van der Vaart has ownership interest in a patent with Ligalli B.V.; is a director of Berman Clinics; receives royalties from NVOG; is participating in clinical trials by UroGyn; and receives research support from ZonMw and NWO-TTW. Benoit Pey-

Table 3 – Recommendations for nocturia

Recommendation	Strength rating
Take a complete medical history from women with nocturia.	Strong
Use a validated questionnaire during the assessment of women with nocturia and for re-evaluation during and/or after treatment.	Weak
Use a 3-d bladder diary to assess nocturia in women.	Strong
Do not use nocturnal-only bladder diaries to evaluate nocturia in women.	Weak
Offer women with LUTS lifestyle advice before or concurrent with treatment.	Strong
Offer PFMT for nocturia (either individually or in the group setting) to women with UI or other storage LUTS.	Strong
Offer women with nocturia and a history suggestive of obstructive sleep apnoea a referral to a sleep clinic for an assessment of suitability for continuous positive airway pressure treatment.	Strong
Offer an anticholinergic treatment for nocturia to women with UUI or other storage LUTS following appropriate counselling regarding the potential benefits and associated risks.	Strong
Inform women with nocturia that the combination of behavioural therapy and anticholinergic drugs is unlikely to provide greater efficacy than either modality alone.	Weak
Offer a combination of anticholinergics and desmopressin to women with OAB and nocturia secondary to nocturnal polyuria following appropriate counselling regarding the potential benefits and associated risks.	Weak
Offer vaginal oestrogen treatment to women with nocturia following appropriate counselling regarding the potential benefits and associated risks.	Weak
Offer timed diuretic treatment to women with nocturia secondary to polyuria following appropriate counselling regarding the potential benefits and associated risks.	Weak

LUTS = lower urinary tract symptoms; OAB = overactive bladder; PFMT = pelvic floor muscle training; UUI = urge urinary incontinence.

ronnet is a company consultant for Boston Scientific. Lazaros Tzelves receives research support from the National and Kapodistrian University of Athens and the ASCAPE Project and is participating in a clinical trial by the ASCAPE Project. The remaining authors have nothing to disclose.

Funding/Support and role of the sponsor: None.

References

- Chapple CR, Osman NI, Birder L, et al. Terminology report from the International Continence Society (ICS) Working Group on Underactive Bladder (UAB). *Neurourol Urodyn* 2018;37:2928–31. <https://doi.org/10.1002/nau.23701>.
- Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn* 2002;21:167–78. <https://doi.org/10.1002/nau.10052>.
- Uren AD, Cotterill N, Harding C, et al. The development of the ICIQ-UAB: A patient reported outcome measure for underactive bladder. *Neurourol Urodyn* 2019;38:996–1004.
- Kira S, Mitsui T, Kobayashi H, Haneda Y, Sawada N, Takeda M. Detrusor pressures in urodynamic studies during voiding in women. *Int Urogynecol J* 2017;28(5):783–7. <https://doi.org/10.1007/s00192-016-3203-5>.
- Osman NI, Esperto F, Chapple CR. Detrusor underactivity and the underactive bladder: a systematic review of preclinical and clinical studies. *Eur Urol* 2018;74:633–43. <https://doi.org/10.1016/j.eururo.2018.07.037>.
- Jeong SJ, Lee JK, Kim KM, Kook H, Cho SY, Oh SJ. How do we diagnose detrusor underactivity? Comparison of diagnostic criteria based on an urodynamic measure. *Investig Clin Urol* 2017;58:247–54. <https://doi.org/10.4111/icu.2017.58.4.247>.
- Griffiths DJ, Constantinou CE, van Mastrigt R. Urinary bladder function and its control in healthy females. *Am J Physiol* 1986;251:R225–30. <https://doi.org/10.1152/ajpregu.1986.251.2.r225>.
- Abrams P. Bladder outlet obstruction index, bladder contractility index and bladder voiding efficiency: three simple indices to define bladder voiding function. *BJU Int* 1999;84:14–5.
- Tan TL, Bergmann Ma, Griffiths D, Resnick NM. Stop test or pressure-flow study? Measuring detrusor contractility in older females. *Neurourol Urodyn* 2004;23:184–9. <https://doi.org/10.1002/nau.20020>.
- Griffiths DJ. Assessment of detrusor contraction strength or contractility. *Neurourol Urodyn* 1991;10:1–18. <https://doi.org/10.1002/nau.1930100102>.
- Van Koeveeringe GA, Vahabi B, Andersson KE, Kirschner-Herrmans R, Oelke M. Detrusor underactivity: A plea for new approaches to a common bladder dysfunction. *Neurourol Urodyn* 2011;30:723–8. <https://doi.org/10.1002/nau.21097>.
- Salvatore S, Rademakers K, DeLancey J. Pathophysiology of urinary incontinence, faecal incontinence and pelvic organ prolapse. In: Abrams P, Cardozo L, Wagg A, Wein A, editors. *Incontinence*. Bristol, UK: Bristol Urological Institute; 2017. p. 361–495.
- El Akri M, Brochard C, Hascoet J, et al. Risk of prolapse and urinary complications in adult spina bifida patients with neurogenic acontractile detrusor using clean intermittent catheterization versus Valsalva voiding. *Neurourol Urodyn* 2019;38:269–77. <https://doi.org/10.1002/nau.23844>.
- Naess I, Bø K. Can maximal voluntary pelvic floor muscle contraction reduce vaginal resting pressure and resting EMG activity? *Int Urogynecol J* 2018;29:1623–7. <https://doi.org/10.1007/s00192-018-3599-1>.
- Mercier J, Morin M, Tang A, et al. Pelvic floor muscle training: mechanisms of action for the improvement of genitourinary syndrome of menopause. *Climacteric* 2020;23:468–73. <https://doi.org/10.1080/13697137.2020.1724942>.
- Ladi-Seyedian S, Kajbafzadeh A-M, Sharifi-Rad L, Shadgan B, Fan E. Management of non-neuropathic underactive bladder in children with voiding dysfunction by animated biofeedback: a randomized clinical trial. *Urology* 2015;85:205–10. <https://doi.org/10.1016/j.urology.2014.09.025>.
- Apostolidis A, Drake MJ, Emmanuel A. Neurological urinary and fecal incontinence. In: Abrams P, Cardozo L, Wagg A, Wein A, editors. *Incontinence*. Bristol, UK: Bristol Urological Institute; 2017. p. 1095–308.
- Huber ER, Kiss G, Berger T, Rehder P, Madersbacher H. The value of intravesical electrostimulation in the treatment of acute prolonged bladder overdistension. *Urologe A* 2007;46:662–6. <https://doi.org/10.1007/s00120-007-1312-y>.
- Barendrecht MM, Oelke M, Laguna MP, Michel MC. Is the use of parasympathomimetics for treating an underactive urinary bladder evidence-based? *BJU Int* 2007;99:749–52. <https://doi.org/10.1111/j.1464-410X.2006.06742.x>.
- Chang SJ, Chiang IN, Yu HJ. The effectiveness of tamsulosin in treating women with voiding difficulty. *Int J Urol* 2008;15:981–5.
- Costantini E, Lazzeri M, Bini V, et al. Open-label, longitudinal study of tamsulosin for functional bladder outlet obstruction in women. *Urol Int* 2009;83:311–5. <https://doi.org/10.1159/000241674>.
- Yamanishi T, Yasuda K, Kamai T, et al. Combination of a cholinergic drug and an α -blocker is more effective than monotherapy for the treatment of voiding difficulty in patients with underactive detrusor. *Int J Urol* 2004;11:88–96. <https://doi.org/10.1111/j.1442-2042.2004.00753.x>.
- Buckley BS, Lapitan MCM. Drugs for treatment of urinary retention after surgery in adults. *Cochrane Database Syst Rev* 2010;2010:CD008023. <https://doi.org/10.1002/14651858.cd008023.pub2>.
- Jonas U, Fowler CJ, Chancellor MB, et al. Efficacy of sacral nerve stimulation for urinary retention: results 18 months after implantation. *J Urol* 2001;165:15–9. <https://doi.org/10.1097/00005392-200101000-00004>.
- Gross C, Habli M, Lindsell C, South M. Sacral neuromodulation for nonobstructive urinary retention: a meta-analysis. *Female Pelvic Med Reconstr Surg* 2010;16:249–53. <https://doi.org/10.1097/spv.0b013e3181df9b3f>.
- Gani J, Hennessey D. The underactive bladder: diagnosis and surgical treatment options. *Transl Androl Urol* 2017;6(Suppl 2):S186–95. <https://doi.org/10.21037/tau.2017.04.07>.
- Swinn MJ, Kitchen ND, Goodwin RJ, Fowler CJ. Sacral neuromodulation for women with Fowler's syndrome. *Eur Urol* 2000;38:439–43. <https://doi.org/10.1159/000020321>.
- Panicker JN, Fowler CJ, Kessler TM. Lower urinary tract dysfunction in the neurological patient: clinical assessment and management. *Lancet Neurol* 2015;14:720–32. [https://doi.org/10.1016/s1474-4422\(15\)00070-8](https://doi.org/10.1016/s1474-4422(15)00070-8).
- Kuo HC. Recovery of detrusor function after urethral botulinum A toxin injection in patients with idiopathic low detrusor contractility and voiding dysfunction. *Urology* 2007;69:57–61. <https://doi.org/10.1016/j.urology.2006.08.1117>.
- Kuo HC. Effect of botulinum A toxin in the treatment of voiding dysfunction due to detrusor underactivity. *Urology* 2003;61:550–4. [https://doi.org/10.1016/s0090-4295\(02\)02541-4](https://doi.org/10.1016/s0090-4295(02)02541-4).
- Lee K-CC, Lee YK, Kuo H-C. Therapeutic efficacy and quality of life improvement in women with detrusor underactivity following transurethral incision of the bladder neck. *Urol Sci* 2019;30:266–71. https://doi.org/10.4103/UROS.UROS_39_19.
- Grouz A, Blaivas JG, Chaikin DC. Bladder outlet obstruction in women: definition and characteristics. *Neurourol Urodyn* 2000;19:213–20. [https://doi.org/10.1002/\(sici\)1520-6777\(2000\)19:3%3C213::aid-nau2%3E3.0.co;2-u](https://doi.org/10.1002/(sici)1520-6777(2000)19:3%3C213::aid-nau2%3E3.0.co;2-u).
- Massey JA, Abrams PH. Obstructed voiding in the female. *Br J Urol* 1988;61:36–9. <https://doi.org/10.1111/j.1464-410x.1988.tb09158.x>.
- Gravina GL, Costa AM, Galatioto GP, Ronchi P, Tubaro A, Vicentini C. Urodynamic obstruction in women with stress urinary incontinence—do nonintubated uroflowmetry and symptoms aid diagnosis? *J Urol* 2007;178:954–9.
- Brucker BM, Shah S, Mitchell S, et al. Comparison of urodynamic findings in women with anatomical versus functional bladder outlet obstruction. *Female Pelvic Med Reconstr Surg* 2013;19:46–50.
- Hsiao SM, Lin HH, Kuo HC. Videourodynamic studies of women with voiding dysfunction. *Sci Rep* 2017;7:6845. <https://doi.org/10.1038/s41598-017-07163-2>.
- Klijer R, Bar K, Biaūčūčlek W. Bladder outlet obstruction in women: difficulties in the diagnosis. *Urol Int* 2004;73:6–10.
- Galica V, Toska E, Saldutto P, Galatioto GP, Vicentini C. Use of transvaginal ultrasound in females with primary bladder neck obstruction. A preliminary study. *Arch Ital Urol Androl* 2015;87:158–60.

- [39] Osman NI, Mangera A, Chapple CR. A systematic review of surgical techniques used in the treatment of female urethral stricture. *Eur Urol* 2013;64:965–73. <https://doi.org/10.1016/j.eururo.2013.07.038>.
- [40] Panicker JN, Anding R, Arlandis S, et al. Do we understand voiding dysfunction in women? Current understanding and future perspectives: ICI-RS 2017. *Neurourol Urodyn* 2018;37(Suppl 4):S75–85. <https://doi.org/10.1002/nau.23709>.
- [41] Webb RJ, Fawcett PR, Neal DE. Electromyographic abnormalities in the urethral and anal sphincters of women with idiopathic retention of urine. *Br J Urol* 1992;70:22–5.
- [42] Tawadros C, Burnett K, Derbyshire LF, Tawadros T, Clarke NW, Betts CD. External urethral sphincter electromyography in asymptomatic women and the influence of the menstrual cycle. *BJU Int* 2015;116:423–31.
- [43] Fowler CJ, Christmas TJ, Chapple CR, Parkhouse HF, Kirby RS, Jacobs HS. Abnormal electromyographic activity of the urethral sphincter, voiding dysfunction, and polycystic ovaries: a new syndrome? *BMJ* 1988;297:1436–8.
- [44] Abdel Raheem A, Madersbacher H. Voiding dysfunction in women: how to manage it correctly. *Arab J Urol* 2013;11:319–30. <https://doi.org/10.1016/j.aju.2013.07.005>.
- [45] Nitti VW. Primary bladder neck obstruction in men and women. *Rev Urol* 2005;7(Suppl 8):S12–7.
- [46] Hoffman DS, Nitti VW. Female Bladder Outlet Obstruction. *Curr Urol Rep* 2016;17:31.
- [47] Blaivas JG, Groutz A. Bladder outlet obstruction nomogram for women with lower urinary tract symptomatology. *Neurourol Urodyn* 2000;19:553–64.
- [48] Akikwala TV, Fleischman N, Nitti VW. Comparison of diagnostic criteria for female bladder outlet obstruction. *J Urol* 2006;176:2093–7.
- [49] Nitti VW, Tu LM, Gitlin J. Diagnosing bladder outlet obstruction in women. *J Urol* 1999;161:1535–40.
- [50] Rademakers K, Apostolidis A, Constantinou C, et al. Recommendations for future development of contractility and obstruction nomograms for women. ICI-RS 2014. *Neurourol Urodyn* 2016;35:307–11.
- [51] Solomon E, Yasmin H, Duffy M, Rashid T, Akinluyi E, Greenwell TJ. Developing and validating a new nomogram for diagnosing bladder outlet obstruction in women. *Neurourol Urodyn* 2018;37:368–78.
- [52] Meier K, Padmanabhan P. Female bladder outlet obstruction: an update on diagnosis and management. *Curr Opin Urol* 2016;26:334–41.
- [53] Deindl FM, Vodusek DB, Bischoff C, Hofmann R, Hartung R. Dysfunctional voiding in women: which muscles are responsible? *Br J Urol* 1998;82:814–9.
- [54] Romanzi LJ, Chaikin DC, Blaivas JG. The effect of genital prolapse on voiding. *J Urol* 1999;161:581–6.
- [55] Bailey C, Matharu G. Conservative management as an initial approach for post-operative voiding dysfunction. *Eur J Obstet Gynecol Reprod Biol* 2012;160:106–9.
- [56] Madjar S, Sabo E, Halachmi S, et al. A remote controlled intraurethral insert for artificial voiding: a new concept for treating women with voiding dysfunction. *J Urol* 1999;161:895–8.
- [57] Madjar S, Halachmi S, Wald M, et al. Long-term follow-up of the In-Flow™ intraurethral insert for the treatment of women with voiding dysfunction. *Eur Urol* 2000;38:161–6. <https://doi.org/10.1159/000020274>.
- [58] Koh JS, Kim SJ, Kim HS, Kim JC. Comparison of alpha-blocker, extracorporeal magnetic stimulation alone and in combination in the management of female bladder outlet obstruction. *Int Urogynecol J* 2011;22:849–54.
- [59] Lee Y-S, Lee K-S, Choo M-S, et al. Efficacy of an alpha-blocker for the treatment of nonneurogenic voiding dysfunction in women: an 8-week, randomized, double-blind, placebo-controlled trial. *Int Neurourol J* 2018;22:30–40.
- [60] Hajebrahimi S, Asrbadr YA, Azaripour A, Sadeghi-Bazargani H. Effect of tamsulosin versus prazosin on clinical and urodynamic parameters in women with voiding difficulty: a randomized clinical trial. *Int J Gen Med* 2011;4:35–9.
- [61] Xu D, Qu C, Meng H, et al. Dysfunctional voiding confirmed by transdermal perineal electromyography, and its effective treatment with baclofen in women with lower urinary tract symptoms: a randomized double-blind placebo-controlled crossover trial. *BJU Int* 2007;100:588–92.
- [62] Datta SN, Kavia RBC, Gonzales G, Fowler CJ. Results of double-blind placebo-controlled crossover study of sildenafil citrate (Viagra) in women suffering from obstructed voiding or retention associated with the primary disorder of sphincter relaxation (Fowler's syndrome). *Eur Urol* 2007;51:487–9.
- [63] Rosario DJ, Woo HH, Parkhouse H, Chapple CR. Effects of intravenous thyrotropin-releasing hormone on urethral closure pressure in females with voiding dysfunction. *Eur Urol* 1995;28:64–7.
- [64] Kao YL, Huang KH, Kuo HC, Ou YC. The therapeutic effects and pathophysiology of botulinum toxin a on voiding dysfunction due to urethral sphincter dysfunction. *Toxins* 2019;11:728. <https://doi.org/10.3390/toxins11120728>.
- [65] Jiang Y-H, Wang C-C, Kuo H-C. OnabotulinumtoxinA urethral sphincter injection as treatment for non-neurogenic voiding dysfunction – a randomized, double-blind, placebo-controlled study. *Sci Rep* 2016;6:38905.
- [66] Pradhan AA. Botulinum toxin: an emerging therapy in female bladder outlet obstruction. *Indian J Urol* 2009;25:318–20.
- [67] Datta SN, Chaliha C, Singh A, et al. Sacral neurostimulation for urinary retention: 10-year experience from one UK centre. *BJU Int* 2008;101:192–6. <https://doi.org/10.1111/j.1464-410X.2007.07282.x>.
- [68] Peeters K, Sahai A, De Ridder D, Van Der Aa F. Long-term follow-up of sacral neuromodulation for lower urinary tract dysfunction. *BJU Int* 2014;113:789–94.
- [69] Espuña Pons M, Cassadó J, Díez Itza I, Valero Fernández EM. Post-void residual and voiding dysfunction symptoms in women with pelvic organ prolapse before and after vaginal surgery. A multicenter cohort study. *Actas Urol Esp* 2021;45:57–63.
- [70] Togo M, Kitta T, Kanno Y, et al. Lower urinary tract function improves after laparoscopic sacrocolpopexy for elderly patients with pelvic organ prolapse. *Low Urin Tract Symptoms* 2020;12:260–5.
- [71] Heidari F, Abbaszadeh S, Ghadian A, Tehrani KF. On demand urethral dilatation versus intermittent urethral dilatation: results and complications in women with urethral stricture. *Nephrourol Mon* 2014;6:e15212.
- [72] Popat S, Zimmern PE. Long-term management of luminal urethral stricture in women. *Int Urogynecol J* 2016;27:1735–41.
- [73] Grivas N, Tsimaris I, Makatsori A, Hastazeris K, Kafarakis V, Stavropoulos NE. The effectiveness of Otis urethrotomy combined with six weeks urethral dilations until 40 Fr in the treatment of bladder outlet obstruction in women: a prospective study. *Urol J* 2014;10:1063–6.
- [74] Fu Q, Xu Y-M. Transurethral incision of the bladder neck using KTP in the treatment of bladder neck obstruction in women. *Urol Int* 2009;82:61–4.
- [75] Jin X, Qu H, Liu H, Li B, Wang J, Zhang Y. Modified transurethral incision for primary bladder neck obstruction in women: a method to improve voiding function without urinary incontinence. *Urology* 2012;79:310–3.
- [76] Shen W, Ji H, Yang C, et al. Controlled transurethral resection and incision of the bladder neck to treat female primary bladder neck obstruction: description of a novel surgical procedure. *Int J Urol* 2016;23:491–5.
- [77] Zhang P, Wu Z-J, Xu L, Yang Y, Zhang N, Zhang X-D. Bladder neck incision for female bladder neck obstruction: long-term outcomes. *Urology* 2014;83:762–6.
- [78] Choudhury A. Incisional treatment of obstruction of the female bladder neck. *Ann R Coll Surg Engl* 1978;60:404–7.
- [79] Hampson LA, Myers JB, Vanni AJ, et al. Dorsal buccal graft urethroplasty in female urethral stricture disease: a multi-center experience. *Transl Androl Urol* 2019;8(Suppl 1):S6–S12.
- [80] Gomez RG, Segura FJ, Saavedra A, Campos RA. Female urethral reconstruction: dorsal buccal mucosa graft onlay. *World J Urol* 2020;38:3047–54.
- [81] Tao T, Xu Q, Hu Q, et al. Novel surgical technique for female distal urethral stricture disease: an evaluation of efficacy and safety compared with urethral dilatation. *Int J Clin Exp Med* 2018;11:12002–7.
- [82] McCreery R, Appell R. Transvaginal urethrolysis for obstruction after antiincontinence surgery. *Int Urogynecol J Pelvic Floor Dysfunct* 2007;18:627–33.

- [83] Mouracade P, Lang H, Jacqmin D, Saussine C. Transvaginal tape lysis for urinary obstruction after suburethral tape placement. When to do an immediate replacement? *Int Urogynecol J Pelvic Floor Dysfunct* 2008;19:1271–4.
- [84] Giannis G, Bousouni E, Mueller MD, et al. Can urethrolisis resolve outlet obstruction related symptoms after Burch colposuspension for stress urinary incontinence? *Eur J Obstet Gynecol Reprod Biol* 2015;195:103–7.
- [85] Leng WW, Davies BJ, Tarin T, Sweeney DD, Chancellor MB. Delayed treatment of bladder outlet obstruction after sling surgery: association with irreversible bladder dysfunction. *J Urol* 2004;172:1379–81.
- [86] Agnew G, Dwyer PL, Rosamilia A, Edwards G, Lee JK. Functional outcomes for surgical revision of synthetic slings performed for voiding dysfunction: a retrospective study. *Eur J Obstet Gynecol Reprod Biol* 2012;163:113–6.
- [87] Van den Broeck T, De Ridder D, Van der Aa F. The value of surgical release after obstructive anti-incontinence surgery: An aid for clinical decision making. *Neurourol Urodyn* 2015;34:736–40.
- [88] Hashim H, Blaker MH, Drake MJ, et al. International Continence Society (ICS) report on the terminology for nocturia and nocturnal lower urinary tract function. *Neurourol Urodyn* 2019;38:499–508. <https://doi.org/10.1002/nau.23917>.
- [89] Weiss JP, Everaert K. Management of nocturia and nocturnal polyuria. *Urology* 2019;133S:24–33.
- [90] Van Kerrebroeck P, Abrams P, Chaikin D, et al. The standardisation of terminology in nocturia: report from the Standardisation Subcommittee of the International Continence Society. *Neurourol Urodyn* 2002;21:179–83.
- [91] Bedretinova D, Ambühl D, Omar MI, et al. What is the most effective treatment for nocturia or nocturnal incontinence in adult women? *Eur Urol Focus* 2021;7:453–63.
- [92] Lo SK, Naidu J, Cao Y. Additive effect of interferential therapy over pelvic floor exercise alone in the treatment of female urinary stress and urge incontinence: a randomized controlled trial. *Hong Kong Physiother J* 2003;21:37–42. [https://doi.org/10.1016/S1013-7025\(09\)70038-7](https://doi.org/10.1016/S1013-7025(09)70038-7).
- [93] Aslan E, Komurcu N, Beji NK, Yalcin O. Bladder training and Kegel exercises for women with urinary complaints living in a rest home. *Gerontology* 2008;54:224–31.
- [94] Johnson 2nd TM, Burgio KL, Redden DT, Wright KC, Goode PS. Effects of behavioral and drug therapy on nocturia in older incontinent women. *J Am Geriatr Soc* 2005;53:846–50.
- [95] Fitzgerald MP, Lemack G, Wheeler T, Litman HJ. Nocturia, nocturnal incontinence prevalence, and response to anticholinergic and behavioral therapy. *Int Urogynecol J Pelvic Floor Dysfunct* 2008;19:1545–50.
- [96] Wang T, Huang W, Zong H, Zhang Y. The efficacy of continuous positive airway pressure therapy on nocturia in patients with obstructive sleep apnea: a systematic review and meta-analysis. *Int Neurourol J* 2015;19:178–84.
- [97] Yamaguchi O, Nishizawa O, Juul KV, Nørgaard JP. Gender difference in efficacy and dose response in Japanese patients with nocturia treated with four different doses of desmopressin orally disintegrating tablet in a randomized, placebo-controlled trial. *BJU Int* 2013;111:474–84.
- [98] Lose G, Lalos O, Freeman RM, van Kerrebroeck P. Efficacy of desmopressin (Minirin) in the treatment of nocturia: a double-blind placebo-controlled study in women. *Am J Obstet Gynecol* 2003;189:1106–13.
- [99] Hilton P, Stanton SL. The use of desmopressin (DDAVP) in nocturnal urinary frequency in the female. *Br J Urol* 1982;54:252–5.
- [100] Rembratt A, Riis A, Nørgaard JP. Desmopressin treatment in nocturia; an analysis of risk factors for hyponatremia. *Neurourol Urodyn* 2006;25:105–9.
- [101] Rovner ES, Raymond K, Andruczyk E, Juul KV. Low-dose desmopressin and tolterodine combination therapy for treating nocturia in women with overactive bladder: a double-blind, randomized, controlled study. *Low Urin Tract Symptoms* 2018;10:221–30. <https://doi.org/10.1111/luts.12169>.
- [102] Lose G, Englev E. Oestradiol-releasing vaginal ring versus oestriol vaginal pessaries in the treatment of bothersome lower urinary tract symptoms. *Br J Obstet Gynaecol* 2000;107:1029–34.
- [103] Reynard JM, Cannon A, Yang Q, Abrams P. A novel therapy for nocturnal polyuria: a double-blind randomized trial of frusemide against placebo. *Br J Urol* 1998;81:215–8. <https://doi.org/10.1046/j.1464-410x.1998.00511.x>.